



Day case parathyroidectomy: is this the right way for the patients?

Rocco Rago¹, Francesco Forfori², Gianluca Frustaci³, Roberta Monzani⁴, Simone Paracchini², Francesca Franceschini², Filomena Cetani⁵, Gabriele Materazzi³

¹Centro Multidisciplinare Ricovero Breve, ²UO Anestesia e Rianimazione Interdipartimentale, ³UO Endocrinochirurgia, Università di Pisa, Pisa, Italy; ⁴UO Anestesia e Day Hospital Chirurgico, Istituto Clinico Humanitas, Rozzano, Milano, Italy; ⁵UO Endocrinologia II, Università di Pisa, Pisa, Italy

Contributions: (I) Conception and design: R Rago; (II) Administrative support: None; (III) Provision of study materials or patients: All authors; (IV) Collection and assembly of data: All authors; (V) Data analysis and interpretation: All authors; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Rocco Rago. Centro Multidisciplinare Ricovero Breve, Università di Pisa, Pisa, Italy. Email: r.rago@ao-pisa.toscana.it.

Background: Minimally-invasive video-assisted parathyroidectomy (MIVAP) can be considered as the primary treatment of choice for single parathyroid adenoma. Often, this technique is performed in a day surgery setting and is associated with regional anaesthesia (RA). Many studies have already reported the feasibility and safety of MIVAP in day surgery. Here our focus has been on the patient's personal experience with these procedures through an assessment of their recovery at home.

Methods: We conducted a prospective observational study in the University Hospital of Pisa Day Surgery Unit. Forty-eight patients were enrolled and divided by personal choice of anaesthesia technique: a regional anaesthesia group (RAg) and general anaesthesia group (GAg). Data were extracted from the medical records and three questionnaires: the first was self-compiled at discharge (Q1), while the second (Q2) and the third (Q3) were administered as telephone surveys.

Results: None of the patients in RAg reported pain longer than 1 day after discharge, whereas 15% of patients in GAg reported pain relief the third day after discharge ($P=0.0065$). Discharge in RAg was within 3 hours in 12.5% of patients, within 4 hours in 78.1%, and within 5 hours in 9.4%. Discharge in GAg was within 5 hours in 53.8% and in more than 5 hours in 46.1% ($P=0.0027$).

Conclusions: Patients highly appreciated day-case parathyroidectomy. Furthermore, the association of RA with MIVAP leads to better results than those of general anaesthesia (GA) and MIVAP. Finally, we point out that it is fundamental that the physicians pay attention to what the patients consider important for them; that is, the personal meaning of the hospitalization.

Keywords: Parathyroid gland; day surgery; patient satisfaction

Submitted Mar 31, 2019. Accepted for publication Aug 21, 2019.

doi: 10.21037/gs.2019.09.02

View this article at: <http://dx.doi.org/10.21037/gs.2019.09.02>

Introduction

As demonstrated in the literature (1-6), minimally-invasive video-assisted parathyroidectomy (MIVAP) can be considered the treatment of choice for single parathyroid adenoma. In the major hospitals of the United States and in a few other countries, this technique is performed in a day surgery (DS) setting (7-9), even on adolescents (10), with high rates of success and a low incidence of complications.

MIVAP is often associated with regional anaesthesia (RA) (11-13), because of the advantages compared with general anaesthesia (GA), including: continuous and direct monitoring of the recurrent laryngeal nerve, better postoperative pain control, and lower risk of postoperative nausea and vomiting (PONV). Owing to these advantages, elderly and/or high-risk patients can be more easily included as candidates to the operation (14).

Table 1 Questionnaires used respectively at discharge, the next day and after 4 weeks

Questionnaire 1
Clinical care
Clinical information
Pain perception
ADLm
The time or missing discharge
Questionnaire 2
Pain perception
IADLm
Minor complications at home
Need for advice or assistance
Unplanned admissions
Questionnaire 3
Pain relief
IADLm and resuming job
Minor complications at home
Need for advice or assistance
Unplanned admissions

ADLm, Activities of Daily Living modified scale; IADLm, Instrumental Activities of Daily Living modified scale.

MIVAP with RA was introduced in Italy in 2002 (11) but has been typically performed as an inpatient procedure. We believe it is time to perform this operation as a day case in our country as well.

Aim of study

Many studies have already reported the feasibility and safety of MIVAP in DS. Therefore, here our focus has been on the patient's personal experience with these procedures through an assessment of their recovery at home. The first set of outcome measures of interest included the quality of care provided and to the well-being of the patient: information received, pain, PONV, and resumption of activities of daily living and work. The second set of outcome measures were related to the patient's safety: complications at home, the need for medical advice and assistance, and unscheduled admissions.

Methods

After Local Ethics Committee approval, we conducted a prospective observational study in the University Hospital of Pisa Day Surgery Unit. Between June 2015 and November 2017, 48 patients were enrolled according to the following inclusion criteria:

- ❖ Anaesthesiology: outpatient criteria, age between 18 and 80 years, American Society of Anaesthesiologists (ASA) Classification I to III, no psychiatric disorders, no chronic therapy with anti-inflammatory drugs, and no osteo-articular pathologies or recent traumas in the cervical region.
- ❖ Surgical: no familial forms of hyperparathyroidism, preoperative ultrasonographic localization of adenoma, no previous neck surgery or irradiation, and no suspicion of thyroiditis or malignancy.

Patients were divided by personal choice of anaesthesia technique: a regional anaesthesia group (RAG) and general anaesthesia group (GAG). As supported in other studies (15), the patient's emotional state is essential in this kind of procedure, so we, therefore, considered patient preference a better strategy than randomization.

Written informed consent to the study procedures was obtained from each patient. Data were extracted from the medical records and from three questionnaires: the first was self-compiled at discharge (Q1), while the second (Q2) and the third (Q3) were administered as a telephone survey, respectively the next day and after 4 weeks (*Table 1*). Most variables were rated as a score from 0 to 4 points. Some questions could be answered "yes" or "no", and for others, the reference times (e.g., time of discharge) were recorded. Finally, the pain was assessed with the Numerical Rating Scale (NRS) as: 9 to 10, severe; 7 to 8, poorly controlled; 4 to 6, controlled; 1 to 3, well-controlled; 0 absent.

Anaesthesia technique

In RAG, a bilateral deep cervical block and an infiltration along the incision line was performed with a total of 30 mL of a mixture of lidocaine 2% (10 mL), ropivacaine 1% (10 mL), and levobupivacaine 0.75% (10 mL). A bilateral block was preferred to improve pain control and to permit exploration of the other side, if necessary, since this does not imply a greater risk (14,15). Remifentanyl was used to provide mild intravenous sedation (0.05–0.10 µg/kg/min). During the operation, the patient remained in a

spontaneous breath, conscious, and contactable.

In GAg, total intravenous anaesthesia was performed. GA was induced with propofol (2 mg/kg) and remifentanyl (0.50 µg/kg/min), then maintained with the same drugs, respectively, at 4 mg/kg/h and 0.30 µg/kg/min. A laryngeal mask (LM) was used to secure the airways and ventilate the patient. At induction, ondansetron (4 mg) and dexamethasone (8 mg) were administered for anti-emetic purposes. Both groups received ketorolac (30 mg) and acetaminophen (1 g) for analgesia. The same drugs were administered in the ward 4 hours after the operation. Patients were discharged at home when they reached a modified Post-Anaesthetic Discharge Scoring System (PADSS) score of at least 9 points.

Surgical technique

All patients in both groups undergone MIVAP procedure. A 1.5-cm transverse minicervicotomy is performed 2 cm above the sternal notch, no skin flap is required, the anterior cervical fascia is opened for 2–3 cm on the midline, and the thyroid gland is exposed. Once reached the thyroid gland, infrahyoid muscles homolateral to the suspicious parathyroid gland is dissected from the thyroid lobe; the thyroid lobe is then retracted medially, and the strap muscles are retracted laterally. A 5-mm, 30-degree endoscope, is inserted through the cervicectomy to explore both the parathyroid of the same side. Once enlarged parathyroid is found, a blunt dissection plus a single ligation are performed for its removal. All patients of the study undergone parathyroid hormone (PTH) intraoperative assay by means of a preoperative PTH level (time 0') and an intraoperative PTH level 5 minutes later than the parathyroid gland removal (time +5') in order to prove a 50% PTH level decrease. If the PTH 50% level dropdown is not obtained, the contralateral side exploration is possible with the same surgical steps.

Statistic analysis

This study was conducted on a per-protocol basis. A descriptive analysis of data was performed using means and standard deviations for continuous variables. Statistical significance was evaluated using Fisher two-tailed tests, with an α error of 0.05. Qualitative and ordinal variables are expressed as percentages and measures of frequency.

Results

Of the total patients, 44 were women (91.7%), and 4 were men (8.3%). Mean age was 56.65 ± 12.09 years (range, 30–79 years), with RAg aged 55.47 ± 11.75 years (range, 30–79 years) and GAg aged 59.5 ± 12.86 (range, 40–77 years). There were 34 patients (70.8%) in the RAg and 14 (29.2%) the GAg. Two patients were transferred to the Surgical Unit for an overnight stay, and another patient was transferred to the Cardiology Unit and discharged after 48 hours. These three patients were excluded from the study. The remaining 45 patients (32 in RAg and 13 in GAg) were discharged the same day. No patient reported PONV. No cases of regional block failure occurred and none of the surgical cases needed to be converted from RA to GA.

Surgical results

All patients in both groups underwent MIVAP; no bilateral exploration was needed. In the RA, there were 15 right side parathyroid adenomas: 7 superior right and 8 inferior right and 19 left side adenomas (8 superior left and 11 inferior left). In the GAg, there were 7 right side parathyroid adenomas: 3 superior and 4 inferior and 7 were on the left side (2 superior and 5 inferior). The parathyroid adenoma removal was demonstrated by the 50% intraoperative PTH level dropdown in all cases. All patients underwent calcium blood assay on the first postoperative day and on the second postoperative day; in all cases, we collected normal calcium level on the first and second postoperative day.

Pain

Considering the small population, none of whom reported severe pain, we grouped the pain intensity in two categories: absent/well-controlled (NRS, 0 and 1 to 3) and controlled/poorly controlled (NRS 4 to 8). Intraoperative pain was reported as absent in 21 patients (65.7%) and well-controlled in 11 patients (34.3%) of RAg.

Pain at Q1 was better controlled in RAg than GAg (*Table 2*). In fact, 31 of 32 patients (96.9%) in RAg reported absent/well-controlled pain against 9 of 13 (69.2%) in GAg ($P=0.0061$). At Q2, all RAg patients (100%) and 11 of 13 GAg patients (84.6%) reported absent/well-controlled pain ($P=0.0085$; *Table 2*).

At Q3, we verified pain relief at home (*Table 3*). None of the patients in RAg reported pain for longer than 1 day

Table 2 Pain results at Q1 and Q2

Level of pain	Pain at discharge (Q1)				Pain at post-operative day 1 (Q2)			
	GAg		RAg		GAg		RAg	
	n	%	n	%	n	%	n	%
Severe	0	0	0	0	0	0	0	0
Poorly controlled	2	15.4	0	0	0	0	0	0
Controlled	2	15.4	1	3.1	2	15.4	0	0
Well-controlled	5	38.5	7	21.9	5	38.5	2	6.3
Absent	4	30.8	24	75.0	6	46.2	30	93.8

GAg, general anaesthesia group; RAg, regional anaesthesia group.

Table 3 Pain data regarding Q3 survey

Time of pain relief (Q3)	GAg, n (%)	RAg, n (%)
The same day of discharge	5 (38.5)	30 (93.8)
1 day	6 (46.2)	2 (6.3)
3 days	2 (15.4)	0
More than 1 week	0	0
More than 2 weeks	0	0

GAg, general anaesthesia group; RAg, regional anaesthesia group.

after discharge, whereas 15% of patients in GAg reported pain relief the third day after discharge ($P=0.0065$).

Resumption of basic activities and cognitive function in the hospital

During hospitalization, recovery of autonomy was assessed with the Activities of Daily Living modified scale (ADLm, *Table 4*). The results were 71.9% of patients in RAg ate within 2 hours after surgery and 28.1% within 3 hours, whereas 84.6% patients in GAg ate within 4 hours and 15.4% within 5 hours ($P=0.0045$; *Table 5*). Return of cognitive function was also faster in RAg: 93.8% of patients within 2 hours, 6.2% within 3 hours in RAg, and 15.4% of patients within 2 hours, 69.2% within 3 hours, and 15.4% within 4 hours in GAg ($P=0.0107$). Discharge in RAg occurred within 3 hours in 12.5% of patients, within 4 hours in 78.1%, and within 5 hours in 9.4%. Discharge in GAg occurred within 5 hours in 53.8% and more than 5 hours in 46.1% ($P=0.0027$).

Table 4 Modified Activities of Daily Living scale

Activities	Score
Dress oneself	
Takes clothes and dresses without the need of help	1
Takes clothes and dresses without help except for tying shoes	1
Needs help for taking clothes or dressing	0
Toilette	
Goes toilet cleans himself and puts clothes on without assistance	1
Needs assistance going to the bathroom, to clean himself, or to put clothes on	0
Does not go to the bathroom	0
Gait	
Moves without assistance	1
Moves if helped	0
Bedridden	0
Nutrition	
No need for assistance	1
Needs assistance only to cut meat or to butter bread	1
Needs assistance to bring the food to the mouth	0
Interaction with people or the environment	
Watches television, reads, uses telephone and talks to people by own initiatives	1
Does at least two of the aforementioned activities	1
Does none of these activities	0
Minimum score	5

Table 5 Modified Activities of Daily Living results

Time after surgery	Eats		Relationship		Discharge	
	GAg	RAg	GAg	RAg	GAg	RAg
Within 2 h	–	71.90%	15.40%	93.80%	–	–
Within 3 h	–	28.10%	69.20%	6.20%	–	12.50%
Within 4 h	84.60%	–	15.40%	–	–	78.10%
Within 5 h	15.40%	–	–	–	53.80%	9.40%
>5 h	–	–	–	–	46.20%	–

GAg, general anaesthesia group; RAg, regional anaesthesia group.

Table 6 Modified Instrumental Activities of Daily Living scale

Activities	Score
Shopping	
Takes care of shopping without assistance	1
Needs help or support for shopping	0
Is completely unable to shop	0
Cooking food	
Organizes prepares and serves food	1
Prepares food only if ingredients are provided	0
Needs food prepared and served	0
Rules the house	
Rules the house alone or requires help only for hard jobs	1
Does only daily jobs, but the level of cleanness is insufficient	0
Does not participate in house works	0
Laundering	
Does the laundry alone	1
Washes only some clothes	1
Needs clothes washed by others	0
Transportations	
Moves along with public transports or own car	1
Takes conveyance with help or takes a taxi by himself	1
Able to take transports only with the help	0
Cannot move	0
Drugs	
Takes prescribed medications without help	1
Takes prescribed medications only if doses are prepared by others	0
Cannot take medications alone	0
Minimum score	6




Figure 1 Kaplan-Meier curve for scores of the Instrumental Activities of Daily Living modified scale (IADL scale; minimum score =6).

Resumption daily activities at home and work

Self-sufficiency at home was evaluated with Instrumental Activities of Daily Living modified scale (IADLm, *Table 6*). None of the patients regained complete self-sufficiency the first day after discharge. The next day, the RAg reached score of 5 in 43.7% of patients, 4 in 46.9%, and 2 in 9.4%, and the GAg scores were 4 in 38.5% of patients, 2 in 46.1%, and 0 in 15.4% ($P=0.025$). We did not find statistically significant differences between the two groups at Q3. However, we point out that 25% of the RAg patients were autonomous at the end of the second day after discharge, reaching 100% within a week. Conversely, in the GAg, 53.8% began to be self-sufficient after 3 days and 100% reached self-sufficiency after 1 week (*Figure 1*).

Resumption of work (*Figure 2*), excluding 31.1% of retired patients, occurred in 30.5% of RAg patients within 1 week, in 47.8% after 1 week, in 13.0% within 2 weeks, and in 8.7% within 3 weeks. Resumption for the GAg occurred in 25% after 1 week, in 50% within 2 weeks, and




Figure 2 Kaplan-Meier curve for the resumption of work activities after discharge (retired patients were excluded).

in 25% within 3 weeks. The difference between the two groups was significant ($P=0.02$).

Complications at home

Complications at home and the need for medical care (home visit, telephone advice, visiting a general practitioner), as well as access to the emergency department or unplanned hospitalization, were the same and not statistically significant in the two groups (2 patients in each group reported mild infection at the wound, and 2 reported gastritis). Eventually, symptomatic hypocalcaemia developed at home in RAg patient despite the consumption of calcium prescribed at discharge. No complications were directly dependent on the anaesthetic choice.

Clinical care satisfaction

Clinical care was reported as excellent in 71.9% of RAg patients and good in 28.1% at Q1, whereas GAg patients rated it excellent in 23.1% and good in 76.9%. Statistics were comparable between the two groups. At Q2 in the RAg, satisfaction was reported as excellent in 96.8% and good in 3.2%, and in GAg excellent in 69.2%, good in 23.1%, and enough in 7.7% ($P=0.02$).

At Q3, we asked patients whether they would return to our department in case of necessity or whether they would advise MIVAP to others. The response was an absolute yes in 96.9% of RAg patients and yes in 3.1%, whereas the response in GAg patients was an absolute yes in 61.5%, yes in 23.1%, and probably in 15.4% ($P=0.018$).

Discussion

The literature agrees on the feasibility and safety of MIVAP in DS performed in RA, considering it superior also to open techniques in some respects. Furthermore, reduced health care expenditure is another reason to prefer it. But what is the patient's perception? Beyond the clinical outcome, is this the right way for the patient? Everything that happens in a hospital has a personal meaning for the patient, compared with the biological significance attributed by physicians (16). We, therefore, focused on the personal sense of hospitalization: perception of the quality of care and well-being (level of treatments and information received, pain, PONV, resumption of ADL), as well as patient safety (need for counselling or assistance, complications, unplanned hospitalization).

Pain

Intraoperative pain control in RAg was very satisfactory, confirming the efficacy of the technique: 65.7% of patients reported the absence of pain, and 34.3% indicated pain as controlled.

According to literature (3,11), postoperative pain assessment was lower in the RAg ($P=0.0061$) than in the GAg, despite the use of LM rather than an endotracheal tube. This choice permitted a decreased risk of severe coughing at the time of extubating and then of bleeding. The subsequent home survey confirmed the best results in the RAg for pain perception (Q2: $P=0.0085$) and on faster resolve (Q3: $P=0.0065$). Our data, therefore, confirm the efficacy of cervical block and intravenous sedation in this kind of operation, reducing pain, PONV, and complications that delay patient well-being and discharge.

Self-sufficiency at hospital and home, work resumption

To evaluate the resumption of ADL and human relations, we modified the Katz scale and Lawton-Brody scale (17-20). They are usually used to assess motor and psychologic skills in people with reduced ability and in the elderly, as well as in occupational therapy to check the patient's progress in recovering from illness or injury. Changing some voices and deleting others, we adapted the scales (ADLm, IADLm) to the study context. Values obtained and the time taken to reach the minimum score of (5 for

ADLm and 6 for IADLm) confirm the superiority of RA compared with GA. Thanks to the reduced psychophysical effect of local anaesthesia, the RAg patients ate earlier ($P=0.0045$), regained cognitive function and resumed human relationships more quickly ($P=0.0107$), and were discharged in a shorter time ($P=0.0027$) compared with the GAg patients, with better results than reported by other studies (21). Despite the use of the LM and short-term drugs (propofol and remifentanyl), the prevalence of RA over GA was confirmed for other ADLm and IADLm entries, although they were not statistically significant. All patients returned to their work within 4 weeks, although RAg patients did so more rapidly in the first days after discharge ($P=0.02$). Eventually, we believe our study shows that RA guarantees greater well-being than GA.

Missing discharge

Three patients were admitted as inpatients for prolonged observation and were dropped from the study. One patient showed signs of heart failure in the afternoon, was transferred to the Cardiology Unit, and was discharged after 48 hours with a diagnosis of Takotsubo disease. The other two patients underwent the operation with an unclear diagnosis: parathyroid was found deep in the mediastinum in one patient, and thyroidectomy was associated with the other patient. Because the operations in both patients extended beyond 2 hours and the bleeding risk, they were transferred to the Surgery Unit and discharged the following morning. Considering these instances were not directly related to procedures, we can conclude that the care pathway is feasible and safe.

Complications at home

We considered complications at home not dangerous for the patient's safety, but rather that could influence the quality of recovery, including mild respiratory disorders (e.g., hoarseness), gastritis, PONV, local edema, or infection. The only true complication recorded was symptomatic hypocalcaemia in one patient, despite calcium integration from the day after the operation (22,23). The patient went to the Emergency Department and was discharged in a few hours. No other complication occurred, and the care at home was limited to telephone counselling with surgeons or one or more accesses to the general practitioner. This confirms what has been highlighted in the literature on the safety of this procedure. No unplanned readmission to the

hospital was necessary.

Clinical care satisfaction

The collected data at the Q3 survey showed the satisfaction of all patients for the clinical care received. Once again, however, the RAg patients were more satisfied than GAg patients ($P=0.02$), for pain relief and short duration, as well as for the feeling of well-being and quick recovery of autonomy after surgery.

Finally, all RAg patients would have made the same care path, whereas a small number of GAg patients (15.38%) appeared doubtful because the pain control was not perfect, and the pain lasted for some days.

Conclusions

Although the study was conducted on a small sample, the data showed that patients highly appreciated day-case parathyroidectomy. This is due to the quality of care provided and to well-being after the operation. Furthermore, the study demonstrates, at the parity of security, that the association of RA with MIVAP leads to better results than the association of GA and MIVAP.

Finally, we point out that, beyond the clinical ability and knowledge, it is fundamental that the physicians pay attention to what the patients consider important for them; that is, the personal meaning of the hospitalization.

Acknowledgments

None.

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. This study was approved by the local ethics committee (No. 12450) and written informed consent was obtained from all patients.

References

1. Bakkar S, Matteucci V, Corsini C, et al. Less is more:

- time to expand the indications for minimally invasive video-assisted parathyroidectomy. *J Endocrinol Invest* 2017;40:979-83.
2. Melfa GI, Raspanti C, Attard M, et al. Comparison of minimally invasive parathyroidectomy under local anaesthesia and minimally invasive video-assisted parathyroidectomy for primary hyperparathyroidism: a cost analysis. *G Chir* 2016;37:61-7.
 3. Rio PD, Vicente D, Maestroni U, et al. A comparison of minimally invasive video-assisted parathyroidectomy and traditional parathyroidectomy for parathyroid adenoma. *J Cancer* 2013;4:458-63.
 4. Garimella V, Yeluri S, Alabi A, et al. Minimally invasive video-assisted parathyroidectomy is a safe procedure to treat primary hyperparathyroidism. *Surgeon* 2012;10:202-5.
 5. Gracie D, Hussain SS. Use of minimally invasive parathyroidectomy techniques in sporadic primary hyperparathyroidism: systematic review. *J Laryngol Otol* 2012;126:221-7.
 6. Augustine MM, Bravo PE, Zeiger MA. Surgical treatment of primary hyperparathyroidism. *Endocr Pract* 2011;17:75-82.
 7. Stack BC Jr, Moore E, Spencer H, et al. Outpatient thyroid surgery data from the University Health System (UHC) Consortium. *Otolaryngol Head Neck Surg* 2013;148:740-5.
 8. Dulfer RR, van Ginhoven TM, Geilvoet W, et al. Operative Treatment of Primary Hyperparathyroidism in Daycare Surgery. *Scand J Surg* 2015;104:196-9.
 9. Rajeev P, Sutaria R, Ezzat T, et al. Changing trends in thyroid and parathyroid surgery over the decade: is same-day discharge feasible in the United Kingdom? *World J Surg* 2014;38:2825-30.
 10. Langusch CC, Norlen O, Titmuss A, et al. Focused image-guided parathyroidectomy in the current management of primary hyperparathyroidism. *Arch Dis Child* 2015;100:924-7.
 11. Miccoli P, Barellini L, Monchik JM, et al. Randomized clinical trial comparing regional and general anaesthesia in minimally invasive video-assisted parathyroidectomy. *Br J Surg* 2005;92:814-8.
 12. Cohen MS, Finkelstein SE, Brunt LM, et al. Outpatient minimally invasive parathyroidectomy using local/regional anesthesia: a safe and effective operative approach for selected patients. *Surgery* 2005;138:681-7.
 13. Materazzi G, Rago R, Berti PF, et al. Paratiroidectomia mini-invasiva con blocco del plesso cervicale. In: *Trattato Italiano di Endocrinochirurgia*. vol. II, Publisher: L. Rosato, Editors: GS, 2008:541-3.
 14. Fang WL, Tseng LM, Chen JY, et al. The management of high-risk patients with primary hyperparathyroidism - minimally invasive parathyroidectomy vs. medical treatment. *Clin Endocrinol (Oxf)* 2008;68:520-8.
 15. Sanjay P, Marioud A, Woodward A. Anaesthetic preference and outcomes for elective inguinal hernia repair: a comparative analysis of public and private hospitals. *Hernia* 2013;17:745-8.
 16. Kiernan CM, Schlegel C, Isom C, et al. Ambulatory bilateral neck exploration for primary hyperparathyroidism: is it safe? *Am J Surg* 2016;212:722-7.
 17. Katz S, Ford AB, Moskowitz RW, et al. Studies of illness in the aged. the index of ADL: a standardized measure of biological and psychosocial function. *JAMA* 1963;185:914-9.
 18. Katz S, Downs TD, Cash HR, et al. Progress in the development of the index of ADL. *Gerontologist* 1970;10:20-30.
 19. Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist* 1969;9:179-86.
 20. Kopp VJ, Shafer A. Anesthesiologists and perioperative Communication. *Anesthesiology* 2000;93:548-55.
 21. Meltzer C, Klau M, Gurushanthaiah D, et al. Safety of Outpatient Thyroid and Parathyroid Surgery: A Propensity Score-Matched Study. *Otolaryngol Head Neck Surg* 2016;154:789-96.
 22. Udelsman R. Six hundred fifty-six consecutive explorations for primary hyperparathyroidism. *Ann Surg* 2002;235:665-70; discussion 670-2.
 23. Gurnell EM, Thomas SK, McFarlane I, et al. Focused parathyroid surgery with intraoperative parathyroid hormone measurement as a day-case procedure. *Br J Surg* 2004;91:78-82.

Cite this article as: Rago R, Forfori F, Frustaci G, Monzani R, Paracchini S, Franceschini F, Cetani F, Materazzi G. Day case parathyroidectomy: is this the right way for the patients? *Gland Surg* 2020;9(Suppl 1):S6-S13. doi: 10.21037/gs.2019.09.02