

regions of white matter microstructural significant changes were detected.

Results: The LPE group had higher mean FA and mean AD in widespread regions. Moreover, the depression and anxiety is negatively correlated with the mean FA of the right posterior thalamic radiation (PTR).

Conclusions: Our study will be helpful for improving our understanding of the mechanism of LPE. Larger and possibly longitudinal studies will be required to confirm these findings and to better specify the link between structural abnormalities and functional data.

Keywords: Premature ejaculation (PE); tract-based spatial statistical (TBSS)

doi: 10.21037/tau.2017.s083

Cite this abstract as: Gao M, Yuan JL. Abnormal white matter microstructural in lifelong premature ejaculation identified by tract-based spatial statistical analysis. *Transl Androl Urol* 2017;6(Suppl 3):AB083. doi: 10.21037/tau.2017.s083

AB084. Sexual anxiety correlate to visual attention bias in sex situation: an eye-tracking study

Yung-Jui Huang, Yen-Chin Lin, Yuan-Hsiang Chu

The Institute of Human Sexuality, Shu-Te University Affiliations, Gaoxiong 807, China

Background: Eye-tracking may aid in understanding the cognitive processes involved in processing sexually appetitive material. The aim of this study was to examine the correlation of visual attention to sexual stimuli and personal sexual anxiety (SA).

Methods: Nineteen heterosexual adults completed Sexual Anxiety Scale questionnaire and an eye-tracking task used free-viewing paradigm, participants were presented with a couple array composed of sexual interaction for 30 trials, recorded throughout the trial to assess the attention processing.

Results: Higher level of SA for “Sexual Communication”

had initial orienting bias and higher fixation probability for faces in hardcore stimuli, but did not exhibit in soft-core and neutral stimuli. No correlation was found for “Solitary and Impersonal Sexual Expression” and “Exposure to Information”.

Conclusions: The present findings suggest that, sexual communicate anxious individuals are characterized by enhanced engagement with sexual threat at an early stage of processing and difficulty in disengaging from emotional clues once their initial attention is located on it.

Keywords: Sexual anxiety (SA); eye tracking; attention

doi: 10.21037/tau.2017.s084

Cite this abstract as: Huang YJ, Lin YC, Chu YH. Sexual anxiety correlate to visual attention bias in sex situation: an eye-tracking study. *Transl Androl Urol* 2017;6(Suppl 3):AB084. doi: 10.21037/tau.2017.s084

AB085. A simple novel surgical technique for penile elongation; compensation for tunical plication in Peyronie's disease

Sun Tae Ahn, Du Geon Moon

Korea University College of Medicine, Seoul 08308, South Korea

Background: The major concern with applying tunical plication procedures for Peyronie curvature is the associated with potential loss of penile length. We describe a novel technique for penile elongation by simple scrotal septum detachment from penile base to compensate penile length loss during penile plication in Peyronie's disease and evaluated the efficacy and safety of our technique.

Methods: From January 2014 to May 2017, we evaluated 16 patients (24–63 years old) with Peyronie's disease underwent penile plication with penile elongation via our novel technique. The penile elongation was performed by release and detachment of scrotal septum from penile base until scrotal fat tissue identified. We assessed the stretched penile length (SPL) preoperatively and 4 weeks