Erratum to investigation of multichannel phased array performance for fetal MR imaging on 1.5T clinical MR system

Xiaoliang Zhang^{1,2}

¹Department of Radiology and Biomedical Imaging, University of California San Francisco, USA; ²UC Berkeley/UCSF Joint Graduate Group in Bioengineering, San Francisco, CA, USA

Corresponding to: Xiaoliang Zhang, Ph.D. Byers Hall, Room 102D, 1700 4th Street, San Francisco, CA 94158, USA. Email: xiaoliang.zhang@ucsf.edu.



Submitted Feb 08, 2013. Accepted for publication Mar 11, 2013. doi: 10.3978/j.issn.2223-4292.2013.03.02 Scan to your mobile device or view this article at: http://www.amepc.org/qims/article/view/1668/2555

Erratum to: Quant Imaging Med Surg 2011;1:24-30

In the published article (1), the wrong B_1 maps were mistakenly inserted into *Figure 3*. The correct *Figure 3*, showing B_1 distributions in the sagittal plane for both the 32 channel fetal array and the 8 channel commercial torso array, should be as follow:



Figure 3 B_1 map of (A) 32 channel fetal array and (B) 8 channel torso array in the central sagittal plane of the phantom calculated by XFDTD. The numbers in the boxes indicated the mean B_1 (10⁻⁸ T) in the 3 cm × 3 cm region

References

 Li Y, Pang Y, Vigneron D, et al. Investigation of multichannel phased array performance for fetal MR imaging on 1.5T clinical MR system. Quant Imaging Med Surg 2011;1:24-30. **Cite this article as:** Zhang X. Erratum to investigation of multichannel phased array performance for fetal MR imaging on 1.5T clinical MR system. Quant Imaging Med Surg 2013;3(2):129. doi: 10.3978/j.issn.2223-4292.2013.03.02