



# Reducing conversion in minimally invasive distal pancreatectomy: patient selection, training, and robotics

Camille Stewart, Yuman Fong

Department of Surgery, City of Hope National Medical Center, Duarte, CA, USA

Correspondence to: Yuman Fong, MD, FACS. Department of Surgery, City of Hope National Medical Center, 1500 E Duarte Rd., Duarte, CA 91010, USA. Email: yfong@coh.org.

Comment on: Casadei R, Ricci C, Pacilio CA, *et al.* Laparoscopic distal pancreatectomy: which factors are related to open conversion? Lessons learned from 68 consecutive procedures in a high-volume pancreatic center. *Surg Endosc* 2018;32:3839-45.

Received: 17 February 2019; Accepted: 28 February 2019; Published: 19 March 2019.

doi: 10.21037/ls.2019.02.05

View this article at: <http://dx.doi.org/10.21037/ls.2019.02.05>

Minimally invasive surgery (MIS) has become the preferred approach to distal pancreatectomy at many high-volume institutions, but only accounts for 15% of distal pancreatectomies performed in the United States (1,2). This approach is used for a variety of histologies, including malignancy (3-5). As with all minimally invasive procedures, there is a risk of conversion to an open procedure, which rises with the complexity of the procedure. The conversion rate for MIS distal pancreatectomy varies in the literature from 3–31% for laparoscopic procedures (6).

In a recent paper, “*Laparoscopic distal pancreatectomy: which factors are related to open conversion? Lessons learned from 68 consecutive procedures in a high-volume pancreas center*”, Casadei and his colleagues describe their outcomes for distal pancreatectomies from 2004–2016 (7). Of note, a laparoscopic approach was attempted in 43% of all distal pancreatectomies at their institution during the study period. They reported a 19% conversion to open rate, which is well within the previously reported range, as were their other complication rates. Regression analysis revealed that the only factor associated with conversion was pancreatic transection to the right of the portal vein. In fact, odds of conversion to open were nearly 10 times higher in this group. There was also a trend towards higher conversion rates with suspicion for malignancy. More difficult cases are not always attempted in a MIS fashion due to risk of conversion. Subtotal distal pancreatectomy and distal pancreatectomy for malignancy are typically considered more difficult/complex procedures, so these risk factors for conversion to open are logical. Based on these results, one might seriously question attempting a

laparoscopic distal pancreatectomy for cancer when need for transection to the right of the portal vein (subtotal distal pancreatectomy) is anticipated. When an MIS approach is chosen for these procedures, the surgeon should counsel the patient of the higher likelihood of conversion to open surgery.

In this paper, the authors question if a robotic approach could potentially overcome the limitations of the laparoscopic equipment during subtotal distal pancreatectomy. The vast majority of MIS distal pancreatectomies in the United States are performed laparoscopically (8). Comparisons of robotic and laparoscopic distal pancreatectomy show outcomes are generally similar, including length of hospital stay, 30-day readmission rate, and overall survival (8). The conversion to open rate, however, tends to be higher for laparoscopic compared to robotic procedures (8-10). The conversion to open rate for robotic distal pancreatectomy ranges from 0–38% (6), but averages 7–8% (9,10). The higher conversion rate for the laparoscopic approach has been found both for malignancy, and in cohorts with mixed indications for surgery. Tumor proximity to major vessels is the most common reason for conversion to open surgery, as would be the case during a subtotal distal pancreatectomy (11). This particular reason for conversion tends to be more common for laparoscopic *vs.* robotic cases (29% *vs.* 14%) (11).

Casadei *et al.* imply that conversion to open is negative, but do not directly compare the post-operative outcomes of their patients who underwent an MIS completed distal pancreatectomy to those who were converted to open (7). A study of the American College of Surgeons-National Surgical Quality Improvement Program identified

1,200 patients who underwent MIS distal pancreatectomy in 2014–2015 (12). In the unadjusted analysis, the patients who underwent conversion to open surgery experienced higher overall morbidity, including serious morbidity, and mortality compared to those who had the operation completed in an MIS fashion. After propensity score matching, risk for overall morbidity and serious morbidity remained higher for the converted to open cohort. Features associated with conversion included male gender, higher body mass index, diabetes, smoking, and qualities pertaining to malignancy (>10% pre-operative weight loss, neoadjuvant chemotherapy, and neoadjuvant radiation) (12). Others have also reported male gender and malignancy as risk factors for conversion to open (11,13).

These data should not discourage the general practice of MIS distal pancreatectomy. A recently published Dutch randomized trial of MIS versus open distal pancreatectomy in fact supports an MIS approach as the standard of care for those with tumors <8 cm without vascular involvement (14). The authors found that functional recovery was faster in the MIS group *vs.* open by 2 days, with better quality of life in the early post-operative period. Again, the vast majority of the MIS operations in this study were laparoscopic (42/47, 89%), and 8% converted to an open procedure. In a large European cohort study of distal pancreatectomies only for adenocarcinoma, conversion to open was 19% (15). Importantly, they found that R0 resection was actually higher in the MIS group, and that overall survival was similar between MIS and open groups.

The conversion to open rate, as well as the patient length of hospitalization is inversely related to surgeon training and experience (13,16). The Dutch initiated a national training program when it was noted that the rate of MIS distal pancreatectomy had stagnated, and conversion to open rates remained >30%. Training consisted of technique description, video training, and on-site proctoring, with a total of ~8 hours of instruction. Patient outcomes were compared to a historical control. Despite the fact that patients treated after training had larger tumors and a higher number of adenocarcinomas, the odds of completing a distal pancreatectomy in a minimally invasive fashion increased 10-fold (16). Rate of conversion is also decreased for surgeons with >1-year experience, and those who have performed >15 operative cases (13). The learning curve as it relates to operative time for laparoscopic and robotic distal pancreatectomy has been reported to be 10–20 procedures (17–20). It has been suggested that learning curve is faster for operative novices with robotics compared to laparoscopy,

and that fewer errors are made with robotics compared to laparoscopy for experienced surgeons performing simple tasks (21,22). Head to head comparisons of skills acquisition as they pertain to distal pancreatectomy, though, have not been made. As such, the best approach is likely the one with which the operative surgeon has the greatest comfort.

## Acknowledgments

*Funding:* None.

## Footnote

*Provenance and Peer Review:* This article was commissioned by the editorial office, *Laparoscopic Surgery*. The article did not undergo external peer review.

*Conflicts of Interest:* C Stewart has received professional honoraria from Verb Surgical, unrelated to this manuscript. Y Fong is a scientific consultant to Medtronic, Johnson & Johnson, Olympus, Avra Robotics, and Perfint Robotics, all unrelated to this manuscript.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

1. Okunrintemi V, Gani F, Pawlik TM. National trends in postoperative outcomes and cost comparing minimally invasive versus open liver and pancreatic surgery. *J Gastrointest Surg* 2016;20:1836-43.
2. Bateni SB, Olson JL, Hoch JS, et al. Drives of cost in pancreatic surgery: It's not about hospital volume. *Ann Surg Oncol* 2018;25:3804-11.
3. Stewart CL, Meguid C, Chapman B, et al. Evolving

- trends towards minimally invasive surgery for solid-pseudopapillary neoplasms. *Ann Surg Oncol* 2016;23:4165-8.
4. Stewart CL, Raeburn C, Edil BH. Laparoscopic distal pancreatectomy for an ACTH secreting pancreatic neuroendocrine tumor. *Video Endocrinology* 2015;2:4.
  5. Stewart CL, Ituarte PHG, Melstrom KA, et al. Robotic surgery trends in general surgical oncology from the National Inpatient Sample. *Surg Endosc* 2018. [Epub ahead of print].
  6. Wright GP, Zureikat AH. Development of minimally invasive pancreatic surgery: An evidence-based systematic review of laparoscopic versus robotic approaches. *J Gastrointest Surg* 2016;20:1658-65.
  7. Casadei R, Ricci C, Pacilio CA, et al. Laparoscopic distal pancreatectomy: which factors are related to open conversion? Lessons learned from 68 consecutive procedures in a high-volume pancreatic center. *Surg Endosc* 2018;32:3839-45.
  8. Raoof M, Nota CLMA, Melstrom LG, et al. Oncologic outcomes after robot-assisted versus laparoscopic distal pancreatectomy: Analysis of the National Cancer Database. *J Surg Oncol* 2018;118:651-6.
  9. Gavriilidis P, Lim C, Menahem B, et al. Robotic versus laparoscopic distal pancreatectomy - The first meta-analysis. *HPB (Oxford)* 2016;18:567-74.
  10. Guerrini GP, Lauretta A, Belluco C, et al. Robotic versus laparoscopic distal pancreatectomy: an up-to-date meta-analysis. *BMC Surg* 2017;17:105-15.
  11. Lee SY, Allen PJ, Sadot E, et al. Distal pancreatectomy: a single institution's experience in open, laparoscopic, and robotic approaches. *J Am Coll Surg* 2015;220:18-27.
  12. Beane JD, Pitt HA, Dolejs SC, et al. Assessing the impact of conversion on outcomes of minimally invasive distal pancreatectomy and pancreatoduodenectomy. *HPB (Oxford)* 2018;20:356-63.
  13. Hua Y, Javed AA, Burkhart RA, et al. Preoperative risk factors for conversion and learning curve of minimally invasive distal pancreatectomy. *Surgery* 2017;162:1040-7.
  14. de Rooij T, van Hilst J, van Santvoort H, et al. Minimally Invasive Versus Open Distal Pancreatectomy (LEOPARD): A Multicenter patient-blinded randomized controlled trial. *Ann Surg* 2019;269:2-9.
  15. van Hilst J, de Rooij T, Klompmaker S, et al. Minimally Invasive versus Open Distal Pancreatectomy for Ductal Adenocarcinoma (DIPLOMA): A Pan-European propensity score matched study. *Ann Surg* 2019;269:10-7.
  16. de Rooij T, van Hilst J, Boerma D, et al. Impact of a nationwide training program in minimally invasive distal pancreatectomy (LAELAPS). *Ann Surg* 2016;264:754-62.
  17. Braga M, Ridolfi C, Balzano G, et al. Learning curve for laparoscopic distal pancreatectomy in a high-volume hospital. *Updates Surg* 2012;64:179-83.
  18. Ricci C, Casadei R, Buscemi S, et al. Laparoscopic distal pancreatectomy: what factors are related to the learning curve? *Surg Today* 2015;45:50-6.
  19. Napoli N, Kauffmann EF, Perrone VG, et al. The learning curve in robotic distal pancreatectomy. *Updates Surg* 2015;67:257-64.
  20. Shakir M, Boone BA, Polanco PM, et al. The learning curve for robotic distal pancreatectomy: an analysis of outcomes of the first 100 consecutive cases at a high-volume pancreatic centre. *HPB (Oxford)* 2015;17:580-6.
  21. Kim HJ, Choi GS, Park JS, et al. Comparison of surgical skills in laparoscopic and robotic tasks between experienced surgeons and novices in laparoscopic surgery: an experimental study. *Ann Coloproctol* 2014;30:71-6.
  22. Zihni A, Gerull WD, Cavallo JA, et al. Comparison of precision and speed in laparoscopic and robot-assisted surgical task performance. *J Surg Res* 2018;223:29-33.

doi: 10.21037/ls.2019.02.05

**Cite this article as:** Stewart C, Fong Y. Reducing conversion in minimally invasive distal pancreatectomy: patient selection, training, and robotics. *Laparosc Surg* 2019;3:11.