



Comparison between robotic Nissen, Toupet and Dor fundoplication: results

Diego Coletta¹, Celeste Del Basso¹, Paolo Ossola², Cristina De Padua¹, Giuseppe Giuliani³

¹Department of General Surgery, Emergency Department, Division of Emergency and Trauma Surgery, ²Department of General Surgery, Umberto I University Hospital, Sapienza University of Rome, Rome, Italy; ³Division of General and Minimally Invasive Surgery, Misericordia Hospital, Grosseto, Italy

Contributions: (I) Conception and design: D Coletta; (II) Administrative support: All authors; (III) Provision of study materials or patients: D Coletta; (IV) Collection and assembly of data: All authors; (V) Data analysis and interpretation: All authors; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Diego Coletta. Department of General Surgery, Emergency Department, Division of Emergency and Trauma Surgery, Umberto I University Hospital, Sapienza University of Rome, Viale del Policlinico 155, 00166 Rome, Italy. Email: diegocoletta1@gmail.com.

Abstract: Minimally invasive surgery has gained a role in the treatment of gastroesophageal reflux disease (GERD). Robotic fundoplication is one of choice of treatment in minimally invasive surgical managements. As for conventional and laparoscopic approaches, Nissen, Toupet and Dor are the mainly performed techniques of fundoplication with the robotic surgery. They differ in wrap construction both for the grades and for the anterior or posterior rotation. Some authors compared the three techniques in laparoscopic field in terms of symptoms control, failure of treatment and reintervention rate. We aimed to attempt a comparison between the three techniques performed by a robotic-assisted approach. The available literature on robotic fundoplication and relative different techniques is still low scarce. Except for intrinsic technical characteristics, the three techniques do not show particular differences between them. Total fundoplication seems to better control the GERD symptoms but has a higher rate of postoperative dysphagia than partial ones. Nissen fundoplication (NF) is the most practiced, standardized and described since its early employments. The lack of specific investigations about robotic Toupet fundoplication (TF) and Dor fundoplication (DF) with relative data preclude to compare the three mentioned techniques accurately. Although robotic surgery had proper characteristics, we could consider it as an advanced laparoscopic approach and thus we could liken the available data concerning the three laparoscopic fundoplication techniques to the corresponding robotic ones.

Keywords: Minimally invasive surgery; robotic fundoplication; Nissen; Toupet; Dor

Received: 26 April 2020; Accepted: 10 June 2020; Published: 25 January 2021.

doi: 10.21037/ls-20-87

View this article at: <http://dx.doi.org/10.21037/ls-20-87>

Introduction

Minimally invasive surgical approaches nowadays are employed as treatment for gastrointestinal malignancies but also for benign diseases. Minimally invasive fundoplication was first described in the early 90's (1) with the intent to reduce the invasiveness of surgical procedure for benign diseases such as gastroesophageal reflux disease (GERD).

About of a decade after, *Cadiere* performed the first robot-assisted laparoscopic Nissen fundoplication (NF) in a groundbreaking moment of the worldwide surgical panorama (2). Since then several studies comparing the robotic-assisted fundoplication with the laparoscopic one, were carried out (3-5). Nissen, Toupet and Dor are the mainly performed techniques of fundoplication with robotic approach, differing in the grade of wrap around esophagus




Figure 1 Techniques of fundoplication. (A) Nissen fundoplication 360° posterior wrap; (B) Toupet 270° posterior wrap; (C) Dor 180° anterior wrap.




Figure 2 Ports placement. Blue: port placements; red: assistant trocar; green: alternative scheme of placement.

of 360°, 270° and 180° respectively (6-8) (Figure 1).

This study aims to give a snapshot on robotic fundoplication, comparing the three techniques.

Technical characteristics

NF represents the most performed surgical procedure for GERD not responsive to pharmacological therapy (9-11). Since the first description, this technique provides a complete mobilization of the stomach fundus followed by a 360° posterior fundoplication. At the beginning of the XXI st. century, the first computer-assisted NF in response to the pure laparoscopic approaches were performed (2,12). Patient is placed in a supine and steep reverse Trendelenburg position on the operating theatre table. After the establishment of pneumoperitoneum, trocars placement follows the same scheme used in other gastro-esophageal surgical procedures, using two or three robotic arms associated to one or two assistant trocars. The scheme is the same for the three techniques and the positions of assistant trocars depends on surgeon preferences. Figure 2

depicts trocar placement.

The standardized laparoscopic approach to floppy NF is generally adopted: opening of the phrenoesophageal ligament in a left-to-right way and preserving of the hepatic branch of the anterior vagus nerve; dissection of crura and transhiatal mobilization is performed to allow approximately 3 cm of intra-abdominal esophagus; in order to ensure a tension free-wrap, short gastric vessel are divided and the crural is closed posteriorly with non-absorbable sutures. The procedure concludes with posterior wrap construction of about 2 cm, placing a bougie and with the most distal suture incorporating the anterior muscular wall of the esophagus (13,14).

Toupet fundoplication (TF) technique is similar to Nissen but stomach wrap finally results in a partial posterior plication of 270°.

In the Dor fundoplication (DF) technique the wrap is constructed anteriorly, with a partial 180° rotation, in which the lateral edges of the plication are represented by the great curve of the stomach sutured to the right and left crura. Generally, DF is associated to the Heller myotomy in the treatment of esophageal achalasia and the wrap is developed covering the miotomized area (15).

Complications and clinical outcomes

Not seem to exist specific intraoperative complications due to the similarity within the three techniques except for the obvious higher risk of esophageal perforation due to myotomy in DF rather than NF and TF. Robotic fundoplication, particularly NF, when compared with laparoscopic one had shown no significant difference in terms of operative time (mean range between 88–137 *vs.* 91–102 minutes), conversion rate (3.4% *vs.* 3.3%), intra and postoperative complications (5% *vs.* 9%) also for dysphagia (16.6 *vs.* 16.1%), reoperation rate (7.4% *vs.* 3.6%), length of hospital stay (mean from 2.9 to 4.4 *vs.* 3.3 to 4.1 days)

and costs (the difference between robotic and laparoscopic procedures was about 500–1,800 euros) (16).

Data available on robotic TF are very limited (5,17). To date the literature available concerns TF performed in association with others procedure, or results pooled in series including other types of fundoplication and then no specific data can be obtained (18,19).

Studies comparing the laparoscopic DF with posterior total or partial fundoplication have been performed without reaching sufficient evidence to support its recommendation as surgical treatment for GERD (20,21). However less postoperative dysphagia in anterior fundoplication were noted, contrariwise a less reflux symptoms control was achieved (22,23). Actually reports dealing with robotic DF are those in which this fundoplication technique is performed in combination with Heller myotomy in the management of esophageal achalasia.

As for postoperative dysphagia, laparoscopic NF and TF were compared in term of variable lengths of fundoplication (1.5 vs. 3 cm) (24). Superior reflux symptoms control in 3-cm-TF over the 1.5-cm t without differences in postoperative dysphagia have been reported. Instead in NF, length did not influence reflux control, but a higher rate of dysphagia was noted in 3-cm wrap group compared with the 1.5-cm ones (25,26).

Recurrence

Need for recurrence has been reported with mixed results. Strate *et al.* reported a higher reoperation rate in NF than TF cause of the persistence of dysphagia probably due to different preoperative esophageal dysmotility (22). Also concerning reinterventions rate, anterior fundoplication requires redo-surgery for scarce control of reflux rather than posterior ones (23).

Pediatric diseases

A recent meta-analysis of randomized controlled trials, comparing RF and LF in children, shows no clear conclusion to suggest robotic or laparoscopic fundoplication in term of safety, short-term efficacy, success rate, post-operative morbidity and redo-surgery rates (27).

Discussion

To data, unfortunately the available literature on robotic fundoplication and relative different techniques is still low.

The lack of specific case series, randomized controlled trials or retrospective study about robotic TF and DF with relative data preclude to compare the three mentioned techniques accurately.

Although robotic surgery had proper characteristics, we could consider it as an advanced laparoscopic approach and thus we could liken the available data concerning the laparoscopic NF, TF and DF to the corresponding robotic ones. Robotic technology has made possible to overcome some of the limitations of laparoscopic surgery including a reduction in tremors, the magnifying images with 3D technology, an increase in the effectiveness of surgical gestures through tools with more degrees of freedom, and mostly a great improvement in fatigue in relation to the continuation of surgery. If compared laparoscopic and robotic surgical approaches to GERD, seemed to be similar in adult and children in terms of operative times, length of hospital stay and post-operative complications. Robotic surgery shown obvious higher costs; some specific complications due to the lack of haptic feedback combined with the overhuman strength robot-assistance like gastric perforation during fundoplication were also reported (15,28,29).

Conclusions

Robot-assisted NF as surgical treatment for GERD is the most performed, codified and reported between robotic approaches since its early employments. Further studies analyzing the different techniques of robotic fundoplication are needed to better compare them and to suggest the best one to perform in routine surgical practice.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the Guest Editor (Andrea Balla) for the series “Minimally Invasive Approach for the Treatment of Gastro-esophageal Reflux Disease” published in *Laparoscopic Surgery*. The article has undergone external peer review.

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/ls-20-87>). The series “Minimally Invasive

Approach for the Treatment of Gastro-esophageal Reflux Disease” was commissioned by the editorial office without any funding or sponsorship. The authors have no other conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Dallemagne B, Weerts JM, Jehaes C, et al. Laparoscopic Nissen fundoplication: preliminary report. *Surg Laparosc Endosc* 1991;1:138-43.
2. Cadière GB, Himpens J, Vertruyen M, et al. Nissen fundoplication done by remotely controlled robotic technique. *Ann Chir* 1999;53:137-41.
3. Meininger D, Byhahn C, Markus BH, et al. Total endoscopic Nissen fundoplication with the robotic device "da Vinci" in children. Hemodynamics, gas exchange, and anesthetic management. *Anaesthesist* 2001;50:271-5.
4. Cadière GB, Himpens J, Vertruyen M, et al. Evaluation of telesurgical (robotic) NISSEN fundoplication. *Surg Endosc* 2001;15:918-23.
5. Melvin WS, Needleman BJ, Krause KR, et al. Computer-enhanced vs. standard laparoscopic antireflux surgery. *J Gastrointest Surg* 2002;6:11-5; discussion 15-6.
6. Nissen R. A simple operation for control of reflux esophagitis. *Schweiz Med Wochenschr* 1956;86:590-2.
7. Toupet A. Technique of Esophago-Gastroplasty With Phrenogastropexy Used in Radical Treatment of Hiatal Hernias as a Supplement to Heller's Operation in Cardiospasm. *Mem Acad Chir (Paris)* 1963;89:384-9.
8. Dor J, Humbert P, Dor V, et al. L'intérêt de la technique de Nissen modifiée la prevention du reflux après cardiomyotomie extra muqueuse de Heller. *Mem AcadChir* 1962;88:87723
9. Frazzoni M, Piccoli M, Conigliaro R, et al. Laparoscopic fundoplication for gastroesophageal reflux disease. *World J Gastroenterol* 2014;20:14272-9.
10. Linzberger N, Berdah SV, Orsoni P, et al. Laparoscopic posterior fundoplication in gastroesophageal reflux: mid-term results. *Ann Chir* 2001;126:143-7.
11. Mardani J, Lundell L, Engström C. Total or posterior partial fundoplication in the treatment of GERD: results of a randomized trial after 2 decades of follow-up. *Ann Surg* 2011;253:875-8.
12. Chapman WH, Young JA, Albrecht RJ, et al. Robotic Nissen fundoplication: alternative surgical technique for the treatment of gastroesophageal reflux disease. *J Laparoendosc Adv Surg Tech A* 2001;11:27-30.
13. Stefanidis D, Hope WW, Kohn GP, et al. Guidelines for surgical treatment of gastroesophageal reflux disease. *Surg Endosc* 2010;24:2647-69.
14. Attwood SE, Lundell L, Ell C, et al. Standardization of surgical technique in antireflux surgery: the LOTUS Trial experience. *World J Surg* 2008;32:995-8.
15. Falkenback D, Lehane CW, Lord RV. Robot-assisted oesophageal and gastric surgery for benign disease: antireflux operations and Heller's myotomy. *ANZ J Surg* 2015;85:113-20.
16. Yao G, Liu K, Fan Y. Robotic Nissen fundoplication for gastroesophageal reflux disease: a meta-analysis of prospective randomized controlled trials. *Surg Today* 2014;44:1415-23.
17. Jensen JS, Antonsen HK, Durup J. Two years of experience with robot-assisted anti-reflux surgery: A retrospective cohort study. *Int J Surg* 2017;39:260-6.
18. Petrucciani N, de'Angelis N, Brunetti F. Robotic Toupet fundoplication following Heller myotomy for achalasia (with video). *J Visc Surg* 2018;155:427-8.
19. Ceccarelli G, Romano A, Esposito G, et al. Robot-assisted Toupet fundoplication and associated cholecystectomy in symptomatic giant hiatal hernia with situs viscerum inversus-A case report and literature review. *Int J Surg Case Rep* 2019;60:371-5.
20. Engström C, Lönroth H, Mardani J, et al. An anterior or posterior approach to partial fundoplication? Long-term results of a randomized trial. *World J Surg* 2007;31:1221-5; discussion 1226-7.
21. Hagedorn C, Jönson C, Lönroth H, et al. Efficacy of an anterior as compared with a posterior laparoscopic partial fundoplication: results of a randomized, controlled clinical trial. *Ann Surg* 2003;238:189-96.
22. Strate U, Emmermann A, Fibbe C, et al. Laparoscopic

- fundoplication: Nissen versus Toupet two-year outcome of a prospective randomized study of 200 patients regarding preoperative esophageal motility. *Surg Endosc* 2008;22:21-30.
23. Spence GM, Watson DI, Jamieson GG, et al. Single center prospective randomized trial of laparoscopic Nissen versus anterior 90 degrees fundoplication. *J Gastrointest Surg* 2006;10:698-705.
 24. Cai W, Watson DI, Lally CJ, et al. Ten-year clinical outcome of a prospective randomized clinical trial of laparoscopic Nissen versus anterior 180(degrees) partial fundoplication. *Br J Surg* 2008;95:1501-5.
 25. Mickevicius A, Endzinas Z, Kiudelis M, et al. Influence of wrap length on the effectiveness of Nissen and Toupet fundoplication: a prospective randomized study. *Surg Endosc* 2008;22:2269-76.
 26. Jobe BA, Wallace J, Hansen PD, et al. Evaluation of laparoscopic Toupet fundoplication as a primary repair for all patients with medically resistant gastroesophageal reflux. *Surg Endosc* 1997;11:1080-3.
 27. Patti MG, Robinson T, Galvani C, et al. Total fundoplication is superior to partial fundoplication even when esophageal peristalsis is weak. *J Am Coll Surg* 2004;198:863-9; discussion 869-70.
 28. Cundy TP, Harling L, Marcus HJ, et al. Meta analysis of robot-assisted versus conventional laparoscopic fundoplication in children. *J Pediatr Surg* 2014;49:646-52.
 29. Dalsgaard T, Jensen MD, Hartwell D, et al. Robotic Surgery Is Less Physically Demanding Than Laparoscopic Surgery: Paired Cross Sectional Study. *Ann Surg* 2020;271:106-13.

doi: 10.21037/ls-20-87

Cite this article as: Coletta D, Del Basso C, Ossola P, De Padua C, Giuliani G. Comparison between robotic Nissen, Toupet and Dor fundoplication: results. *Laparosc Surg* 2021;5:10.