

Full-length article

Effects of AMP579 and adenosine on L-type Ca^{2+} current in isolated rat ventricular myocytesXiong WANG¹, Bo-wei WU, Dong-mei WU*Department of Physiology, Shanxi Medical University, Taiyuan 030001, China***Key words**

AMP579; adenosine; heart ventricles; cardiac myocytes; L-type calcium channels; patch-clamp techniques

¹ Correspondence to Prof Xiong WANG.
Phn 86-351-469-0162.
E-mail Wxiong@public.ty.sx.cnReceived 2004-08-23
Accepted 2004-12-11

doi: 10.1111/j.1745-7254.2005.00107.x

Abstract

Aim: To compare the effects of AMP579 and adenosine on L-type Ca^{2+} current ($I_{\text{Ca-L}}$) in rat ventricular myocytes and explore the mechanism by which AMP579 acts on $I_{\text{Ca-L}}$. **Methods:** $I_{\text{Ca-L}}$ was recorded by patch-clamp technique in whole-cell configuration. **Results:** Adenosine (10 nmol/L to 50 $\mu\text{mol/L}$) showed no effect on basal $I_{\text{Ca-L}}$, but it inhibited the $I_{\text{Ca-L}}$ induced by isoproterenol 10 nmol/L in a concentration-dependent manner with the IC_{50} of 13.06 $\mu\text{mol/L}$. Similar to adenosine, AMP579 also showed an inhibitory effect on the $I_{\text{Ca-L}}$ induced by isoproterenol. AMP579 and adenosine (both in 10 $\mu\text{mol/L}$) suppressed isoproterenol-induced $I_{\text{Ca-L}}$ by 11.1% and 5.2%, respectively. In addition, AMP579 had a direct inhibitory effect on basal $I_{\text{Ca-L}}$ in a concentration-dependent manner with IC_{50} (1.17 $\mu\text{mol/L}$). PD116948 (30 $\mu\text{mol/L}$), an adenosine A_1 receptor blocker, showed no action on the inhibitory effect of AMP579 on basal $I_{\text{Ca-L}}$. However, GF109203X (0.4 $\mu\text{mol/L}$), a special protein kinase C (PKC) blocker, could abolish the inhibitory effect of AMP579 on basal $I_{\text{Ca-L}}$. So the inhibitory effect of AMP579 on basal $I_{\text{Ca-L}}$ was induced through activating PKC, but not linked to adenosine A_1 receptor. **Conclusion:** AMP579 shows a stronger inhibitory effect than adenosine on the $I_{\text{Ca-L}}$ induced by isoproterenol. AMP579 also has a strong inhibitory effect on basal $I_{\text{Ca-L}}$ in rat ventricular myocytes. Activation of PKC is involved in the inhibitory effect of AMP579 on basal $I_{\text{Ca-L}}$ at downstream-mechanism.

Introduction

Recent studies showed that AMP579 was a novel adenosine agonist with high affinity for adenosine A_1 and A_2 receptors^[1,2]. Experiments in animal models have demonstrated that AMP579 reduced infarct size by 50% to 98% when administered before a final ischemic event (mediation of ischemic preconditioning) or just before reperfusion (attenuation of reperfusion injury)^[3,4]. Further experiments on pigs, dogs, and rabbits suggested that AMP579 was more powerful than adenosine in attenuating polymorphonuclear neutrophil-mediated inflammatory responses, dilating the coronary artery, reducing myocardial contracture and limiting infarct size^[5,6]. Although the protective effect of AMP579 required adenosine receptor activation, adenosine could not duplicate the effects.

The difference between pharmacologic effect of AMP579

and adenosine might reflect the differences in ionic mechanisms. It has been established that adenosine could cause an attenuation of basal $I_{\text{Ca-L}}$ only in unstimulated atrial myocytes, but under conditions of isoproterenol stimulation, adenosine could markedly attenuate isoproterenol induced- $I_{\text{Ca-L}}$ in both atrial and ventricular myocytes. However, little is known about the electrophysiological effects of AMP579 so far. This study will examine the effects of AMP579 and adenosine on L-type calcium channel and elucidate the mechanisms underlying the cardioprotective effect of AMP579 and its utility in treatment of myocardial ischemia-reperfusion injury.

Materials and methods

Rat myocardial cell isolation Ventricular myocytes were obtained from Wistar male rats (250–300 g) by enzymatic

isolation procedure. In brief, rats were killed by cervical dislocation and the heart was then immediately removed, cannulated through the aorta and perfused through the coronary artery with Ca^{2+} -free Tyrode's solution for 10 min. The composition of Ca^{2+} -free Tyrode's solution was: NaCl 140.0 mmol/L, KCl 5.4 mmol/L, MgCl_2 1.0 mmol/L, NaH_2PO_4 0.3 mmol/L, glucose 10.0 mmol/L, HEPES 5.0 mmol/L; pH adjusted to 7.4 with NaOH at room temperature. The heart was then perfused with enzymatic solution, which was low Ca^{2+} (CaCl_2 150 $\mu\text{mol/L}$) Tyrode's solution with collagenase P (0.3g/L) for about 8–10min. The left ventricle was then removed. The cells were isolated by gentle agitation and kept in Krebs buffer (KB) solution, which contained: KOH 85.0 mmol/L, *L*-glutamic acid 50.0 mmol/L, KCl 30.0 mmol/L, taurine 20.0 mmol/L, KH_2PO_4 30.0 mmol/L, MgCl_2 1.0 mmol/L, HEPES 10.0 mmol/L, glucose 10.0 mmol/L and egtazic acid 0.5 mmol/L; pH adjusted to 7.4 by KOH.

Electrophysiological measurement Whole-cell patch-clamp was used to record $I_{\text{Ca-L}}$ (L-type Ca^{2+} currents) and membrane capacitance was measured with a P-clamp 5.51 software package (Axon Instruments, USA). Patch electrodes were made from thin-walled glass capillaries (1.5 mm outside diameter) using a two-stage vertical microelectrode puller (model PP-83, Narishige Scientific Instruments, Japan). The electrode resistance ranges 3 $\text{M}\Omega$, when filled with pipette solution.

For the measurement of $I_{\text{Ca-L}}$, the extracellular solution contained: NaCl 140.0 mmol/L, CaCl_2 1.8 mmol/L, MgCl_2 1.0 mmol/L, KCl 5.4 mmol/L, glucose 10.0 mmol/L, NaH_2PO_4 0.3 mmol/L, and HEPES 10.0 mmol/L; pH adjusted to 7.4 with NaOH. The pipette solution contained: egtazic acid 10.0 mmol/L, KCl 140.0 mmol/L, Na_2ATP 2.0 mmol/L, HEPES 5.0 mmol/L, 4-AP 5.0 mmol/L, MgCl_2 1.0 mmol/L; pH adjusted to 7.4 with KOH. The calcium current was expressed as membrane current density (pA/pF). The cell capacitance was measured by the method previously described by Coetzee *et al*^[9]. $I_{\text{Ca-L}}$ was measured according to the method described by Hartzell *et al*^[10]. The AMP579 was a gift from Department of Cardiothoracic Surgery Research Laboratory, Emory University School of Medicine, USA. AMP579 was dissolved in small volumes of Me_2SO , then diluted to the desired final concentration before each experiment.

Statistic analysis Data were expressed as mean \pm SD. Statistical significance was determined by Student's *t*-test and $P < 0.05$ was considered significant.

Results

Detection of L-type calcium channel current The cal-

cium current was activated by depolarizing pulse from a holding potential of -40 mV to +10 mV at 50 mV step-voltage. This inward current could be completely inhibited by 1 $\mu\text{mol/L}$ verapamil, the basic characteristics indicated that the current present in rat ventricular myocytes was L-type Ca^{2+} current.

Effect of AMP579 and adenosine on L-type calcium current In the presence of adenosine at 10 nmol/L, 1, 10, and 50 $\mu\text{mol/L}$, $I_{\text{Ca-L}}$ varied from 4.9 ± 0.9 to 4.8 ± 0.9 , 4.9 ± 0.9 , 4.9 ± 0.9 , 4.7 ± 0.9 pA/pF, respectively ($n=5$, $P > 0.05$). Adenosine had no effect on basal $I_{\text{Ca-L}}$. However, when $I_{\text{Ca-L}}$ was augmented to 2.7 ± 0.6 pA/pF by 10 nmol/L isoproterenol, adenosine at 10 nmol/L, 1, 10, and 50 $\mu\text{mol/L}$ significantly reduced it to 2.4 ± 0.6 , 2.1 ± 0.6 , 2.0 ± 0.5 , and 1.9 ± 0.5 pA/pF, respectively ($n=4$, $P < 0.05$). Adenosine showed an inhibitory effect on isoproterenol-induced $I_{\text{Ca-L}}$ in a concentration-dependent manner with the IC_{50} of 13.06 $\mu\text{mol/L}$ (Figure 1, 2).




Figure 1. Effect of adenosine on $I_{\text{Ca-L}}$ in isolated rat ventricular myocytes. (a) Control; (b) 50 $\mu\text{mol/L}$ adenosine.

Effect of AMP579 on $I_{\text{Ca-L}}$ Isoproterenol 10 nmol/L augmented $I_{\text{Ca-L}}$ to 3.8 ± 0.7 pA/pF. AMP579 10 $\mu\text{mol/L}$ reduced $I_{\text{Ca-L}}$ to 2.4 ± 0.1 pA/pF ($P < 0.05$, $n=3$, Figure 3), AMP579 also showed an inhibitory effect on isoproterenol-induced $I_{\text{Ca-L}}$. AMP579 and adenosine (both 10 $\mu\text{mol/L}$) suppressed isoproterenol-induced $I_{\text{Ca-L}}$ by 11.1% and 5.2%, respectively. AMP579 had a stronger inhibitory effect. In contrast to adenosine, AMP579 possessed a direct inhibitory effect on basal $I_{\text{Ca-L}}$ in a concentration-dependent manner with the IC_{50} of 1.17 $\mu\text{mol/L}$ (Table 1, Figure 4).

AMP579 10 $\mu\text{mol/L}$ markedly reduced basal $I_{\text{Ca-L}}$ from 2.5 ± 1.2 to 2.0 ± 1.0 pA/pF ($n=5$, $P < 0.05$). Infusion of PD116948 30 $\mu\text{mol/L}$, an adenosine A_1 receptor blocker, did not abolish the inhibitory effects of AMP579 on $I_{\text{Ca-L}}$ (1.9 ± 0.6 vs 2.0 ± 1.0 pA/pF, $P > 0.05$). But under the same conditions AMP579 10




Figure 2. The inhibitory effect of adenosine on I_{Ca-L} induced by isoproterenol in isolated rat ventricular myocytes. (a) Control; (b) isoproterenol 10 nmol/L; (c) adenosine 50 μ mol/L.




Figure 3. The inhibitory effect of AMP579 on I_{Ca-L} induced by isoproterenol in isolated ventricular myocytes. (a) Control; (b) isoproterenol 10 nmol/L; (c) AMP579 10 μ mol/L.

μ mol/L markedly reduced the I_{Ca-L} from 2.4 ± 0.4 to 1.8 ± 0.4 pA/pF ($n=4$, $P<0.01$). Infusion of 0.4μ mol/L GF109203X, a PKC blocker, significantly reversed it to 2.2 ± 0.4 pA/pF ($P<0.05$, Figure 5). So GF109203X could abolish the inhibitory effect of AMP579, indicating that the inhibitory effect on basal I_{Ca-L} by AMP579 was induced through activating PKC but not linked to the adenosine A_1 receptor.

Table 1. Effect of AMP579 on basal I_{Ca-L} in rat ventricular myocytes. $n=5$. Mean \pm SD. ^b $P<0.05$, ^c $P<0.01$ vs corresponding control group.

AMP579 concentration	I_{Ca-L} value/ pA·pF ⁻¹	Change rate/ %
0 (Control)	2.80 ± 0.75	
10 nmol/L	2.66 ± 0.75^b	-5.0
1 μ mol/L	2.36 ± 0.71^b	-15.7
10 μ mol/L	2.03 ± 0.72^c	-27.5
50 μ mol/L	1.78 ± 0.70^c	-36.4

Change rate=(the current value after administration of drug-control value)/control value \times 100%




Figure 4. Effect of AMP579 on I_{Ca-L} in isolated rat ventricular myocytes. (a) Control; (b) AMP579 50 μ mol/L.




Figure 5. Abolition of inhibitory effects of AMP579 on I_{Ca-L} by a PKC blocker in isolated rat ventricular myocytes. (a) Control; (b) AMP579 10 μ mol/L; (c) GF109203X 0.4 μ mol/L.

Discussion

In cardiac tissue, a direct inhibition of basal I_{Ca-L} by adenosine has only been demonstrated in guinea-pig atrial and ferret ventricular myocytes^[11,12]. But in the presence of isoproterenol stimulation, adenosine has prominent inhibitory effects on I_{Ca-L} in ventricular myocytes^[13]. These may reflect differences in receptor-effector coupling mechanisms, the level of basal adenylate cyclase activity, the basal phosphorylated state of Ca^{2+} channels and/or the effect of phosphorylation on the gating of L-type Ca^{2+} channel. Consistent with previous reports, our experiment shows that adenosine has no direct inhibitory effect on basal I_{Ca-L} in the rat ventricle, but in the condition that isoproterenol was previously administered, adenosine shows an inhibitory effect on the I_{Ca-L} induced by isoproterenol with an IC_{50} of 13.06 $\mu\text{mol/L}$, suggesting that adenosine exerts an indirect inhibitory effect on I_{Ca-L} in the rat ventricle by inhibition of isoproterenol stimulation.

In contrast to adenosine, AMP579 shows a direct inhibitory effects on basal I_{Ca-L} in the rat ventricle with IC_{50} of 1.17 $\mu\text{mol/L}$. The blocking of Ca^{2+} influx by L-type Ca^{2+} channel could serve as an efficient method for protecting the ischemic myocyte by minimizing ischemia-induced Ca^{2+} overload and irreversible cell contracture and autodigestion by Ca^{2+} -dependent proteases^[14]. Therefore, by reducing both basal I_{Ca-L} and isoproterenol-induced I_{Ca-L} , AMP579 will play a more important role in negative chronotropic and negative dromotropic effects. These action mechanism differences between AMP579 and adenosine may account for the contribution of AMP579 in reducing neutrophil-mediated inflammatory reaction, inhibiting cardiac contraction, dilating coronary vessels, attenuating ischemia and reperfusion injury.

Our study does not show that adenosine A_1 receptor is linked to inhibition of AMP579 on basal I_{Ca-L} . At present, available data indicate that three pathways are involved in receptor-linked downstream mechanisms for inhibition of I_{Ca-L} by adenosine. The first is cAMP-PKA, as PKA increase I_{Ca-L} by phosphorylation on the gating of the L-type calcium channel, inhibitions of adenylate cyclase and reductions of cAMP and PKA levels by adenosine result in attenuation on I_{Ca-L} ^[12]. Second is that activation of guanylate cyclase results in increments of intracellular cGMP and PKG concentration, which in turn inhibits phosphorylation on the gating of the L-type calcium channel^[15]. The third is modulated by PKC, because there are different PKC subunits which result in different effects^[16]. Our experiment finds that special PKC antagonist GF109203X can totally eliminate inhibitory effects of AMP579 on I_{Ca-L} , suggesting that AMP579 exerts a direct inhibitory effects on the L-type

calcium channel through the PKC pathway.

References

- 1 Nakamura M, Zhao ZQ, Clark KL, Velez DV, Guyton RA, Vinter-Johansen J. A novel adenosine analog, AMP579, inhibits neutrophil activation, adherence and neutrophil-mediated injury to coronary vascular endothelium. *Eur J Pharmacol* 2000; 397: 197–205.
- 2 Sledeski AW, Kubiak GG, O'Brien MK, Powers MR, Powener TH, Truessedale LK. Efficient synthesis of AMP579, a novel adenosine A_1/A_2 receptor agonist. *J Org Chem* 2000; 65: 8114–9.
- 3 Budde JM, Velez DA, Zhao ZQ, Clark KL, Morris CD, Muraki S, *et al*. Comparative study of AMP579 and adenosine inhibition of neutrophil-mediated vascular and myocardial injury during 24 h of reperfusion. *Cardiovasc Res* 2000; 47: 294–305.
- 4 Xu Z, Downey JM, Cohen MV. AMP579 reduces contracture and limits infarction in rabbit heart by activating adenosine A_2 receptors. *J Cardiovasc Pharmacol* 2001; 38: 474–81.
- 5 Mcvey MJ, Smiths GJ, Cox BF, Kitzen JM, Clark KL, Perrone MH. Cardiovascular pharmacology of the adenosine A_1/A_2 -receptor agonist AMP579: coronary hemodynamic and cardioprotective effects in the canine myocardium. *J Cardiovasc Pharmacol* 1999; 33: 701–10.
- 6 Smits GJ, Mcvey M, Cox BF, Perrone MH, Clark KL. Cardioprotective effects of the novel A_1/A_2 receptor agonist AMP579 in a porcine model of myocardial infarction. *J Pharmacol Exp Ther* 1998; 286: 611–8.
- 7 Pelleg A, Belardinelli C. Cardiac electrophysiology and pharmacology of adenosine: basic and clinical aspects. *Cardiovasc Res* 1993; 27: 54–61.
- 8 Belardinelli L, Linden J, Berne RM. The cardiac effects of adenosine. *Prog Cardiovasc Dis* 1989; 32: 73–97.
- 9 Coetzee WA, Ichikawa H, Hearse DJ. Oxidant stress inhibits Na^+/Ca^{2+} exchange in cardiac myocytes: mediation by sulfhydryl groups? *Am J Physiol* 1994; 266: H909–19.
- 10 Hartzell HC, Simmons MA. Comparison of effects of acetylcholine on calcium and potassium currents in frog atrium and ventricle. *J Physiol* 1987; 89: 411–22.
- 11 Cerbai E, Klockner U, Isenberg G. Ca^{2+} -antagonistic effects of adenosine in guinea-pig atrial cell. *Am J Physiol* 1988; 255: H872–8.
- 12 Qu Y, Campbell DL, Whorton AR, Strauss HC. Modulation of basal L-type Ca^{2+} current by adenosine in ferret isolated right ventricular myocytes. *J Physiol* 1993; 471: 269–93.
- 13 Isenberg G, Belardinelli L. Ionic basis for the antagonism between adenosine and isoproterenol on isolated mammalian ventricular myocytes. *Circ Res* 1984; 55: 309–25.
- 14 Eckert R, Utz J, Trautwin W, Mentzer RM, Wis M, Saar H. Involvement of intracellular Ca^{2+} release mechanism in adenosine-induced cardiac Ca^{2+} current inhibition. *Surgery* 1993; 114: 334–42.
- 15 Shen JB, Pappano AJ. On the role of phosphatase in regulation of cardiac L-type calcium current by cyclic GMP. *J Pharmacol Exp Ther* 2002; 301: 501–6.
- 16 Kameyama M, Hofmann F, Trautwein W. On the mechanism of beta-adrenergic regulation of the Ca^{2+} channel in the guinea-pig heart. *Pflugers Arch* 1985; 405: 285–93.