

## Full-length article

# High-throughput screening assay for new ligands at human melatonin receptors<sup>1</sup>

Jian-hua YAN<sup>2</sup>, Hao-ran SU<sup>2</sup>, Jean A BOUTIN<sup>3</sup>, M Pierre RENARD<sup>3</sup>, Ming-wei WANG<sup>2,4</sup>

<sup>2</sup>The National Center for Drug Screening and the State Key Laboratory of Drug Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 201203, China; <sup>3</sup>Les Laboratoires Servier, 22 rue Garnier, 92200 Neuilly-sur-Seine, France

## Key words

melatonin; receptor; selectivity; agonist; antagonist

<sup>1</sup> Project supported in part by the Shanghai Municipality Science and Technology Development Fund (No 06DZ22907 and 07DZ22920), the Ministry of Science and Technology (No 2004CB518902), and Servier Beijing Pharmaceutical Research and Development.

<sup>4</sup> Correspondence to Dr Ming-wei WANG.  
Phn 86-21-5080-0598.  
Fax 86-21-5080-0721.  
E-mail wangmw@mail.shnc.ac.cn

Received 2008-07-31  
Accepted 2008-09-22

doi: 10.1111/j.1745-7254.2008.00903.x

## Abstract

**Aim:** Melatonin (MT) is a neurohormone produced and secreted primarily by the pineal gland in a circadian manner, and mainly acts through 2 receptor subtypes: MT<sub>1</sub> and MT<sub>2</sub> in humans. The diversity in their tissue distribution is in favor of different functions for each receptor subtype. Selective modulators are therefore required to determine the physiological roles of these melatonin receptor subtypes and their implications in pathological processes. **Methods:** A homogenous MT<sub>1</sub>/MT<sub>2</sub> receptor binding assay was established for high-throughput screening of new ligands at the hMT<sub>1</sub> and/or hMT<sub>2</sub> receptors. The functional properties (agonists or antagonists) were assessed by a conventional guanosine-5' [ $\gamma$ -<sup>35</sup>S] triphosphate (GTP- $\gamma$ S) assay. **Results:** Three hMT<sub>1</sub> receptor-selective small molecule antagonists and 1 hMT<sub>2</sub> receptor-selective small molecule antagonist with novel structural features were identified following a high-throughput screening campaign of 48 240 synthetic and natural compounds. **Conclusion:** The findings may assist in the expansion of chemical probes to these 2 receptor subtypes.

## Introduction

Melatonin (*N*-acetyl-5-methoxytryptamine, MT) is a neurohormone synthesized and secreted primarily by the pineal gland in a circadian manner, with peak levels in all species occurring during the period of darkness. The synthesis of melatonin within the pineal gland is mainly regulated by the daily and seasonal changes in the environmental light/dark cycle. Melatonin, which is released into the blood circulation, is thought to transduce photoperiodic information to all the tissues expressing its receptors or other binding sites<sup>[1]</sup>. After its discovery and chemical characterization in 1959<sup>[2]</sup>, many different physiological and behavioral responses have been ascribed to melatonin. It not only regulates the sleep/wake cycle<sup>[3]</sup>, but also affects cardiovascular<sup>[4]</sup>, reproductive<sup>[5]</sup>, cell growth<sup>[6]</sup>, and retinal processes<sup>[7]</sup>.

The discovery of different melatonin receptors was facilitated by the introduction of the agonist radioligand 2-[<sup>125</sup>I]iodomelatonin<sup>[8]</sup>. Two human melatonin receptor

subtypes, hMT<sub>1</sub> and hMT<sub>2</sub>, have been cloned and shown to be G-protein-coupled receptors (GPCR)<sup>[9–11]</sup>. MT<sub>3</sub>, a third melatonin binding site, has been subsequently described in hamsters as the human homolog of the cytoplasmic quinone reductase-2<sup>[12,13]</sup>. The homology between hMT<sub>1</sub> and hMT<sub>2</sub> receptors at the amino acid level is 55% with approximately 70% overall identity within the transmembrane domains. The activation of these 2 receptor subtypes leads to the inhibition of adenylyl cyclase, probably through the activation of a G<sub>i</sub>-protein<sup>[9,10]</sup>. Studies of their tissue distribution in mammals have shown that these receptors are localized in different areas of the brain (suprachiasmatic nucleus, pars tuberalis, hypothalamus, cerebellum, cortex, hippocampus, and cerebral vessels), and at the peripheral level in the kidney, small intestine, and caudal arteries<sup>[9,10,14–19]</sup>. The differences in tissue distribution are in favor of different functions for each receptor subtype. Pharmacological investigations with selective agonists or antagonists are therefore necessary to determine the physiological roles of these melatonin

receptor subtypes and their implications in pathological processes. This information will probably open new therapeutic perspectives for selective ligands, different from the chronobiotic properties of melatonin clearly demonstrated in humans<sup>[20,21]</sup>.

Scintillation proximity assay (SPA) technology is a homogeneous approach that does not involve post-reaction liquid handling steps and is well-suited to automation and high-throughput screening (HTS). In the SPA system, membranes that express a particular receptor are attached to a microbead coated with wheat germ agglutinin. A radioisotope atom (eg [<sup>125</sup>I]) is brought very close to a scintillant-impregnated microbead by binding to the receptor. Because the emitted  $\beta$  particles can only travel short distances in the bulk solution, the microbeads preferentially capture electrons from the bound radiolabeled ligand. Thus, the amount of light emitted from the scintillant in the microbeads is directly proportional to the amount of bound radiolabeled ligand<sup>[22]</sup>.

In this study, we screened a library of 48 240 synthetic (91.2%) and natural compounds (8.8%) made in house, collected from various research institutions across China or purchased from commercial sources, with the goal of finding novel selective and potent hMT<sub>1</sub> and/or hMT<sub>2</sub> receptor agonists and antagonists. Using SPA technology, binding affinities were determined with 2-[<sup>125</sup>I]iodomelatonin and their functional activities by guanosine-5'-[ $\gamma$ -<sup>35</sup>S]triphosphate ([<sup>35</sup>S]-GTP- $\gamma$ S) binding assay<sup>[23]</sup>. As a result, we discovered a series of selective antagonists for hMT<sub>1</sub> and hMT<sub>2</sub> receptors.

## Materials and methods

**Reagents** The radioligand 2-[<sup>125</sup>I]iodomelatonin (specific activity: 81.4 TBq/mmol) and FlashBlue GPCR beads were purchased from PerkinElmer (Boston, MA, USA). [<sup>35</sup>S]GTP- $\gamma$ S (specific activity: 37 TBq/mmol) was obtained from Amersham Biosciences (Buckinghamshire, England). Guanosine diphosphate (GDP), GTP- $\gamma$ S, saponin, bovine serum albumin (BSA), HEPES, and melatonin were from Sigma-Aldrich (St Louis, MO, USA). Fetal bovine serum (FBS) was bought from Hyclone (Logan, UT, USA), and HamF12 medium was purchased from GIBCO BRL (Rockville, MD, USA).

**Membrane preparation** Chinese hamster ovary (CHO) cells stably expressing hMT<sub>1</sub> or hMT<sub>2</sub> receptors were provided by Servier (Neuilly-sur-Seine, France)<sup>[24]</sup>. The cells were cultured in HamF12 medium supplemented with 10% FBS, 2 mmol/L glutamine, 100 IU/mL penicillin,

and 100  $\mu$ g/mL streptomycin. They were grown at confluence, harvested in phosphate-buffered saline, and centrifuged at 1000 $\times$ g for 5 min (4 °C). The resulting pellet was suspended in isotonic buffer (5 mmol/L Tris/HCl, 0.2 mmol/L MgCl<sub>2</sub>, and 0.25 mol/L sucrose, pH 7.4), and homogenized using the BioNeb Cell Disruption System (Terre Haute, IN, USA). The homogenate was then centrifuged (20 000 $\times$ g for 30 min at 4 °C), and the resulting pellet was suspended in binding buffer 1 (50 mmol/L Tris/HCl, pH 7.4, 5 mmol/L MgCl<sub>2</sub>, 1 mmol/L EDTA). The protein content was determined using the Bradford assay<sup>[25]</sup>. Aliquots of membrane preparations were stored at -80 °C until use.

**Homogenous binding assay** The membranes were incubated overnight in binding buffer 1 containing 18 pmol/L 2-[<sup>125</sup>I]iodomelatonin, 0.75 mg/mL FlashBlue GPCR beads, various titrations of melatonin from a stock solution of 40  $\mu$ mol/L, and library compounds with an average concentration of 6.7  $\mu$ mol/L (final volume: 100  $\mu$ L). Non-specific binding was defined with 1  $\mu$ mol/L melatonin. Data were analyzed with GraphPad PRISM (GraphPad Software, San Diego, CA, USA). The competitive inhibition constant ( $K_i$ ) was calculated according to the Cheng-Prusoff equation:  $K_i = IC_{50} / (1 + L / K_d)$ , where  $IC_{50}$  is the concentration that produced 50% inhibition,  $K_d$  is the dissociation constant of inhibitor-binder reaction, and  $L$  is the concentration of the radiolabeled ligand used<sup>[26]</sup>.


**[<sup>35</sup>S]GTP- $\gamma$ S binding assay** The membranes and test compounds were diluted in binding buffer 2 (20 mmol/L HEPES, pH 7.4, 100 mmol/L NaCl, 3 mmol/L MgCl<sub>2</sub>, 12  $\mu$ mol/L GDP, and 0.1% BSA) in the presence of 20  $\mu$ g/mL saponin in order to enhance the agonist-induced stimulation level<sup>[27]</sup>, and pre-mixed with 1 mg/mL FlashBlue GPCR beads. For the agonist test, incubation was started by the addition of 0.2 nmol/L [<sup>35</sup>S]GTP- $\gamma$ S to the membranes and test compounds, and continued for 4 h at room temperature in a final volume of 100  $\mu$ L. To study the antagonist activity, the membranes were pre-incubated for 2 h with melatonin (30 nmol/L) in conjunction with a given concentration of a test compound. Reaction begun with the addition of 0.2 nmol/L [<sup>35</sup>S]GTP- $\gamma$ S followed by 4 h incubation at room temperature. Non-specific binding was assessed using non-radiolabeled GTP- $\gamma$ S (10  $\mu$ mol/L). Data were analyzed with PRISM (GraphPad, USA) to calculate the 50% effective concentration ( $EC_{50}$ ) and the maximal effect expressed as a percentage of that observed with 1  $\mu$ mol/L melatonin, 100% ( $E_{max}$ ) for agonists.  $K_B$  was used to describe antagonist potency:  $K_B = IC_{50} / (1 + [agonist] / EC_{50})$ , where  $IC_{50}$  is the antagonist concentration that gives

50% inhibition of [<sup>35</sup>S]GTP-γS binding in the presence of a fixed concentration of agonist, and EC<sub>50</sub> is the 50% effective concentration of an agonist when tested alone<sup>[26]</sup>. The maximal inhibitory effect (*I*<sub>max</sub>) was expressed as a percentage of that observed with 30 nmol/L melatonin for hMT<sub>1</sub> or hMT<sub>2</sub> receptors.


## Results

**HTS campaign** In the validation process, various assay parameters were studied. The optimal concentrations of radioligand and microbeads were 18 pmol/L and 0.75 mg/mL, respectively, for both receptor subtypes, while protein concentrations were optimized to 2.6 μg/mL for the hMT<sub>1</sub> receptor and 0.9 μg/mL for the hMT<sub>2</sub> receptor, respectively (data not shown). The control compound, melatonin, was tested at both receptors to evaluate the system. As shown in Figure 1, the *K*<sub>i</sub> values of melatonin were 0.87 nmol/L for the hMT<sub>1</sub> receptor and 2.07 nmol/L for the hMT<sub>2</sub> receptor, respectively, compatible to that reported in the literature<sup>[24]</sup>. In order to apply the assay to a HTS format, the *Z*' value and the signal/background (S/B; ie specific vs non-specific binding) were studied. The *Z*' factor, a screening window coefficient, is reflective of

both the assay signal dynamic range and the data variation associated with the signal measurements. Therefore, it is suitable for the assay quality assessment<sup>[28]</sup>. As shown in Figure 2, the *Z*' value for the hMT<sub>1</sub> receptor was 0.81 with a S/B ratio of 43.2 and a coefficient of variation value of 5.7%. In the case of the hMT<sub>2</sub> receptor, the corresponding values were 0.76, 9.08, and 5.4%, respectively. These characteristics suggest that the 2 assay systems are of high quality and suitable for HTS<sup>[28]</sup>. The number of hits at different inhibition rates discovered from a parallel HTS campaign directed towards hMT<sub>1</sub> and hMT<sub>2</sub> receptors are shown in Figure 3. Of the 48 240 samples screened (which did not include 4 P-PDOT and other 17 ligands known to react with hMT<sub>1</sub> and hMT<sub>2</sub> receptors<sup>[24]</sup>), 205 and 199 initial hits were identified for hMT<sub>1</sub> and hMT<sub>2</sub> receptors (≥85% binding activities), respectively. Among them, 63 (hMT<sub>1</sub>) and 54 (hMT<sub>2</sub>) hits were subsequently confirmed



**Figure 1.** Displacement of 2-[<sup>125</sup>I]iodomelatonin binding to hMT<sub>1</sub> receptor- (A) or hMT<sub>2</sub> receptor-expressing CHO cell membrane preparation (B) by melatonin. Data shown are mean±SD of triplicate measurements and representative of at least 3 independent experiments.



**Figure 2.** *Z*' value and S/B ratio determination. Homogenous hMT<sub>1</sub> (A) or hMT<sub>2</sub> (B) receptor binding assay was performed at optimized conditions with respective receptor-expressing CHO cells, 2-[<sup>125</sup>I]iodomelatonin (18 pmol/L) and melatonin (1 μmol/L). Background is the radioactivity counts in 2.5% DMSO solution where melatonin is dissolved. Forty replicates of S/B readouts were studied.

and subjected to  $IC_{50}$  determination, and the values of these confirmed hits were within the range of dozens of nmol/L to a few  $\mu\text{mol/L}$  for both  $hMT_1$  and  $hMT_2$  receptors (Table 1 and data not shown).

**Functionality characterization** In order to examine the pharmacological properties (agonist or antagonist) of these compounds at each melatonin receptor subtype, [ $^{35}\text{S}$ ]GTP- $\gamma\text{S}$  binding assays were performed. For both subtypes, the optimal concentrations of saponin,  $\text{MgCl}_2$ , and GDP were 20  $\mu\text{g/mL}$ , 3 mmol/L and 12  $\mu\text{mol/L}$ , respectively (data not shown), similar to that reported previously<sup>[24]</sup>. The agonist activities of melatonin at  $hMT_1$  ( $EC_{50}=1.91$  nmol/L) and  $hMT_2$  receptors ( $EC_{50}=1.37$  nmol/L) are shown in Figure 4. Based on the novelty in molecular structures and potency in binding activities, we selected 18 confirmed hits identified from the HTS campaign for simultaneous functionality studies with both receptor subtypes. As shown in Table 1, melatonin acted as a full agonist for both  $hMT_1$  and  $hMT_2$  receptors. Of the 18 compounds, NC00090702, NC00096199, and NC00099763 displayed selective  $hMT_1$  receptor antagonist activities with  $K_B$  values of 0.26, 0.50, and 1.06  $\mu\text{mol/L}$ , respectively; NC00080421 exhibited selective  $hMT_2$

receptor antagonist activities with a relatively more potent  $K_B$  value of 0.15  $\mu\text{mol/L}$ . The chemical structures of the 4 selective antagonists, all obtained from a commercial source (Specs, Delft, the Netherlands), are presented in Figure 5. Figure 6 depicts both the agonist and antagonist features of these compounds, as defined by  $E_{\text{max}}$  and  $K_B/I_{\text{max}}$ . Apart from NC00080421, which demonstrated some partial agonist activities (at the  $hMT_1$  receptor in particular), these antagonists are highly selective for a particular receptor subtype.


## Discussion

SPA is an innovative approach to assay development and biochemical screening that allows the rapid and sensitive measurement of a wide variety of molecular interactions in a homogeneous system. In the present study, we employed this technology in the HTS of 48 240 samples against 2 melatonin receptor subtypes:  $hMT_1$  and  $hMT_2$ . Not only did it show strong reliability in terms of consistent  $EC_{50}$  values measured for the control ligand (melatonin) and assay parameters exemplified by the high  $Z'$  values (0.81 for the  $hMT_1$  receptor and 0.76 for the

**Table 1.** Binding properties and functionality characteristics of 18 confirmed hits at two melatonin receptor subtypes ( $MT_1$  and  $MT_2$ ).

Compound	Receptor binding $K_i$ (nmol/L)		GTP $\gamma$ S assay $hMT_1$				GTP $\gamma$ S assay $hMT_2$				Structural feature
	$hMT_1$	$hMT_2$	$EC_{50}$ (nmol/L)	$E_{\text{max}}$ (%)	$K_B$ (nmol/L)	$I_{\text{max}}$ (%)	$EC_{50}$ (nmol/L)	$E_{\text{max}}$ (%)	$K_B$ (nmol/L)	$I_{\text{max}}$ (%)	
Melatonin	0.87	2.07	1.91	100	NA	NA	1.37	100	NA	NA	Indole
NC00004068	43	61	101	85	NA	24	98	71	NA	<10	Indole
NC00076880	56	21	NA	<10	463	62	> $10^4$	47	NA	33	Thieno[2,3-d]pyrimidine
NC00080421	238	348	NA	22	6324	75	NA	11	153	71	Tetrahydroisoquinoline
NC00085323	1573	ND	NA	<10	15401	128	NA	<10	9028	119	Benzenesulfonamide
NC00085862	ND	34	NA	<10	> $10^4$	57	62	130	NA	<10	Naphthalene
NC00085864	79	ND	NA	12	514	83	442	78	NA	<10	Naphthalene
NC00085892	475	ND	727	58	NA	33	300	97	NA	<10	Thiophene
NC00086791	ND	86	620	77	NA	39	257	71	NA	<10	Thiophene
NC00087441	ND	208	NA	<10	6588	92	NA	62	> $10^4$	39	Thieno[2,3-d]pyrimidine
NC00089216	ND	2535	NA	<10	2591	103	NA	8.5	4750	88	Benzoimidazole
NC00090038	18	ND	NA	83	NA	41	NA	19	NA	42	Dibenzo[a,c]cycloheptene
NC00090702	512	ND	NA	<10	264	120	NA	<10	NA	35	Dibenzylamide
NC00094488	1402	1216	NA	<10	NA	55	NA	4.9	12605	75	2-Oxo-[1,3]dithiole
NC00096199	1370	ND	NA	<10	501	120	NA	<10	NA	50	Benzulfamide
NC00096661	222	306	1290	126	NA	NA	97	120	NA	<10	Phenylethylamide
NC00098918	842	ND	NA	<10	NA	35	NA	16	NA	15	2-Hydroxy-naphthalene
NC00099763	1609	ND	NA	<10	1056	100	NA	<10	NA	<10	Propionamide
NC00100636	124	ND	> $10^4$	58	> $10^4$	47	NA	<10	NA	30	Dibenzo[a,c]cycloheptene


ND, not determined as the compound was not an HTS hit at this receptor subtype; NA, not available, unable to calculate or the value was more than 20  $\mu\text{mol/L}$ .




**Figure 3.** HTS of 48 240 compounds using hMT<sub>1</sub> (A) or hMT<sub>2</sub> (B) receptor binding assays. Results are expressed as percentage inhibition of radiolabeled 2-[<sup>125</sup>I]iodomelatonin binding to the receptors by the number of hit compounds at each 5% interval.

hMT<sub>2</sub> receptor), this affinity detection-based HTS method successfully identified a series of active compounds with a confirmed hit rate of 0.13% and 0.11% for the hMT<sub>1</sub> and hMT<sub>2</sub> receptors, respectively. Subsequent IC<sub>50</sub> determination and functionality studies with GTP-γS assays verified the results obtained from the HTS campaign. It was noted that the S/B window of our GTP-γS assay was somehow reduced (data not shown), probably due to the lack of wash step, as compared to that reported elsewhere<sup>[24]</sup>. Nevertheless, the system was robust enough for the eventual identification of several novel melatonin receptor modulators.

Of the 4 functionally-selective compounds, 1 was characterized as a hMT<sub>2</sub> antagonist and 3 as hMT<sub>1</sub> receptor antagonists. NC00080421, with a similar structure to known selective hMT<sub>2</sub> receptor antagonist 4P-PDOT<sup>[29]</sup>, showed approximately 40-fold selectivity at the hMT<sub>2</sub> receptor. However, NC00090702, NC00096199, and NC00099763, with selective hMT<sub>1</sub> receptor antagonist activities, exhibited little structural similarity to melatonin. Because hMT<sub>1</sub> receptor modulators are scarcely available,




**Figure 4.** Dose-response curves of melatonin at hMT<sub>1</sub> (A) and hMT<sub>2</sub> (B) receptors measured by respective GTP-γS assay from which EC<sub>50</sub> values were calculated. Data shown are mean±SD of triplicate measurements and representative of at least 3 independent experiments.



**Figure 5.** Structures of newly-discovered selective melatonin receptor antagonists and melatonin.

our findings certainly provide some new knowledge for continued pursuit in this direction using the dual melatonin receptor subtype assay system<sup>[30]</sup>. Unfortunately, we have not found selective hMT<sub>1</sub> or hMT<sub>2</sub> receptor agonists in this study, but further structure–activity relationship analyses on these leads and relevant structural modifications may





**Figure 6.** Subtype selectivity of 4 newly-discovered melatonin receptor (hMT<sub>1</sub> and hMT<sub>2</sub>) modulators. All compounds show antagonist activities.  $K_B$  values at the hMT<sub>1</sub> receptor were compared with that of hMT<sub>2</sub> (hMT<sub>1</sub>/hMT<sub>2</sub>). (A) agonist feature ( $E_{max}$ ); (B) antagonist feature ( $K_B/I_{max}$ ).

dial in agonist activity.

In humans, it has been suggested that melatonin has a variety of clinical applications, such as the treatment of delayed sleep phase syndrome, jet lag, work shift disturbances<sup>[31]</sup>, seasonal affective disorders<sup>[32]</sup>, and aging<sup>[33]</sup>. Melatonin was also implicated to have immunomodulating properties<sup>[34]</sup>. Recent studies demonstrated that melatonin may play a role in Alzheimer's disease<sup>[35]</sup>, anesthesia<sup>[36]</sup>, depression<sup>[37]</sup>, and obesity<sup>[38]</sup>. However, the very short biological half-life (20–30 min), extensive metabolism, and lack of selectivity of melatonin at its target sites prompt the search for metabolically-stable agonists and antagonists. Although studies involving animal models relevant to human pathologies and appropriate clinical trials have indicated that circadian sleep disorders and insomnia are probably the sole therapeutic indications today for selective hMT<sub>1</sub> or hMT<sub>2</sub> receptor ligands<sup>[39]</sup>, the selective antagonists described in this study may provide

important tools in further understanding the physiological functions of the different receptor subtypes (hMT<sub>1</sub> or hMT<sub>2</sub>) *in vivo*.

## Acknowledgements

We are indebted to Mr Xin HUI, Ms Hong YU, and Mr Qiang SHEN for their technical assistance, and to Dr Dale E MAIS for valuable discussions.

## Author contribution

Ming-wei WANG and Jean A BOUTIN designed research; Jian-hua YAN and Hao-ran SU performed research; Jean A BOUTIN and M Pierre RENARD contributed new analytical tools and reagents; Jian-hua YAN and Ming-wei WANG analyzed data; Ming-wei WANG wrote the paper.

## References

- Reiter RJ. Pineal melatonin: cell biology of its synthesis and of its physiological interactions. *Endocr Rev* 1991; 12: 151–80.
- Lerner AB, Case JD. Structure of melatonin. *J Am Chem Soc* 1959; 81: 6084–5.
- Cassone VM. Melatonin's role in vertebrate circadian rhythms. *Chronobiol Int* 1998; 15: 457–73.
- Krause DN, Geary GG, Doolen S, Duckles SP. Melatonin and cardiovascular function. *Adv Exp Med Biol* 1999; 460: 299–310.
- Reiter RJ. Melatonin and human reproduction. *Ann Med* 1998; 30: 103–8.
- Jones MP, Melan MA, Witt-Enderby PA. Melatonin decreases cell proliferation and transformation in a melatonin receptor-dependent manner. *Cancer Lett* 2000; 151: 133–43.
- Dubocovich ML. Characterization of a retinal melatonin receptor. *J Pharmacol Exp Ther* 1985; 234: 395–401.
- Dubocovich ML, Takahashi JS. Use of 2-[<sup>125</sup>I]iodomelatonin to characterize melatonin binding sites in chicken retina. *Proc Natl Acad Sci USA* 1987; 84: 3916–20.
- Reppert SM, Weaver DR, Ebisawa T. Cloning and characterization of a mammalian melatonin receptor that mediates reproductive and circadian responses. *Neuron* 1994; 13: 1177–85.
- Reppert SM, Godson C, Mahle CD, Weaver DR, Slaugenhaupt SA, Gusella JF. Molecular characterization of a second melatonin receptor expressed in human retina and brain: the Mel1b melatonin receptor. *Proc Natl Acad Sci USA* 1995; 92: 8734–8.
- Hunt AE, Ghoul WM, Dubocovich ML. Activation of melatonin MT<sub>2</sub> receptors in rat suprachiasmatic nucleus phase advances the circadian clock. *Am J Physiol Cell Physiol* 2001; 280: C110–C8.
- Nosjean O, Ferro M, Cogé F, Beauverger P, Henlin JM, Lefoulon F, *et al*. Identification of the melatonin binding site MT<sub>3</sub> as the quinone reductase 2. *J Biol Chem* 2000; 275: 31311–7.
- Nosjean O, Nicolas JP, Klupsch F, Delagrangre P, Canet E, Boutin JA. Comparative pharmacological studies of melatonin receptors: MT<sub>1</sub>, MT<sub>2</sub> and MT<sub>3</sub>/QR<sub>2</sub>. Tissue distribution of MT<sub>3</sub>/QR<sub>2</sub>. *Biochem Pharmacol* 2001; 61: 1369–79.

- 14 Dubocovich ML, Cardinali DP, Delagrange P, Krause DN, Strosberg D, Sugden D, *et al*. The IUPHAR compendium of receptor characterization and classification. IUPHAR Media, London 2000: 270–7.
- 15 Al-Ghoul WM, Herman MD, Dubocovich ML. Melatonin receptor subtype expression in human cerebellum. *Neuroreport* 1998; 9: 4063–8.
- 16 Ting KN, Blaylock NA, Sugden D, Delagrange P, Scalbert E, Wilson VG. Molecular and pharmacological evidence for MT<sub>1</sub> melatonin receptor subtype in tail artery of juvenile Wistar rats. *Br J Pharmacol* 1999; 127: 987–95.
- 17 Mazzucchelli C, Pannacci M, Nonno R, Lucini V, Fraschini F, Stankov BM. The melatonin receptor in the human brain: cloning experiments and distribution studies. *Brain Res Mol Brain Res* 1996; 39: 117–26.
- 18 Weaver DR, Reppert SM. The Mel1a melatonin receptor gene is expressed in human suprachiasmatic nuclei. *Neuroreport* 1996; 8: 109–12.
- 19 Song Y, Chan CW, Brown GM, Pang SF, Silverman M. Studies of the renal action of melatonin: evidence that the effects are mediated by 37 kDa receptors of the Mel1a subtype localized primarily to the basolateral membrane of the proximal tubule. *FASEB J* 1997; 11: 93–100.
- 20 Delagrange P, Guardiola-Lemaitre B. Melatonin, its receptors and relationships with biological rhythms disorders. *Clin Neuropharmacol* 1997; 20: 482–510.
- 21 Mahle CD, Takaki KS, Watson AJ. Melatonin receptor ligands and their potential clinical applications. *Annu Rep Med Chem* 1997: 31–40.
- 22 Wu B, Gao J, Wang MW. Development of a complex scintillation proximity assay for high-throughput screening of PPAR $\gamma$  modulators. *Acta Pharmacol Sin* 2005; 26: 339–44.
- 23 Harrison C, Traynor JR. The [<sup>35</sup>S]GTP $\gamma$ S binding assay: approaches and applications in pharmacology. *Life Sci* 2003; 74: 489–508.
- 24 Audinot V, Mailliet F, Lahaye-Brasseur C, Bonnaud A, Gall AL, Amossé C, *et al*. New selective ligands of human cloned melatonin MT<sub>1</sub> and MT<sub>2</sub> receptors. *Naunyn Schmiedeberg's Arch Pharmacol* 2003; 367: 553–61.
- 25 Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976; 72: 248–54.
- 26 Cheng YC, Prussoff WH. Relationship between the inhibition constant ( $K_i$ ) and the concentration of inhibitor which causes 50 per cent inhibition (IC<sub>50</sub>) of an enzymatic reaction. *Biochem Pharmacol* 1973; 22: 3099–108.
- 27 Cohen FR, Lazareno S, Birdsall NJM. The effects of saponin on the binding and functional properties of the human adenosine A1 receptor. *Br J Pharmacol* 1996; 117: 1521–9.
- 28 Zhang J-H, Chung TDY, Oldenburg KR. A simple statistical parameter for use in evaluation and validation of high throughput screening assays. *J Biomol Screen* 1999; 4: 67–73.
- 29 Dubocovich ML, Masana MI, Iacob S, Sauri DM. Melatonin receptor antagonists that differentiate between the human Mel1a and Mel1b recombinant subtypes are used to assess the pharmacological profile of the rabbit retina ML1 presynaptic heteroreceptor. *Naunyn-Schmiedeberg's Arch Pharmacol* 1997; 355: 365–75.
- 30 Silvia R, Simone L, Marco M, Pier VP, Gilberto S, Annalida B, *et al*. Analysis of structure-activity relationships for MT<sub>2</sub> selective antagonists by melatonin MT<sub>1</sub> and MT<sub>2</sub> receptor models. *J Med Chem* 2005; 48: 4049–60.
- 31 Lewy AJ, Emens J, Jackman A, Yuhas K. Circadian uses of melatonin in humans. *Chronobiol Int* 2006; 23: 403–12.
- 32 Srinivasan V, Smits M, Spence W, Lowe AD, Kayumov L, Pandi-Perumal SR, *et al*. Melatonin in mood disorders. *World J Biol Psychiatry* 2006; 7: 138–51.
- 33 Poeggeler B. Melatonin, aging, and age-related diseases: perspectives for prevention, intervention, and therapy. *Endocrine* 2005; 27: 201–12.
- 34 Srinivasan V, Maestroni G, Cardinali D, Esquifino A, Perumal SP, Miller S. Melatonin, immune function and aging. *Immun Ageing* 2005; 2: 17.
- 35 Brunner P, Sozer-Topcular N, Jockers R, Ravid R, Angeloni D, Fraschini F, *et al*. Pineal and cortical melatonin receptors MT<sub>1</sub> and MT<sub>2</sub> are decreased in Alzheimer's disease. *Eur J Histochem* 2006; 50: 311–6.
- 36 Naguib M, Gottumukkala V, Goldstein PA. Melatonin and anesthesia: a clinical perspective. *J Pineal Res* 2007; 42: 12–21.
- 37 Banki MC. Agomelatin: the first “melatoninergetic” antidepressant. *Neuropsychopharmacol Hung* 2006; 8: 105–12.
- 38 Hussein MR, Ahmed OG, Hassan AF, Ahmed MA. Intake of melatonin is associated with amelioration of physiological changes, both metabolic and morphological pathologies associated with obesity: an animal model. *Int J Exp Pathol* 2007; 88: 19–29.
- 39 Delagrange P, Boutin JA. Therapeutic potential of melatonin ligands. *Chronobiol Int* 2006; 23: 413–8.