

superoxide dismutase to study endothelium-dependent and nitric oxide-elicited vascular smooth muscle relaxation. *J Pharmacol Exp Ther* 1988; **244**: 181-9.

8 Li YJ, Duckles SP. Presence of endothelium masks direct vasodilator effects of pyrogallol and methylthionium chloride in perfused rat mesenteric artery. *Acta Pharmacol Sin* 1993; **15**: 21-7.

9 Li YJ, Duckles SP. Effect of endothelium on the actions of sympathetic and sensory nerves in the perfused rat mesentery. *Eur J Pharmacol* 1992; **210**: 23-30.

10 Wolin MS, Cherry PD, Rodenburg JM, Messina EJ, Kaley G. Methylene blue inhibits vasodilation of skeletal muscle arterioles to acetylcholine and nitric oxide via the extracellular generation of superoxide anion. *J Pharmacol Exp Ther* 1990; **254**: 872-6.

11 Wood KS, Buga GM, Byrns RE, Ignarro LJ. Vascular smooth muscle-derived relaxing factor (MDRF) and its close similarity to nitric oxide. *Biochem Biophys Res Commun* 1990; **170**: 80-8.

12 Bult H, Boeckxstaens GE, Pelckmans PA, Jordaens FH, Van Maercke YM, Herman AG. Nitric oxide as an inhibitory non-adrenergic non-cholinergic neurotransmitter. *Nature* 1990; **345**: 346-7.

13 Chen X, Hu YC, Gillis CN. Free radicals make vasodilator(s) release from aortic endothelial and smooth muscle cells. *Chinese J Pharmacol Toxicol* 1991; **5**: 269-72.

14 Gustafsson LE, Wiklund CU, Wiklund NP, Persson MG, Moncada S. Modulation of autonomic neuroeffector

transmission by nitric oxide in guinea pig ileum. *Biochem Biophys Res Commun* 1990; **173**: 106-10.

15 Tate RM, Morris HG, Schroeder WR, Repine JE. Oxygen metabolites stimulate thromboxane production and vasoconstriction in isolated saline-perfused rabbit lungs. *J Clin Invest* 1984; **74**: 608-13.

142-147

12

邻苯三酚与亚甲蓝舒张大鼠肠系膜动脉的不同机制¹

R965.2

李元建², Sue Piper DUCKLES (Department of Pharmacology, College of Medicine, University of California, Irvine CA 92717, USA)

A 摘要 在去内皮肠系膜血管, 邻苯三酚与亚甲蓝产生浓度依赖性舒张, SOD 可取消邻苯三酚对 TNS 所致缩血管效应的抑制作用, 但不影响亚甲蓝的效应. 邻苯三酚与亚甲蓝的舒血管效应不被过氧化氢酶、去铁胺、吲哚美辛和辣椒素所影响. 结果提示, 邻苯三酚的舒血管效应是其产生超氧阴离子所致, 而亚甲蓝的效应为直接作用于血管平滑肌.

关键词 肠系膜动脉; 邻苯三酚; 亚甲蓝; 亚硝基精氨酸甲酯; 超氧化物歧化酶; 吲哚美辛; 辣椒素; 过氧化氢酶; 去铁胺

Effects of toosendanin on electric and mechanical properties of guinea pig papillary muscles¹

GAO Xiao-Dong, TANG Shu-Ben², LÜ Jian (Department of Physiology, He-nan Medical University, Zhengzhou 450052, China) SHI Yu-Liang (Shanghai Institute of Physiology, Chinese Academy of Sciences, Shanghai 200031, China)

ABSTRACT Effects of toosendanin (TS) on the action potentials and contractile force in guinea pig papillary muscles were examined using a standard microelectrode technique.

TS concentration-dependently increased the action potential duration at 90% repolarization (APD₉₀) of the fast action potentials. In the presence of a I_{K1} channel blocker BaCl₂, the effects of TS on lengthening the APD₉₀ were completely abolished, thereby suggesting that TS inhibited the inward rectifier K⁺ current I_{K1}. The APD and contractile force of amino-

Received 1992-11-02 Accepted 1993-09-03
¹ Project supported by the National Natural Science Foundation of China, No 38970831.
² Correspondence should be addressed to TANG Shu-Ben.

phylline-induced slow action potentials were potentiated by TS in a concentration-dependent manner. In the presence of BaCl_2 , both effects of TS were completely abolished. The effect of TS on enhancing contractile force was abolished by the addition of CdCl_2 , with the prolongation of APD preserved. Thus, TS selectively inhibited the inward rectifier K^+ current I_{K1} with a positive inotropic effect, resulting from a delay in Ca channel inactivation which was secondary to delay in ventricular repolarization.

KEY WORDS toosendanin; papillary muscles; action potentials; myocardial contraction; ion channels

Toosendanin (TS) is a triterpenoid derivative ($\text{C}_{30}\text{H}_{38}\text{O}_{11}$) extracted from the bark of *Melia toosendan* Seib et Zucc, which has been elucidated in structure¹⁾ and used as an ascaris vermifuge in China. TS influenced not only quantal but also non-quantal release of acetylcholine²⁾. As an effective anti-botulinic agent, TS showed an extracellular Ca^{2+} dependent initial facilitative action on miniature end-plate potentials (MEPP) discharge before the inhibition of MEPP³⁾. The effects of TS on the Ca channels may be attributed to an increase in calcium conductance⁴⁾. But there were no report available to show the effects of TS on the cardiac ion channels. Hence, using a standard microelectrode technique, we investigated the ionic mechanisms of the effect of TS in isolated guinea pig papillary muscles.

MATERIALS AND METHODS

Reagents The TS used were crystals from ethanol, >98% in purity, the same as described previously²⁾, and gifted from Prof LI Pei-Zhong, Academy of Military Medical Sciences; aminophylline (Shanghai XinYi Pharmaceutical Factory, lot 881006); BaCl_2 (Tianjin 3rd Chemical Reagent Factory, lot 820912);

CdCl_2 (Beijing 57601 Chemical Engineering Factory, lot 850827).

Protocol Guinea pigs (either sex) weighing 250 ± 50 g were stunned. Papillary muscle was taken from the right ventricle and was pinned in a 2-ml chamber. For recording fast action potentials (FAP), the Krebs solution (NaCl 135, KCl 5.0, CaCl_2 2.0, MgCl_2 1.0, NaH_2PO_4 2.2, NaHCO_3 11.9, and glucose $5.5 \text{ mmol}\cdot\text{L}^{-1}$) was gassed with 95% O_2 + 5% CO_2 (34 ± 0.5 °C, pH 7.4 ± 0.5). Bath solution was circulated at a rate of $11 \text{ ml}\cdot\text{min}^{-1}$. Force of contraction (FC) was measured with an isometric force transducer (Model HN-79, Shanghai Institute of Physiology, Chinese Academy of Sciences) which was connected to the tendinous end of the preparation. Initial length was adjusted to yield an optimal contraction amplitude upon stimulation. Transmembrane action potentials were recorded with conventional glass microelectrodes filled with KCl $3 \text{ mol}\cdot\text{L}^{-1}$ having a resistance of 15–25 M Ω . The maximal upstroke velocity (V_{max}) was obtained by an electronic differentiator (Department of Physiology, Beijing University). Stimulation parameters were: stimulus duration 1 ms, frequency 1 Hz, voltage 20% above threshold. After recording the control FAP for 1 h, data were obtained cumulatively by increasing the concentration of TS every 20 min.

For slow action potentials (SAP), high K^+ ($22 \text{ mmol}\cdot\text{L}^{-1}$) Krebs solution was prepared by equimolar substitution of KCl for NaCl , containing aminophylline $0.75 \text{ mmol}\cdot\text{L}^{-1}$ to elicit SAP after depolarization of the resting potential and inactivation of fast Na^+ channels. Stimulation parameters were: stimulus duration 3 ms, frequency 0.5 Hz, voltage 20% above threshold. Experiments started after the recording of control SAP for 1 h.

Solutions of TS were prepared in either normal ($\text{K}^+ 5 \text{ mmol}\cdot\text{L}^{-1}$) or high K^+ ($\text{K}^+ 22 \text{ mmol}\cdot\text{L}^{-1}$) Krebs solutions and were diluted to the final concentrations of 5, 10, 20 $\mu\text{g}\cdot\text{ml}^{-1}$ (pH 7.4). The electric and mechanical signals were displayed on the screen of an oscilloscope and photographed at regular intervals. The effects of TS on the FAP and SAP parameters (V_{max} , action potential amplitude APA, APD), and FC were observed. Data were analyzed for statistical significance by *t* test for paired values.

RESULTS

With TS at concentrations of $\leq 5 \mu\text{g} \cdot \text{ml}^{-1}$, no effects on FAP parameters were seen. At higher concentrations, TS caused a concentration-dependent prolongation in the APD, with an increase at APD_{90} during the terminal phase of repolarization, while the V_{max} and APA were practically unaffected (Tab 1, Fig 1).




Fig 1. Effect of toosendanin on fast action potential (FAP) configuration in guinea pig papillary muscles. A: control; B: in the presence of toosendanin $20 \mu\text{g} \cdot \text{ml}^{-1}$, duration of FAP at 90 % repolarization was somewhat lengthened.

In another 6 papillary muscles, after recording the control FAP for 1 h, the addi-

tion of BaCl_2 $50 \mu\text{mol} \cdot \text{L}^{-1}$ caused a depolarization of the resting potential from $68 \pm 3 \text{ mV}$ to $66 \pm 3 \text{ mV}$ ($n=5$, $P < 0.01$), a reduction of the APA from $103 \pm 3 \text{ mV}$ to $100 \pm 4 \text{ mV}$ ($n=6$, $P < 0.01$), and a prolongation of APD_{90} from $185 \pm 43 \text{ ms}$ to $220 \pm 29 \text{ ms}$ ($n=6$, $P < 0.01$). Similar results have also been reported^{15,16}, in which BaCl_2 was used to block I_{K1} . In the presence of BaCl_2 , the effects of TS on lengthening APD was completely abolished (Tab 1).

With TS $\leq 5 \mu\text{g} \cdot \text{ml}^{-1}$, no significant effects on the aminophylline-induced SAP parameters were seen. At higher concentrations, TS concentration-dependently prolonged the APD, without effecting the V_{max} and APA (Tab 2). TS (10 and $20 \mu\text{g} \cdot \text{ml}^{-1}$) potentiated the accompanying FC of the aminophylline - induced SAP concentration - dependently (Tab 2, Fig 2). TS $20 \mu\text{g} \cdot \text{ml}^{-1}$ increased the FC to $173 \pm 42 \%$ of the control ($n=6$, $P < 0.01$).

In another 6 papillary muscles, after recording the aminophylline-induced SAP for 1 h, the addition of BaCl_2 $70 \mu\text{mol} \cdot \text{L}^{-1}$ caused an

Tab 1. Effects of toosendanin on fast action potentials in guinea pig papillary muscles. $n=6$, $\bar{x} \pm s$. * $P > 0.05$, ^b $P < 0.05$, ^c $P < 0.01$ vs control. ^d $P > 0.05$ vs control + BaCl_2 $50 \mu\text{mol} \cdot \text{L}^{-1}$.

Toosendanin/ $\mu\text{g} \cdot \text{ml}^{-1}$	$V_{\text{max}} /$ $\text{V} \cdot \text{s}^{-1}$	APA/ mV	$\text{APD}_{30} /$ ms	$\text{APD}_{50} /$ ms	$\text{APD}_{90} /$ ms
0	195 ± 36	99 ± 6	102 ± 21	135 ± 25	163 ± 27
5	185 ± 29^a	101 ± 3^a	106 ± 19^a	138 ± 25^a	165 ± 27^a
10	180 ± 39^a	102 ± 4^a	108 ± 18^a	143 ± 25^c	171 ± 28^c
20	181 ± 54^a	101 ± 3^a	112 ± 19^b	150 ± 22^c	180 ± 24^c
0	167 ± 32	103 ± 3	109 ± 31	154 ± 40	185 ± 43
+ BaCl_2	172 ± 35^d	100 ± 4^d	116 ± 20^d	167 ± 29^d	220 ± 29^d
5	173 ± 35^d	100 ± 4^d	115 ± 23^d	167 ± 31^d	222 ± 31^d
10	179 ± 36^d	100 ± 4^d	117 ± 19^d	168 ± 29^d	222 ± 27^d
20	169 ± 35^d	101 ± 4^d	118 ± 16^d	172 ± 22^d	228 ± 23^d

V_{max} = maximal upstroke velocity; APA = action potential amplitude; APD_{30} , APD_{50} , and APD_{90} = action potential duration at 30 %, 50, and 90 %, repolarization.

Tab 2. Effects of toosendanin on slow action potentials induced by aminophylline in guinea pig papillary muscles. $\bar{x} \pm s$. ^a $P > 0.05$, ^b $P < 0.05$, ^c $P < 0.01$ vs control ($n = 6$). ^d $P > 0.05$, ^e $P < 0.05$, ^f $P < 0.01$ vs control + BaCl₂ 70 $\mu\text{mol} \cdot \text{L}^{-1}$ ($n = 5$) or vs control + CdCl₂ 50 $\mu\text{mol} \cdot \text{L}^{-1}$ ($n = 8$).

Toosendanin/ $\mu\text{g} \cdot \text{ml}^{-1}$	$V_{\text{max}} /$ $\text{V} \cdot \text{s}^{-1}$	APA/ mV	APD ₅₀ / ms	APD ₉₀ / ms	FC/ %
0	9.4 ± 4.0	68 ± 3	125 ± 27	145 ± 28	100
5	10.6 ± 6.8 ^a	68 ± 3 ^a	128 ± 34 ^a	149 ± 34 ^a	111 ± 30 ^a
10	12.3 ± 7.6 ^a	69 ± 4 ^a	135 ± 36 ^b	156 ± 38 ^b	147.8 ± 52.6 ^b
20	12.7 ± 6.3 ^a	69 ± 4 ^a	145 ± 31 ^b	167 ± 33 ^c	173 ± 42 ^c
<hr/>					
0	11.3 ± 4.7	68 ± 3	149 ± 33	180 ± 34	100
+ BaCl ₂	14 ± 4 ^a	70 ± 4 ^c	193 ± 41 ^c	236 ± 45 ^e	93 ± 28 ^c
5	13.5 ± 4.6 ^d	70 ± 4 ^d	192 ± 38 ^d	236 ± 43 ^d	99 ± 2 ^d
10	13.5 ± 4.6 ^d	70 ± 4 ^d	193 ± 39 ^d	236 ± 42 ^d	112 ± 23 ^d
20	15 ± 4 ^d	70 ± 4 ^d	191 ± 41 ^d	237 ± 43 ^d	126.6 ± 58.3 ^d
<hr/>					
0	12.9 ± 4.6	78 ± 8	122 ± 38	174 ± 56	100
+ CdCl ₂	10.9 ± 5.1 ^a	67 ± 7 ^c	113 ± 36 ^a	177.5 ± 72.0 ^a	22.4 ± 9.7 ^c
5	10.5 ± 4.9 ^d	66 ± 6 ^d	117 ± 28 ^d	179 ± 66 ^d	99 ± 12 ^d
10	10.7 ± 4.7 ^d	67 ± 5 ^d	123 ± 28 ^c	191.2 ± 66.4 ^f	98 ± 6 ^d
20	10.7 ± 4.4 ^d	67 ± 5 ^d	126 ± 27 ^c	201 ± 63 ^f	105 ± 20 ^c




Fig 2. Effects of toosendanin on aminophylline - induced slow action potentials (SAP, upper tracing) and force of contraction (FC, lower tracing) in guinea pig papillary muscles. Duration of SAP was lengthened and FC increased, in a concentration - dependent manner.

increase of the APA from 68 ± 3 mV to 70 ± 4 mV ($n = 5$, $P < 0.01$), a prolongation of APD₅₀ and APD₉₀ from 149 ± 33 and 180 ± 34 ms to 193 ± 41 and 236 ± 45 ms, respectively ($n = 5$, $P < 0.01$), without effecting the FC. The effects of TS (10 and 20 $\mu\text{g} \cdot \text{ml}^{-1}$) on lengthening APD and enhancing FC were completely abolished when BaCl₂ was used to block the I_{K1} (Tab 2).

In another 6 experiments, after recording the aminophylline-induced SAP for 1 h, the

addition of CdCl₂ 50 $\mu\text{mol} \cdot \text{L}^{-1}$ caused a reduction of APA from 78 ± 8 mV to 67 ± 7 mV ($n = 8$, $P < 0.01$), an attenuation of FC to 22.4 ± 9.7 % of the control ($n = 8$, $P < 0.01$), without effecting the V_{max} and APD. The effect of TS on enhancing FC was abolished when CdCl₂ was used to reduce the I_{K1} , whereas the concentration-dependent prolongation of the APD persisted with increasing concentrations of TS (Tab 2, Fig 3).

DISCUSSION

Our results indicated that the main effect of TS was to inhibit the inward rectifier K⁺ current I_{K1} . Suppressing both the inward rectifier current and the delayed rectifier current by TS could lead to prolongation of the APD, but the primary currents blocked by TS were I_{K1} , since the current intensity of I_{K} was less than that of I_{K1} and APD at voltages of 0 mV or above did not increase substantially^{1,7,8}. Thus, the I_{K} system appeared to play an insignificant role, at least in the case of the TS -




Fig 3. I_{K1} -blocking effect of $CdCl_2$ on toosendanin-induced prolongation of slow action potential duration (upper tracing) and potentiation of force of contraction (FC, lower tracing) in guinea pig papillary muscles. A: control slow action potential (SAP) induced by aminophylline and the accompanying contraction; B: after the addition of $CdCl_2$, $50 \mu mol \cdot L^{-1}$, FC was suppressed and without much effect on the duration of SAP; C: in the presence of Cd^{2+} , effect of toosendanin on enhancing FC was abolished, with the prolongation of APD preserved.

induced prolongation of APD in multicellular preparations.

The APD of aminophylline-induced SAP and FC were increased by TS. Following the application of $BaCl_2$ to block I_{K1} , both the effects of TS on lengthening APD and enhancing FC could be completely abolished. Addition of $CdCl_2$ abolished the effects of TS on enhancing FC, while the APD-prolonging effects of TS were preserved. Thus, the positive inotropic effects of TS on the accompanying FC of the SAP were not likely to be due to increased Ca^{2+} influx into the cells at depolarization phase of the SAP (no changes in V_{max} occurred), but resulting from slowing the calcium channel inactivation secondary to the delay in ventricular repolarization.

REFERENCES

1 Shu GX, Liang XT. A correction of the structure of chuanjiansu. Acta Chim Sin 1980; 38: 196-8.
 2 Shih YL. Abolishment of non-quantal release of acetylcholine from the mouse phrenic nerve endings by toosendanin.

Jap J Physiol 1986; 36: 601-5.
 3 Shih YL, Hsu K. Anti-botulismic effect of toosendanin and its facilitatory action on miniature end-plate potentials. Jap J Physiol 1983; 33: 677-80.
 4 Shih YL, Furuya K. Calcium conductance produced by toosendanin in neuroblastoma X glioma hybrid cells. Proceedings of Shanghai Society for Physiological Sciences CAPS News Communication 1991; 10 suppl 6: S9-S10.
 5 Hirano Y, Hiraoka M. Changes in K^+ currents induced by Ba^{2+} in guinea pig ventricular muscles. Am J Physiol 1986; 251: H24-H33.
 6 Imoto Y, Ehara T, Matsuura H. Voltage- and time-dependent block of i_{K1} underlying Ba^{2+} -induced ventricular automaticity. Am J Physiol 1987; 252: H325-H333.
 7 Martin CL, Chinn K. Contribution of delayed rectifier and inward rectifier to repolarization of the action potential: pharmacologic separation. J Cardiovasc Pharmacol 1992; 19: 830-7.
 8 Shimoni Y, Clark RB, Giles WR. Role of an inwardly rectifying potassium current in rabbit ventricular action potential J Physiol (Lond) 1992; 448: 709-27.

147-151

川楝素对豚鼠乳头状肌电和机械特性的影响

高晓东, 汤树本, 吕健 (河南医科大学生理教研室, 郑州450052, 中国) 施玉慧 (中国科学院上海生理研究所, 上海200031, 中国)

A 摘要 川楝素(TS)浓度依赖性地使快反应电位复极至90%的时程(APD_{90})延长。用 $BaCl_2$ 阻断 I_{K1} , 可取消 TS 延长 APD_{90} 的作用。TS 使慢反应电位的 APD 延长和收缩力(FC)增强。用 $BaCl_2$ 后, 可取消 TS 的上述作用。 $CdCl_2$ 取消 TS 增强 FC 的作用, 但延长 APD 的作用存在。提示, TS 抑制 I_{K1} , 其正性肌力作用是继发于 APD 的延长及钙通道的失活减慢。

关键词 川楝素; 乳头状肌; 动作电位; 心肌收缩力; 离子通道

豚鼠

13

R965.2