

©2004, Acta Pharmacologica Sinica  
Chinese Pharmacological Society  
Shanghai Institute of Materia Medica  
Chinese Academy of Sciences  
<http://www.ChinaPhar.com>

## Effects of 15-deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub> on cell proliferation and apoptosis in ECV304 endothelial cells<sup>1</sup>

Yu-gang DONG<sup>2</sup>, Dan-dan CHEN, Jian-gui HE, Yong-yuan GUAN<sup>3</sup>

*Department of Cardiology, First Affiliated Hospital, Zhongshan University, Guangzhou 510080;*

*<sup>3</sup>Department of Pathology, Medical College of Zhongshan University, Guangzhou 510089, China*

**KEY WORDS** peroxisome proliferator-activated receptor; atherosclerosis; endothelial cells; apoptosis; transcription factor AP-1; NF-kappa B; prostaglandins

### ABSTRACT

**AIM:** To investigate the effects of 15-deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub> (15d-PGJ<sub>2</sub>) on cell proliferation and apoptosis in ECV304 endothelial cells and related molecular mechanism. **METHODS:** MTT, Hoechst33258, TUNEL, Flow cytometry, DNA ladder, RT-PCR, Western blot, and electrophoretic mobility shift assay (EMSA) analysis were employed. **RESULTS:** The 15d-PGJ<sub>2</sub> induced apoptosis in ECV304 endothelial cells in a dose-dependent manner (the percentage of apoptosis was enhanced from 10.0 %±1.3 % to 32.8 %±1.6 %), which was accompanied by inhibition of NF- $\kappa$ B and AP-1 DNA binding activity, down-regulation of *c-myc*, upregulation of *Gadd45* and *p53*, and activation of p38 kinase. However, the expression of *p21* was found no significant change. **CONCLUSION:** peroxisome proliferator-activated receptor gamma ligand, 15d-PGJ<sub>2</sub>, can inhibit proliferation and induce apoptosis in ECV304 endothelial cells through different mechanisms.

### INTRODUCTION

Atherosclerosis is a multifactorial disease in which the occurrence of lesions may result in ischemia of the heart, brain, and extremities. Nevertheless, the mechanism of atherosclerosis remains unclear. Endothelial damage or dysfunction is considered the initiating cause of atherogenesis. Recent studies have shown that increased apoptosis of endothelial cell (EC) has been observed in atheromatous plaques<sup>[1]</sup>. To observe the change of EC apoptotic pathways in atherosclerosis may enable a greater understanding of disease pathogenesis.

Expression of peroxisome proliferator-activated receptor (PPAR) has been shown in atherosclerotic lesions and macrophage foam cells, which suggests that PPAR may affect atherosclerogenic process<sup>[2]</sup>. PPAR form a subfamily of the nuclear receptor superfamily, and three isoforms of PPAR have been identified thus far: PPAR $\alpha$ , PPAR $\delta$ , and PPAR $\gamma$ . The PPAR is ligand-dependent transcription factor that regulate target gene expression by binding to specific peroxisome proliferator response elements (PPRE) in enhancer sites of regulated genes. Each receptor binds to its PPRE as a heterodimer with a retinoid X receptor (RXR). Upon binding an agonist, the conformation of a PPAR is altered and stabilized such that a binding cleft is created and recruitment of transcriptional coactivators occurs. The result showed an increase in gene transcription<sup>[3]</sup>. Activation of peroxisome proliferator-activated receptors induces apoptosis of endothelial cells, human mono-

<sup>1</sup> Project supported by the Provincial Science Foundation of Guangdong, China (2000).

<sup>2</sup> Correspondence to Yu-gang DONG. Phn 86-20-8775-5766, ext 8151. E-mail [yugangdong@gzsums.edu.cn](mailto:yugangdong@gzsums.edu.cn)

Received 2002-11-22

Accepted 2003-04-30

cyte-derived macrophages and smooth muscle cells<sup>[4-6]</sup>. Transcription of affected genes may also indirectly modulated by PPAR via interference with other transcription factor pathways. Activation of PPAR negatively interferes with the NF- $\kappa$ B, a signal transducer and activator of transcription (STAT), and Activator protein 1 (AP-1) pathways<sup>[5-7]</sup>. 15-Deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub> (15d-PGJ<sub>2</sub>) has been demonstrated to be a natural ligand of the PPAR $\gamma$ . It has been reported that 15d-PGJ<sub>2</sub> induced endothelial apoptosis<sup>[4]</sup>, however the mechanisms have been unclear yet. The aim of the present work was to measure the effects of 15d-PGJ<sub>2</sub> on cell proliferation and apoptosis in ECV304 endothelial cells, and further investigate its possible molecular mechanisms.

## MATERIALS AND METHODS

**Drugs and reagents** RPMI-1640 medium, RNase A, agarose gel, 1 kb DNA ladder, proteinase K, propidium iodide (PI), and MTT were purchased from Gibco Co. The 15d-PGJ<sub>2</sub> was purchased from Cayman Chemical Co. Other reagents, antibodies and reagent boxes were purchased from Sigma Chemical Co.

**Cell culture** The ECV304 human endothelial cell line was purchased from China Center for Type Culture Collection (CCTCC). The cell line was cultured in RPMI-1640 medium containing 10 % fresh bovine serum, benzylpenicillin (100 kU/L), and kanamycin (0.1 g/L) at 37 °C in 5 % CO<sub>2</sub>-95 % air atmosphere.

**MTT assay** ECV304 human endothelial cells growing on 96-well plates were treated with 15d-PGJ<sub>2</sub> (5-20  $\mu$ mol/L) for 48 h, and untreated cells served as a control. Ten  $\mu$ L of the stock solution (2.5 g/L) of 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl-tetrazolium bromide (MTT) was added to each well. After 1 h of incubation at 37 °C, the medium was removed, 50  $\mu$ L of the extraction buffer (10 % Triton-X100; HCl 0.1 mol/L) was added, and plates were gently shaken for 30 min at room temperature. The optical densities were measured at 570 nm.

**Hoechst 33258 staining** Cells were fixed with 4 % formaldehyde in phosphate buffered saline (PBS) for 10 min, stained by Hoechst 33258 for 1 h, and were subjected to fluorescence microscopy. Apoptotic cells were identified by nuclear condensation and/or fragmentation.

**TUNEL assay** TUNEL assay was performed by the apoptosis detection system. ECV304 cells were

fixed by 4 % paraformaldehyde in PBS overnight at 4 °C, and were washed three times with PBS and then were permeabilized by 0.2 % Triton X-100 in PBS for 15 min on ice. After washing twice, cells were equilibrated at room temperature for 15-30 min in equilibration buffer (potassium cacodylate 200 mmol/L, dithiothreitol 0.2 mmol/L, bovine serum albumin 0.25 g/L, and cobalt chloride 2.5 mmol/L in Tris-HCl 25 mmol/L, pH 6.6) and then incubated in the presence of fluorescein-12-dUTP 5  $\mu$ mol/L, dATP 10  $\mu$ mol/L, edetic acid 100  $\mu$ mol/L, and terminal deoxynucleotidyl transferase at 37 °C for 1.5 h in dark. The tailing reaction was terminated by standard saline citrate (SSC). The samples were washed three times with PBS and were analyzed by fluorescence microscopy. At least  $1 \times 10^3$  cells were counted, and the percentage of TUNEL-positive cells was determined.

**Flow cytometry** For DNA content analysis, 15d-PGJ<sub>2</sub> (5-20  $\mu$ mol/L) was added to ECV304 endothelial cells in mid-logarithmic phase ( $1 \times 10^9$  cells/L). Cells ( $1 \times 10^6$ ) were harvested at indicated time, pelleted, washed with phosphate-buffered saline (PBS), and re-suspended in PBS containing PI 20 mg/L and RNase A 1 mg/L. Fixed cells ( $1 \times 10^6$ ) were examined per each experimental condition by flow cytometry, and the percentage of degraded DNA was determined by the number of cells displaying subdiploid (sub-G<sub>1</sub>). DNA divided by the total number of cells was examined. Cell cycle analysis was performed in the same experimental conditions and distributions used the CellFit program. All measurements were carried out under the same instrumental settings.

**Ladder detection assay** After induction of apoptosis, cells ( $5 \times 10^6$  per sample, both attached and detached cells) were lysed with 150  $\mu$ L hypotonic lysis buffer (edetic acid 10 mmol/L, 0.5 % Triton X-100, Tris-HCl, pH 7.4) for 15 min on ice and were precipitated with 2.5 % polyethylene glycol and NaCl 1 mol/L for 15 min at 4 %. After centrifugation at  $25\,000 \times g$  for 10 min at room temperature, the supernatant was incubated in the presence of proteinase K (0.3 g/L) at 37 °C for 1 h and precipitated with isopropanol at -20 °C. After centrifugation, each pellet was dissolved in 10  $\mu$ L of Tris-edetic acid (pH 7.6) and electrophoresed on a 1.5 % agarose gel containing ethidium bromide. Ladder formation of oligonucleosomal DNA was detected under ultraviolet light.

**Western blot analysis** The cells were lysed in lysis buffer Hepes 25 mmol/L, 1.5 % Triton X-100, 1 %

sodium deoxycholate, 0.1 % SDS, NaCl 0.5 mol/L, edetic acid 5 mmol/L, NaF 50 mmol/L, sodium vanadate 0.1 mmol/L, phenylmethylsulfonyl fluoride (PMSF) 1 mmol/L, and leupeptin 0.1 g/L (pH 7.8) at 4 °C with sonication. The lysates were centrifuged at 20 000×*g* for 15 min and the concentration of the protein in each lysate was determined with Coomassie brilliant blue G-250. Loading buffer (Tris-HCl 42 mmol/L, 10 % glycerol, 2.3 % SDS, 5 % 2-mercaptoethanol and 0.002 % bromophenol blue) was then added to each lysate, and then the samples were electrophoresed on a SDS-polyacrylamide gel. Proteins were transferred to nitrocellulose and incubated sequentially with antibodies against *c-fos*, *c-jun*, IκB, *c-myc*, *p53*, *Gadd45*, and *p21*, and then with peroxidase-conjugated secondary antibodies in the second reaction. Detection was performed with enhanced chemiluminescence reagent.

**RT-PCR** Total RNA was extracted from cells using TRIzol™. RT-PCR for *c-myc*, *p21*, *p53*, *Gadd45*, and *beta-actin* mRNA was performed as described<sup>[7]</sup>.  
*beta-actin*: sense 5' ATCTGGCACCACCTTCTACAATGAG-CTGCG-3',  
 antisense 5' CGTCATACTCCTGCTTGCTGATCCACATCTGC-3'.  
*c-myc*: sense 5' ATCTGGCACACCTTCTACAATGAGCTGCG 3'  
 antisense 5' CCCCTCAGTGGTCTTCCCTAC 3'  
*p21*: sense 5' ATGACTGAGTATAAACTTGTGG 3'  
 antisense 5' TCACATGACTATACACCTTGTC 3'  
*p53*: sense 5' ATGGAGGATTCACAGTCGGA 3'  
 antisense 5' TCAGTCTGAGTCAGGCCCA 3'  
*Gadd45*: sense 5' ATGACTTTGGAGGAATTCTCGG 3'  
 antisense 5' TCACCGTTCGGGAGATTAATC 3'

#### Electrophoretic mobility shift assay (EMSA)

Nuclear extracts were prepared from ECV304 cells treated with 15d-PGJ<sub>2</sub>. Synthetic double-strand oligonucleotides of consensus NF-κB binding sequence, GATCCCAACGGCAGGGGA, and the oligonucleotide of consensus AP-1 binding sequence, CGCTTGATGAGTCAGCCGAA, were end-labeled with γ[<sup>32</sup>P]ATP using T4 polynucleotide kinase. Nuclear extract was incubated with the labeled probe in the presence of poly (dI-dC) in a binding buffer containing 20 mmol/L *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid at room temperature for 30 min. For supershift assays, a total of 0.2 μg of antibodies against p65 subunit of NF-κB were included in the reaction. DNA-protein complexes were resolved by electrophoresis in a 5 % non-denaturing polyacrylamide gel, which was dried and visualized

by autoradiography.

**P38 activity assays**<sup>[8]</sup> P38 MAP kinase activity assays were carried out using an upstate biotechnology MAPKAP kinase 2 immunoprecipitation-kinase assay kit, according to the manufacturer's instructions and using [<sup>32</sup>P]ATP (Amersham UK).

**Statistical analysis** The data were mean values of at least three different experiments and expressed as mean±SD. Student's *t*-test was used to compare data. *P*<0.05 was considered statistically significant.

## RESULTS

**Effect of 15d-PGJ<sub>2</sub> on cell proliferation** The 15d-PGJ<sub>2</sub> induced a dose-dependent inhibition of ECV304 endothelial cell growth in the range of 5-20 μmol/L. Compared with the control group, greater than 50 % mean inhibition was achieved at 15d-PGJ<sub>2</sub> 20 μmol/L (Fig 1).





Fig 1. Effect of 15d-PGJ<sub>2</sub> on proliferation in ECV304 endothelial cells. (1): control; (2-4): 15d-PGJ<sub>2</sub> 5, 10, 20 μmol/L. *n*=3. Mean±SD. <sup>c</sup>*P*<0.01 vs control.


**Hoechst staining** Under the fluorescent microscope, ECV304 endothelial cell at 48 h after treatment with 15d-PGJ<sub>2</sub> showed the morphological changes apoptosis characteristic (Fig 2).

**TUNEL assay** To confirm that the effect of various concentrations of 15d-PGJ<sub>2</sub> (5-20 μmol/L) on apoptosis in ECV304 endothelial cells, TUNEL analysis was performed. The 15d-PGJ<sub>2</sub> induced apoptosis of ECV304 cells in a dose-dependent manner, and the percentage of apoptosis was enhanced from 10.0 %±1.3 % to 32.8 %±1.6 % (Fig 3).

**Cell cycle phase distribution** ECV304 endothe-



**Fig 2.** The fluorescent microphotograph: showing condensation and segmentation of nucleus in ECV304 endothelial cell at 48 h after treatment with 15d-PGJ<sub>2</sub>. A) Control; B) 15d-PGJ<sub>2</sub> 10 μmol/L. Hoechst stain, ×200.




**Fig 3.** Effect of 15d-PGJ<sub>2</sub> on apoptosis in ECV304 endothelial cells. (1): control; (2-4): 15d-PGJ<sub>2</sub> 5, 10, and 20 μmol/L. *n*=4. Mean±SD. <sup>c</sup>*P*<0.01 vs control.

lial cells were accumulated in G<sub>0</sub>/G<sub>1</sub> phase, after exposure to different concentrations of 15d-PGJ<sub>2</sub> (5-20 μmol/L) for 24 h. There was a significant decrease in the number of cells in S phase. No significant change of cell number was observed in the number of cells in G<sub>2</sub>/M phase (Tab 1).

**Tab 1.** Effect of 15d-PGJ<sub>2</sub> on ECV304 cells cycle. *n*=3. Mean±SD. <sup>b</sup>*P*<0.05 vs control.

15d-PGJ <sub>2</sub> /μmol/L	Cell number/%		
	G <sub>0</sub> /G <sub>1</sub>	S	G <sub>2</sub> /M
0	29±2	41±3	29±1
5	38±3 <sup>b</sup>	32±4 <sup>b</sup>	29±2
10	51±3 <sup>b</sup>	21±1 <sup>b</sup>	28±2
20	74±5 <sup>b</sup>	2±2 <sup>b</sup>	25±3

**Ladder detection assay** Agarose gel electrophoresis exhibited DNA ladder formation in ECV304 endothelial cells after exposure to different concentrations of 15d-PGJ<sub>2</sub> (5-20 μmol/L) for 48 h. Compared with the control group, the DNA laddering was clearly observed by the treatment with 15d-PGJ<sub>2</sub> (Fig 4).



**Fig 4.** DNA ladder pattern formation of ECV304 endothelial cells. Lane M: DNA marker; Lane 1: control; Lanes (2-4): 15d-PGJ<sub>2</sub> 5, 10, and 20 μmol/L.

**Effects of 15d-PGJ<sub>2</sub> on the expression of *c-myc*, *p21*, *p53*, and *Gadd45*** RT-PCR analysis showed that the expression of *c-myc* mRNA was down-regulated, whereas the expressions of *p53* mRNA and *Gadd45* mRNA were up-regulated (Fig 5). Meanwhile, Western blot showed the same changes in protein expressions (Fig 6). Both RT-PCR analysis and Western blot analysis showed no significant changes in the expression of *p21*.

**Effects of 15d-PGJ<sub>2</sub> on DNA binding activity of NF-κB and AP-1, and on expression of *c-jun*, *c-fos* and *IκBα*** EMSA was performed with nuclear extracts prepared from control or treated cells exposed to 15d-




Fig 5. Effect of 15d-PGJ<sub>2</sub> on the expression of *c-myc*, *p21*, *p53*, and *Gadd45* mRNA in ECV304 endothelial cells. Lane M : DNA marker; Lane 1: control; Lanes 2-4: 15d-PGJ<sub>2</sub> 5, 10, and 20  $\mu\text{mol/L}$ .




Fig 6. Effect of 15d-PGJ<sub>2</sub> on the expression of *c-myc*, *p21*, *p53*, and *Gadd45* protein in ECV304 endothelial cells. Lane M : DNA marker; Lane 1: control; Lanes 2-4: 15d-PGJ<sub>2</sub> 5, 10, and 20  $\mu\text{mol/L}$ .

PGJ<sub>2</sub> at various concentrations for 6 h. DNA binding activities of NF- $\kappa$ B and AP-1 were almost completely inhibited in 15d-PGJ<sub>2</sub>-treated cells compared with untreated cells (Fig 7). Western blotting showed that the degradation of I $\kappa$ B $\alpha$  and the expression of *c-jun* were inhibited by 15d-PGJ<sub>2</sub> in a dose-dependent manner (Fig 8).

#### Effect of 15d-PGJ<sub>2</sub> on activity of p38 kinase

The activity of p38 kinase in ECV304 endothelial cells was increased with the increase of 15d-PGJ<sub>2</sub> 5 -20  $\mu\text{mol/L}$  (Fig 9).

## DISCUSSION

Endothelial cell (EC) apoptosis is an initiating event in atherogenesis since EC apoptosis compromises vasoregulation and increases SMC proliferation,




Fig 7. Effect of 15d-PGJ<sub>2</sub> on DNA binding activity of AP-1 (A), NF- $\kappa$ B (B). (A): Lane 1: control; Lanes 2-4: 15d-PGJ<sub>2</sub> 5, 10, and 20  $\mu\text{mol/L}$ ; (B): Lane 1: supershift; lane 2: control; Lanes 3-5: 15d-PGJ<sub>2</sub> 5, 10, and 20  $\mu\text{mol/L}$ .




Fig 8. Effect of 15d-PGJ<sub>2</sub> on the expression of *c-jun*, *c-fos*, and I $\kappa$ B $\alpha$  protein in ECV304 endothelial cells. Lane 1: control; Lanes 2-4: 15d-PGJ<sub>2</sub> 5, 10, and 20  $\mu\text{mol/L}$ .




Fig 9. Effect of 15d-PGJ<sub>2</sub> on activity of p38 kinase in ECV304 endothelial cells.  $n=4$ . Mean $\pm$ SD. <sup>c</sup> $P<0.01$  vs control. (1): control; (2-4): 15d-PGJ<sub>2</sub> 5, 10, and 20  $\mu\text{mol/L}$ .

migration, and blood coagulation. In addition, EC overlying vascular lesions increase expression of pro-apoptotic proteins, such as Fas and Bax, while decrease levels of anti-apoptotic factors. Therefore, understanding EC apoptotic pathways altered in atherosclerosis may enable a greater understanding of disease pathogenesis and foster the development of new therapies<sup>[1]</sup>.

PPAR have an impact on the development of atherosclerosis. The expression of the PPAR was found in endothelial cells, macrophage foam cells, smooth muscle cells of human, and atherosclerotic lesions<sup>[2]</sup>. PPAR play important roles in fatty acid metabolism and storage in liver and adipose tissue, respectively. Furthermore, PPAR have a potentially important effect of inhibiting growth and migration of vascular cells<sup>[9]</sup>. PPAR not only control plasma levels of cholesterol and triglyceride concentrations, but also exert an activity at the level of the vascular wall, which might contribute to the atherosclerotic processes.

Our present study demonstrated that 15d-PGJ<sub>2</sub> inhibited the growth of ECV304 endothelial cells in a dose-dependent manner. This finding is in agreement with previous studies demonstrating the ability of 15d-PGJ<sub>2</sub> to inhibit proliferation in HUVEC and ECV304 cells<sup>[4]</sup>. The possible mechanism might include two aspects: 1) 15d-PGJ<sub>2</sub> had an impact on cell cycle phase. In our present study, flow cytometry analysis of ECV304 cells treated with 15d-PGJ<sub>2</sub> for 24 h revealed the accumulation of the cells at the G<sub>0</sub>/G<sub>1</sub> peak, and the reduction of cells in S phase. So we could infer that 15d-PGJ<sub>2</sub> inhibited the proliferation of ECV304 endothelial cells by arresting cell cycle; 2) 15d-PGJ<sub>2</sub> induced apoptosis in ECV304 endothelial cells. Hoechst 33258 staining showed that the morphological change characteristic for apoptosis could be observed in ECV304 cells at 48 h after treatment with 15d-PGJ<sub>2</sub>. Meanwhile, TUNEL and DNA ladder analysis both confirmed that 15d-PGJ<sub>2</sub> induced apoptosis of ECV304 endothelial cells in a dose-dependent manner.

The mechanism of 15d-PGJ<sub>2</sub> inhibition on cell proliferation and induction on apoptosis is still unclear. Some oxidation-sensitive transcription factors, such as NF- $\kappa$ B, the AP-1 complex, and *c-myc* are activated in atherosclerotic lesions and are involved in expression of early response genes that propagate atherogenic vascular changes<sup>[9-11]</sup>.

Signal-dependent activation of I $\kappa$ B kinase (IKK) results in phosphorylation and rapid degradation of I $\kappa$ B. Free NF- $\kappa$ B migrates into the nucleus and activates ex-

pression target gene<sup>[12]</sup>. AP-1 consists of a variety of dimers composed of member of the Fos, Jun, and ATF families of proteins. All kinds of nuclear receptor can affect the activity of AP-1 by inhibition of the JUN family member (c-jun) and the FOS family member (c-fos)<sup>[13]</sup>. NF- $\kappa$ B and AP-1 play important roles in cell apoptosis<sup>[5,14]</sup>. In our study, EMSA analysis showed that 15d-PGJ<sub>2</sub> strongly inhibited the DNA binding activity of NF- $\kappa$ B. Accordingly, Western blot analysis confirmed that 15d-PGJ<sub>2</sub> inhibited the degradation of I $\kappa$ B $\alpha$ . So we could infer that 15d-PGJ<sub>2</sub> inhibited the DNA binding activity of NF- $\kappa$ B by inhibiting the degradation of I $\kappa$ B $\alpha$ . In addition, EMSA analysis also showed that 15d-PGJ<sub>2</sub> distinctly inhibited the DNA binding activity of AP-1. Furthermore, Western blot analysis showed that 15d-PGJ<sub>2</sub> could decrease the expression of c-jun protein. It suggests that 15d-PGJ<sub>2</sub> inhibit the DNA binding activity of AP-1 by inhibiting the expression of c-jun protein. All results suggested that 15d-PGJ<sub>2</sub> induced apoptosis of ECV304 endothelial cells, which may be through inhibiting NF- $\kappa$ B and AP-1 activation pathways.

The proto-oncogene, *c-myc* plays a pivotal role in cell proliferation. *C-myc* has two coupled function: proliferation and apoptosis. These opposing roles of *c-myc* require that other products should interact with *c-myc* to determine the final outcome of cell<sup>[15]</sup>. *P53* is one of the most frequently studied tumor suppressing genes. *P53* seems to be responsible for the arrest of cell in the G<sub>1</sub> phase of the cell cycle. *P53* induces cell-cycle arrest or apoptosis in many types of cells and activates the transcription of target genes including *p21* and *Gadd45*<sup>[16]</sup>. *Gadd45* is a *p53*-regulated growth arrest gene. Evidence suggests that it plays an important role in regulation of cell growth. *Gadd45* may be involved in the G<sub>2</sub>/M checkpoint<sup>[17]</sup>. Both *Gadd45* and *p53* are the target genes of *c-myc*. *c-myc* has been shown to down-regulate both basal and stress-induced *Gadd45* expression<sup>[18]</sup>. Therefore, in order to understand the mechanism of 15d-PGJ<sub>2</sub> inducing apoptosis in ECV304 cells, we applied RT-PCR and Western blot to investigate the expression of *c-myc*, *p53*, *Gadd45*, and *p21*. Both RT-PCR and Western blot analysis showed that 15d-PGJ<sub>2</sub> could decrease the expression of *c-myc*, and induce the expression of *p53* and *Gadd45*. Whereas, 15d-PGJ<sub>2</sub> has no effect on the expression of *p21*. So we were able to infer that the down-regulation of *c-myc*, the up-regulation of *p53* and *Gadd45* play important roles in the apoptosis of ECV304 cells induced by 15d-PGJ<sub>2</sub>.

Recent studies have shown the possible involvement of MAPK (mitogen-activated protein kinase) pathway. p38 kinase belongs to the MAPK superfamily, which acts as a signaling molecule in cellular apoptosis. Takekawa and Saitohave<sup>[19]</sup> proposed the following model to the JNK/p38-mediated apoptotic pathways: *p53* is activated in response to genotoxic stress, then causes transcriptional up regulation of *Gadd45*, and *Gadd45* interacts with MTK1 (mitogen-activated protein kinase 1) to set into motion the JNK/p38-mediated apoptotic pathway. The proposed model implicates *p53* reside at upstream of JNK/p38 pathways and suggests transcriptional and translational dependence for JNK and p38 kinase activation. In our study, p38 kinase analysis showed that the activity of p38 kinase in ECV304 endothelial cells increased after the treatment with 15d-PGJ<sub>2</sub>. It was possible that 15d-PGJ<sub>2</sub> induced apoptosis in ECV304 cells by activating p38 kinase, which was activated by the up-regulation of *p53* and *Gadd45*.

In conclusion, 15d-PGJ<sub>2</sub> was able to inhibit ECV304 cell proliferation and induce apoptosis of ECV304. The effect was related to the inhibition of the DNA binding activities of NF- $\kappa$ B and AP-1; on the other hand, the effect was associated with the down-regulation of *c-myc*, up-regulation of *p53* and *Gadd45*, and activation of p38 kinase.

## REFERENCES

- 1 Choy JC, Granville DJ, Hunt DW, Mcmanus BM. Endothelial cell apoptosis: biochemical characteristics and potential implications for atherosclerosis. *J Mol Cell Cardiol* 2001; 33: 1637-90.
- 2 Ricote M, Huang J, Fajas L, Li A, Welch J, Najib J, *et al*. Expression of the peroxisome proliferation-activated receptor Proc Natl Acad Sci Proc Natly (PPAR gamma) in human atherosclerosis and regulation in macrophages by colony stimulating factors and oxidized low density lipoprotein. *Proc Natl Acad Sci USA* 1998; 95: 7614-9.
- 3 Toel B, Daid EM. The mechanisms of action of PPARs. *Annu Rev Med* 2002; 53: 409-35.
- 4 Bishop-Bailey D, Hla T. Endothelial cell apoptosis induced by the peroxisome proliferator-activated receptor (PPAR) ligand 15-deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub>. *J Biol Chem* 1999; 274: 17042-8.
- 5 Chinetti G, Griglio S, Antonucci M, Torra IP, Delerive P, Majd Z, *et al*. Activation of proliferator-activated receptors  $\alpha$  and  $\gamma$  induces apoptosis of human monocyte-derived macrophages. *J Biol Chem* 1998; 273: 25573-80.
- 6 Staels B, Koenig W, Habib A, Merval R, Lebrun M, Torra IP, *et al*. Activation of human aortic smooth-muscle cells is inhibited by PPAR  $\alpha$  but not by PPAR  $\alpha$  activators. *Nature* 1998; 393: 790-3.
- 7 Philippe D, Karolien DB, Sandrine B, Wim VB, Jeffrey MP, Trank JG, *et al*. Peroxisome proliferation-activated receptor alpha negatively regulates the vascular inflammatory gene response by negative cross-talk with transcription factors NF-kappaB and AP-1. *J Biol Chem* 1999; 274: 32048-54.
- 8 Cosulich S, James N, Roberts R. Role of MAP kinase signalling pathways in the mode of action of peroxisome proliferators. *Carcinogenesis* 2000; 21: 579-83.
- 9 Willa AH, Ronald EL. PPAR  $\gamma$  and Atherosclerosis effects on cell growth and movement. *Arterioscler Thromb Vasc Biol* 2001; 21: 1891-5.
- 10 Maulik N, Sasaki H, Addya S, Das DK. Regulation of cardiomyocyte apoptosis by redox-sensitive transcription factors. *FEBS Lett* 2000; 485: 2-12
- 11 Dang CV. *C-myc* target genes involved in cell growth, apoptosis, and metabolism. *Mol Cell Biol* 1999; 19: 1-11.
- 12 Baeuerle PA, Henkel T. Function and activation of NF- $\kappa$ B in the immune system. *Annu Rev Immunol*, 1994; 12: 141-79.
- 13 Caelles C, Gonzalez-Sancho JM., Munoz A. Nuclear hormone receptor antagonism with AP-1 by inhibition of the JNK pathway. *Genes Dev* 1997; 11: 3351-64.
- 14 Wang C, Mayo MW, Baldwin AS. TNF and cancer therapy-induced apoptosis: potentiation by inhibition of NF- $\kappa$ B. *Science* 1996; 274: 784-9.
- 15 Hoffman B, Lieberman DA. The proto-oncogene *c-myc* and apoptosis. *Oncogene* 1998; 17: 3351-7.
- 16 Wang XW, Zhan Q, Coursen JD, Khan MA, Kontrry HU, Yu L, *et al*. GADD45 induction of a G<sub>2</sub>/M cell cycle checkpoint. *Proc Natl Acad Sci* 1999; 96: 3706-11.
- 17 Li HL, Ren XD, Zhang HW, Ye CL, Lu JH, Zheng PE, *et al*. Synergism between heparin and adriamycin on cell proliferation and apoptosis in human nasopharyngeal carcinoma CNE2 cells. *Acta Pharmacol Sin* 2002; 23: 167-72.
- 18 Aundson SA, Zhan Q, Penn LZ, Fownace AJ. Myc suppresses induction of the growth arrest *gadd34*, *gadd45* and *gadd153* by DNA-damaging agents. *Oncogene* 1998; 17: 2149-54.
- 19 Takekawa M, Saito H. A family of stress-inducible GADD45-like proteins mediate activation of the stress-responsive MTK1/MEKK4 MAPKKK. *Cell* 1998; 95: 521-30.