

Inhibitory effect of dopamine on Ca^{2+} -calmodulin-dependent protein kinase II activity in rat hippocampal slices¹

HOU Xiao-Yu, ZHANG Guang-Yi² (Research Center of Biochemistry and Molecular Biology, Xuzhou Medical College, Xuzhou, Jiangsu 221002, China)

KEY WORDS dopamine; Ca^{2+} -calmodulin dependent protein kinase; cerebral ischemia; calcium; dopamine agonists; dopamine antagonists; hippocampus

ABSTRACT

AIM: To study the effect of dopamine (DA) on Ca^{2+} -calmodulin dependent protein kinase II (CCDPK II) activity in rat hippocampus. **METHODS:** Using rat hippocampal slices as an *in vitro* model, the activity of CCDPK II was examined by the method of ³²P-incorporation. **RESULTS:** Exogenous DA reduced CCDPK II activity in hippocampal slices in a concentration- and time-dependent manner. Removal of extracellular calcium antagonized the DA-induced inhibition of CCDPK II activity, partially or completely. The activity of CCDPK II was markedly decreased by apomorphine (a nonselective DA receptor agonist), SKF38393 (a selective D₁-like DA receptor agonist), or quinpirole (a selective D₂-like DA receptor agonist). The inhibition of CCDPK II activity induced by exogenous DA was abolished by preincubation with Sch-23390, a selective D₁-like DA receptor antagonist, or domperidone, a selective D₂-like DA receptor antagonist. **CONCLUSION:** DA has an inhibitory effect on CCDPK II activity in rat hippocampus, related to stimulation of D₁-like and D₂-like receptors and calcium influx.

INTRODUCTION

The hippocampus was the most vulnerable brain

region to ischemic injury. The excessive release of DA in the hippocampus during cerebral ischemia played an important role in the development of ischemic cerebral injury^(1,2). However, the mechanism underlying dopaminergic neurotoxicity remained unclear.

Overload of intracellular calcium was thought to be a pivotal factor in ischemic cerebral injury. Elevation of intracellular calcium concentration led to activation of calcium-dependent lipases, nucleases, proteases, nitric oxide synthases, and protein kinases such as CCDPK II, etc., and consequent neuronal cell death. CCDPK II, which was highly enriched in neural tissues and composed up to 2% of total proteins in the hippocampus, was a multifunctional protein kinase and regulated many physiological responses to calcium mobilization including gene expression, neurotransmitter synthesis and exocytosis, postsynaptic responses and cytoskeletal interactions, etc. Previously, we reported that hypoxia could result in the significant inhibition of CCDPK II activity in rat hippocampus⁽³⁾. The inhibition of the enzyme activity had been implicated in contributing to neuronal cell damage⁽⁴⁾. Therefore, elucidating the effect of DA on CCDPK II activity would provide insight into the molecular mechanism of dopaminergic neurotoxicity in cerebral ischemic injury. In this paper, the effect of DA on CCDPK II activity in rat hippocampus was firstly investigated.

MATERIALS AND METHODS

Materials Dopamine hydrochloride, *d,l*-SKF38393 hydrochloride, *l*-quinpirole hydrochloride, R(+)-Sch-23390 hydrochloride, and domperidone (RBI, USA); apomorphine (Sigma, USA). They were dissolved in a small amount of distilled water, except quinpirole was dissolved in H₂SO₄ (0.1 mol · L⁻¹) and domperidone in 1% lactate. They were

¹ Project supported by the National Natural Science Foundation of China, No 39470168.

² Correspondence to Prof ZHANG Guang-Yi.
Ptm 86-516-574-8423. Fax 86-516-574-8429.
E-mail xmcb@public.xz.js.cn

Received 1998-11-27. Accepted 1999-05-18

diluted with glucose-free Krebs-Ringer solution immediately before use.

[γ - 32 P]ATP was purchased from Yahui Biological and Medical Engineering Co, Beijing; sodium metabisulfite and ATP from Sigma; phenylmethane-sulfonylfluorid (PMSF), and mercapto ethenol (β -ME) from E Merck. Other reagents were of AR.

Hippocampal slice preparation and drug application Male Sprague-Dawley rats (Grade II, Certificate No 02-49-02, purchased from Sippr-BK Experimental Animal Ltd Co, Shanghai) weighing 160 - 200 g were decapitated and the brains were placed in ice-cold Krebs-Ringer solution: NaCl 122, KCl 3.1, KH_2PO_4 0.4, MgSO_4 1.2, CaCl_2 1.3, NaHCO_3 25, glucose 10 $\text{mmol}\cdot\text{L}^{-1}$; pH 7.4 equilibrated with 95 % O_2 + 5 % CO_2 . Parasagittal hippocampal slices (350 μm) were prepared with a McIlwain tissue chopper and preincubated in normal Krebs-Ringer solution at 36 $^\circ\text{C}$ for 90 min. Following preincubation, slices were transferred to glucose-free Krebs-Ringer solution and incubated with or without drugs for 30 min or as indicated. During whole course of preincubation and drug application, incubation medium was continuously bubbled with 95 % O_2 + 5 % CO_2 . Sodium metabisulfite (200 $\mu\text{mol}\cdot\text{L}^{-1}$) was added to the dopamine-containing solution to prevent oxidation of dopamine. In pilot experiments, sodium metabisulfite (200 $\mu\text{mol}\cdot\text{L}^{-1}$) did not affect the activity of CCDPK II. DA receptor antagonists were added 15 min before exposure to DA. After experiments, the slices were quickly frozen in liquid nitrogen until use.

Assay for CCDPK II activity The slices were homogenized in a glass homogenizer on ice with ice-cold homogenization buffer: Tris-HCl 20, edetic acid 2, NaF 2, PMSF 0.5, β -ME 10 $\text{mmol}\cdot\text{L}^{-1}$; pH 7.5. The homogenate was spun at 10 000 $\times g$ at 4 $^\circ\text{C}$ for 5 min, and the supernatant was assayed for Ca^{2+} -calmodulin-dependent activity of CCDPK II by the method of ^{32}P -incorporation⁽³⁾. The radioactivity was measured by a liquid scintillation spectrometer (LS 6500, Beckman). Protein concentration of the supernatant was determined with bovine serum albumin (BSA) as standard⁽⁵⁾. The CCDPK II activity was expressed in the amount of ^{32}P -incorporation ($\text{nmol}\cdot\text{min}^{-1}\cdot\text{g}^{-1}$).

Statistical analysis Values were expressed as $\bar{x} \pm s$ and compared by ANOVA followed by the

Duncan's new multiple range method or Newman-Keuls test.

RESULTS

Inhibitory effect of DA on CCDPK II activity After exposure of rat hippocampal slices to exogenous DA at 100 $\mu\text{mol}\cdot\text{L}^{-1}$, the activity of CCDPK II decreased gradually with incubation time. Much inhibition of CCDPK II activity was observed 20 min after the addition of DA (Fig 1).




Fig 1. Inhibitory effect of DA (100 $\mu\text{mol}\cdot\text{L}^{-1}$) on CCDPK II activity in rat hippocampal slices. $n = 4$ rats. $\bar{x} \pm s$. ^c $P < 0.01$ vs control.

Incubation with DA at 30 - 2000 $\mu\text{mol}\cdot\text{L}^{-1}$ for 30 min markedly depressed CCDPK II activity. The maximal inhibition was observed with DA at 500 $\mu\text{mol}\cdot\text{L}^{-1}$, with the enzyme activity falling to 54.4 % of the control. Incubation of slices in calcium-free medium (with 1 $\text{mmol}\cdot\text{L}^{-1}$ egtazic acid) containing DA at the same concentrations antagonized the DA-induced inhibition of CCDPK II activity, partially or completely (Tab 1).

Effect of DA receptor agonists on CCDPK II activity Incubation with apomorphine (1 - 20 $\mu\text{mol}\cdot\text{L}^{-1}$), a nonselective DA receptor agonist, for 30 min resulted in an inhibition of CCDPK II activity. The maximal inhibition was observed with apomorphine at 10 $\mu\text{mol}\cdot\text{L}^{-1}$, with the enzyme activity falling to 53.5 % of the control. Incubation for 30 min with SKF38393, a selective D_1 -like DA receptor agonist, or quinpirole, a selective D_2 -like DA receptor agonist, reduced the activity of CCDPK II. SKF38393 at 1 - 200 $\mu\text{mol}\cdot\text{L}^{-1}$ and quinpirole at 20 - 200 $\mu\text{mol}\cdot\text{L}^{-1}$

Tab 1. Inhibitory effect of exogenous DA on CCDPK II activity in rat hippocampal slices with or without extracellular calcium ($[Ca^{2+}]_o$). $n=6$ rats. $\bar{x} \pm s$. $^aP > 0.05$, $^cP < 0.01$ vs control. $^fP < 0.01$ vs the respective values obtained with $[Ca^{2+}]_o$.

DA/ $\mu\text{mol} \cdot \text{L}^{-1}$	CCDPK II activity/ $\text{nmol} \cdot \text{min}^{-1} \cdot \text{g}^{-1}$ + $[Ca^{2+}]_o$	$\text{nmol} \cdot \text{min}^{-1} \cdot \text{g}^{-1}$ - $[Ca^{2+}]_o$
0	169 ± 12	169 ± 11
30	138 ± 11 ^c	179 ± 22 ^{af}
100	116 ± 5 ^c	148 ± 7 ^{cf}
500	92 ± 8 ^c	150 ± 6 ^{cf}
1000	132 ± 9 ^c	161 ± 4 ^{af}
2000	124 ± 8 ^c	171 ± 16 ^{af}

resulted in an inhibition of the enzyme activity (Tab 2).

Tab 2. Effect of apomorphine, SKF38393, and quinpirole on CCDPK II activity in rat hippocampal slices. $n=6$ rats. $\bar{x} \pm s$. $^cP < 0.01$ vs control.

Treatment	Concentration/ $\mu\text{mol} \cdot \text{L}^{-1}$	CCDPK II activity/ $\text{nmol} \cdot \text{min}^{-1} \cdot \text{g}^{-1}$
Control	-	170 ± 11
Apomorphine	1	130 ± 11 ^c
	5	110 ± 11 ^c
	10	91 ± 6 ^c
	20	123 ± 7 ^c
SKF38393	1	140 ± 12 ^c
	10	136 ± 10 ^c
	100	123 ± 8 ^c
	200	103 ± 10 ^c
Quinpirole	10	171 ± 17
	20	147 ± 8 ^c
	50	107 ± 9 ^c
	100	111 ± 9 ^c
	200	134 ± 11 ^c

Effect of DA receptor antagonists on DA-induced inhibition of CCDPK II activity

Preincubation of slices with specific DA receptor antagonists, Sch-23390 (D_1 -like DA receptor antagonist) or domperidone (D_2 -like DA receptor antagonist), antagonized the inhibition of CCDPK II activity induced by exogenous DA ($500 \mu\text{mol} \cdot \text{L}^{-1}$). Sch-23390 at $50 \mu\text{mol} \cdot \text{L}^{-1}$ and domperidone at $10 \mu\text{mol} \cdot \text{L}^{-1}$ abolished the DA-induced inhibition of CCDPK II activity (Tab 3).

Tab 3. Protection of Sch-23390 and domperidone against inhibition of CCDPK II activity induced by exogenous DA in rat hippocampal slices. $n=6$ rats. $\bar{x} \pm s$. $^cP < 0.01$ vs DA ($500 \mu\text{mol} \cdot \text{L}^{-1}$). $^dP > 0.05$ vs control.

Treatment	Concentration/ $\mu\text{mol} \cdot \text{L}^{-1}$	CCDPK II activity/ $\text{nmol} \cdot \text{min}^{-1} \cdot \text{g}^{-1}$
Control	-	169 ± 12
DA	500	92 ± 8
DA + Sch-23390	1	106 ± 15
	5	121 ± 13 ^c
	20	136 ± 6 ^c
	50	163 ± 16 ^{cd}
DA + domperidone	0.1	114 ± 8 ^c
	1	132 ± 11 ^c
	5	152 ± 5 ^c
	10	169 ± 14 ^{cd}

DISCUSSION

For the first time, the present study shows that DA has an inhibitory effect on CCDPK II activity in the hippocampus, depending on both duration of DA exposure and DA concentration. This DA-mediated effects were blocked by removal of extracellular calcium. The result suggests that the inhibitory effect of DA on CCDPK II activity is partially related to influx of extracellular calcium. Maybe there is release of intracellular calcium storage because the inhibitory effects of DA at some concentrations were not antagonized completely.

It has been demonstrated that the inhibition of CCDPK II activity is concerned with calcium-stimulated autophosphorylation of the enzyme at a threonine residue, Thr²⁶⁶. The autophosphorylation converts the kinase from the Ca^{2+} -calmodulin-dependent form to Ca^{2+} -calmodulin-independent form, which results in the inhibition of Ca^{2+} -calmodulin-dependent activity and the enhancement of Ca^{2+} -calmodulin-independent activity. Excessive activated CCDPK II can potentiate the presynaptic release of glutamate by phosphorylating synapsin I^[6], but also phosphorylate postsynaptic *N*-methyl-*D*-aspartate (NMDA) receptors or non-NMDA receptors and promote their subsequent responsiveness to released glutamate^[7,8]. Excessive stimulation of glutamate receptors causes massive calcium influx and results in

neuronal cell damage.

All above results may support our hypothesis that neurotoxic effects of DA would partially due to DA-induced the inhibition of CCDPK II activity, which enhances excitotoxicity of excitory amino acid and increases calcium influx.

Radioligand assay proves the distribution of DA receptors in the hippocampus. Direct dopamine receptor stimulation may mediate inhibitory effect of DA on CCDPK II activity. Apomorphine, a non-selective DA receptor agonist had an inhibitory effect on CCDPK II activity, similar to the effect of DA. The result suggests the effect of DA is mediated by stimulating DA receptors. The DA receptors are now classified as "D₁-like" (D₁ and D₅) and "D₂-like" (D₂, D₃, and D₄). In order to assess the receptor subtypes mediating DA inhibitory effect, selective DA receptor agonists and antagonists were used. The results suggest stimulation of D₁-like or D₂-like receptor subtypes can both result in inhibition of the enzyme activity. The inhibition of CCDPK II activity is calcium-dependent. Therefore, our results demonstrate that stimulation of D₁-like or D₂-like receptor subtypes can elevate the intracellular calcium concentration in rat hippocampus. Intracellular calcium accumulation causes autophosphorylation of the enzyme, which makes the kinase convert to Ca²⁺-calmodulin-independent form and results in inhibition of Ca²⁺-calmodulin-dependent activity. However, The mechanism of D₁-like or D₂-like receptor mediating intracellular calcium accumulation remains to be elucidate.

At higher concentration, DA, apomorphine, quinpirole-induced inhibitory rates of CCDPK II activity decreased. This may result from activation of Ca²⁺-calmodulin-dependent protein phosphatases, such as calcineurin. Calcineurin is abundant in the hippocampus. Several proteins including synapsin I and NMDA receptors are common to CCDPK II and calcineurin as substrates for phosphorylation and dephosphorylation^[6-10]. It had been reported that stimulation of D₂-like receptors can activate calcineurin^[11]. The activation of calcineurin leads to dephosphorylation of synapsin I and NMDA receptors^[9,10] and consequent decrease of calcium influx induced by excessive stimulation of glutamate receptors. As a result of D₂-like receptors stimulation

at higher concentration, the DA-induced inhibitory rates of CCDPK II activity is attenuated. The above results indicated that the inhibition of CCDPK II activity induced by exogenous DA might be mainly due to stimulation of D₁-like receptors.

In conclusion, DA has an inhibitory effect on CCDPK II activity in rat hippocampus, related to stimulation of D₁-like or D₂-like receptor subtypes and calcium influx.

ACKNOWLEDGMENTS The authors wish to thank Prof JIN Guo-Zhang (Shanghai Institute of Materia Medica, Chinese Academy of Sciences) for his helpful advice and suggestion.

REFERENCES

- 1 Yamamoto Y, Tanaka T, Shibata S, Watanabe S. Involvement of D₁ dopamine receptor mechanism in ischemia-induced impairment of CA1 presynaptic fiber spikes in rat hippocampal slices. *Brain Res* 1994; 665: 151-4.
- 2 Maginn M, Kelly JP, Leonard BE. Protective effects of vanoxamine (GBR12909) against ischaemia-induced hyperactivity and neurodegeneration in the gerbil model of cerebral ischaemia. *Pharmacol Biochem Behav* 1997; 56: 727-35.
- 3 Pei L, Zong YY, Zhang L, Sun YF, Zhang GY. Studies on mechanism of the inhibition of Ca²⁺/CaMPK II activity induced by hypoxia in rat hippocampal slices. *Acta Physiol Sin* 1997; 49: 61-6.
- 4 Hajimohammadreza I, Probert AW, Coughenour LL, Borosky SA, Marcoux FW, Boxer PA, *et al.* A specific inhibitor of calcium/calmodulin-dependent protein kinase-II provides neuroprotection against NMDA- and hypoxia/hypoglycemia-induced cell death. *J Neurosci* 1995; 15: 4093-101.
- 5 Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. *J Biol Chem* 1951; 193: 265-75.
- 6 Stefani G, Onofri F, Valtorta F, Vaccaro P, Greengard P, Benfenati F. Kinetic analysis of the phosphorylation-dependent interactions of synapsin I with rat brain synaptic vesicles. *J Physiol (Lond)* 1997; 504: 501-15.
- 7 Omlkumar RV, Kiely MJ, Rosenstein AJ, Min KT, Kennedy MB. Identification of a phosphorylation site for calcium/calmodulin-dependent protein kinase II in the NR2B subunit of the N-methyl-D-aspartate receptor. *J Biol Chem* 1996; 271: 31670-8.
- 8 Barria A, Muller D, Derkach V, Griffith LC, Soderling TR. Regulatory phosphorylation of AMPA-type glutamate receptors by CaM-K II during long-term potentiation. *Science* 1997; 276: 2042-5.

- 9 Victor RG, Thomas GD, Marban E, O'Rourke B. Presynaptic modulation of cortical synaptic activity by calcineurin. Proc Natl Acad Sci USA 1995; 92: 6269 - 73.
- 10 Tong G, Shepherd D, Jahr CE. Synaptic desensitization of NMDA receptors by calcineurin. Science 1995; 267: 1510 - 2.
- 11 Nishi A, Snyder GL, Greengard P. Bidirectional regulation of DARPP-32 phosphorylation by dopamine. J Neurosci 1997; 17: 8147 - 55.

102-906

8

目的: 研究多巴胺(DA)对大鼠海马脑片 Ca^{2+} -钙调素依赖性蛋白激酶 II (CCDPK II)活性的影响。
方法: 采用大鼠海马脑片体外培养模型,以 ^{32}P 掺入法测定CCDPK II的活性。
结果: 外源性 DA 可显著降低大鼠海马脑片CCDPK II活性,并有一定的浓度依赖性和时间依赖性。去除胞外的 Ca^{2+} 对不同浓度 DA 诱导的CCDPK II 活性抑制有部分或完全保护作用。阿扑吗啡(非特异性 DA 受体激动剂)、SKF38393 (特异性 D_1 样 DA 受体激动剂)和哇吡罗(特异性 D_2 样 DA 受体激动剂)均可显著降低CCDPK II的活性。Sch-23390 (特异性 D_1 样 DA 受体拮抗剂)和多潘立酮(特异性 D_2 样 DA 受体拮抗剂)均可拮抗 DA 所诱导的酶活性抑制。
结论: DA 抑制海马CCDPK II 的活性,其作用机制与 D_1 样和 D_2 样受体以及胞外 Ca^{2+} 的内流有关。

多巴胺诱导大鼠海马脑片 Ca^{2+} -钙调素依赖性蛋白激酶 II 活性的抑制作用¹

R972.1

侯筱宇, 张光毅² (徐州医学院生物化学与分子生物学研究中心, 江苏 221002, 中国)

关键词 多巴胺; Ca^{2+} -钙调素依赖性蛋白激酶; 脑缺血; 钙; 多巴胺激动剂; 多巴胺拮抗剂; 海马

(责任编辑 李颖)

CCDPK II 药理

读者、订阅者注意

本刊《中国药理学报》(英文刊名 Acta Pharmacologica Sinica)数年来坚持走国际化道路,发表国内外全英文来稿,关持续进入国际著名的 SCI 核心榜中及其他国际检索系统如 BA、CA、IM 等。报道内容具有国际先进水平及我国药理学研究特色。受到国内外同行好评。现经上级有关部门批准,2000 年起,将登记为英文刊。刊号(ISSN 0253-9756, CN 31-1347/R)和邮发代号(4-295)不变,内容和现有格式基本不变。但邮发目录中将标注(英文)字样。请读者注意通知有关订购部门继续订购《中国药理学报》(英文)。本刊并无另外中文版本。

国内年定价人民币 300 元, 每期定价 25 元, 欢迎订阅。

联系地址: 200031 上海市太原路 294 号《中国药理学报》编辑部

电话: 021-6474-2629, 021-6431-1833 转 200, 202 分机

传真: 021-6474-2629, 021-6437-0269

电子邮件 (E-mail): aps@server.shnc.ac.cn hxiao@mail.shnc.ac.cn