

S1.1**Anti-cancer activity of *Glycosmis parva* leaf extract on human colorectal cancer HT29 cell**

Nattaporn BURANABUNWONG¹, Nijisiri RUANGRUNGSI², Wacharee LIMPANASITHIKUL³.
¹Interdisciplinary Program of Pharmacology, Graduate School, Chulalongkorn University, Bangkok, Thailand; ²College of Public Health Sciences, Chulalongkorn University, Bangkok, Thailand; ³Department of Pharmacology, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

Aim: *Glycosmis parva* leaf extract was reported to have cytotoxicity and anti-inflammatory activities with the reduction of COX2 expression *in vitro*. This study aimed to investigate the effect of ethyl acetate extracts from leaves of *G. parva* (GPE) on human colorectal cancer expressing COX2-HT29 cells and its underlying mechanisms of action. **Methods:** Cytotoxicity of 6.25–100 µg/mL GPE against HT29 was done by resazurin reduction assay. Induction of apoptosis was determined by AnnexinV/FITC and PI staining. Anti-proliferative study was performed using automated cell counter. Cell cycle pattern was analyzed by PI staining. The expression of Bcl-2 family proteins was examined by qPCR. COX2 expression was determined by RT-PCR. **Results:** GPE was cytotoxic to HT29 in a dose-dependent manner with IC₅₀ values 69.50, 55.86, and 49.04 µg/mL at 24, 48, and 72 h, respectively. GPE at 50–100 µg/mL significantly induced HT29 apoptosis. At 25–100 µg/mL, it dose-dependently inhibited HT29 cell proliferation. GPE at 100 µg/mL caused G2M accumulation. It changed the expression of Bcl-2 family proteins. It also significantly decreased COX2 expression at 25–100 µg/mL. **Conclusion:** GPE exhibited cytotoxic activity, induction of apoptosis, inhibition of cell proliferation and arrest cell cycle in HT29 cells. The underlying mechanisms may involve decrease in COX2 expression and changes in the expression of Bcl-2 family proteins and which favor cancer cell death.

Keywords: *Glycosmis parva*; Colorectal cancer; COX2; Bcl-2; HT29

Acknowledgements: Chulalongkorn University Graduate Scholarship to commemorate the 72nd anniversary of His Majesty King Bhumibol Adulyadej and The 90th Anniversary of Chulalongkorn University Fund.

S1.2**Role of tissue factor (TF)/protease-activated receptor-2 (PAR-2) signaling in pancreatic cancer progression**

Li-Hsun CHANG¹, Shioh-Lin PAN², Che-Ming TENG¹. ¹Pharmacological Institute, College of Medicine, National Taiwan University, Taipei, Taiwan, China; ²The PhD program for Cancer Biology and Drug Discovery, College of Medical Science and Technology, Taipei Medical University, Taipei, Taiwan, China

Aim: Patients with advanced pancreatic cancer are associated with venous thromboembolism (VTE). Tissue factor (TF) is a biomarker in initiation of thrombosis and cancer progression. The aim of this study was to investigate the pathological role of TF/protease-activated receptor-2 (PAR-2) signaling in pancreatic cancer. **Methods:** First, effects of PAR-2 activation on gene expression investigated by microarray analysis. Then, PAR-2-mediated genes were confirmed by RT-PCR and Western blot analysis. VEGF-A levels were detected by an ELISA kit. For *in vitro* angiogenesis assay, HUVECs proliferation was measured by crystal violet staining; tube formation was examined by Matrigel tube formation assay. **Results:** According to the results of microarray assay, TF/PAR-2 signaling regulated an oncogenic pathway through increasing both HIF-1α and HIF-2α, resulting in enhanced transcription of TGF-α. Downregulation of HIF-α by siRNA or YC-1 resulted in depleted levels of TGF-α protein. Furthermore, the culture medium from PAR-2-treated pancreatic cancer cells enhanced HUVEC proliferation and tube formation, which was blocked by the MEK inhibitor PD98059. We also found that activated PAR-2 enhanced tumor angiogenesis through the release of VEGF from cancer cells, independent of the HIF pathways. **Conclusion:** The TF/PAR-2 signaling induced human pancreatic cancer progression through the induction of TGF-α expression by HIF-α, as well as through MEK/VEGF-mediated angiogenesis, and it plays a role in the interaction between cancer progression and cancer-related thrombosis.

Keywords: protease-activated receptor-2 (PAR-2); tissue factor (TF); HIF; VEGF; pancreatic cancer

S1.3**Cisplatin downregulates the aryl hydrocarbon receptor nuclear translocator to inhibit MDR1 expression**

Wen-Chang CHANG¹, Ben-Kuen CHEN². ¹Graduate Institute of Medical Sciences, College of Medicine, Taipei Medical University, Taipei 110, Taiwan, China; ²Institute of Bioinformatics and Biosignal Transduction, College of Bioscience and Biotechnology, National Cheng Kung University, Tainan 701, Taiwan, China

Aim: Identification of molecular pathways essential for cancer cells is critical for understanding their underlying biology and designing effective cancer therapies. In this study, effect of expression of the aryl hydrocarbon receptor nuclear translocator (ARNT) on the development of acquired drug resistance in cancer cells was investigated. **Methods:** cisplatin-sensitive or acquired resistant cancer cells were treated with cisplatin for a period of time in ARNT knockdown or overexpression conditions. The cell survival and apoptotic markers such as expression of p53, caspase-3 activation and DNA fragmentation were then detected by CCK-8 assay, MTT assay, flow cytometric and Western blot analysis. The effect of ARNT on MDR1 expression was determined by luciferase reporter analysis, RT-PCR and Western blot analysis. In addition, the cooperation between Sp1 and ARNT in the regulation of MDR1 expression was examined by DAPA and luciferase reporter analysis. In a xenograft analysis of SCID mice, the effect of cisplatin on ARNT-deficient c4 tumors and ARNT-containing vT2 tumor formation was studied. **Results:** Cisplatin dramatically induced p53 expression and caspase-3 activation, resulting in degradation of the ARNT and cell apoptosis in drug-sensitive cells. ARNT interacted with Sp1 to enhance the expression of MDR1 through Sp1-binding sites in the MDR1 promoter, resulting in a reversal of the effect of cisplatin on cell death. In contrast, suppression of ARNT reversed the characteristics of cisplatin-resistant cells, making them cisplatin-sensitive, and significantly enhancing cell death with cisplatin treatment. In addition, cisplatin also efficiently inhibited ARNT-deficient c4 tumors but not ARNT-containing vT2 tumor formation. **Conclusion:** These results reveal previously unrecognized multifaceted functions of ARNT in establishing drug-resistant properties of cancer cells, highlighting its potential as a therapeutic target for an important subset of cancers.

Keywords: cisplatin; ARNT; MDR1; drug resistance

S1.4**The effect of high-efficiency immune modulators on inhibiting tumor growth and regulating immunity in S₁₈₀ sarcoma mice**

Xin-feng CHANG^{1,2}, Chun-hua SONG², Hong-yan LI², Yu-fu LIN², Yan GONG², Zhi-hua HUANG², Li-min HAN^{1,2,*}. ¹Hunan Chinese Traditional Medicine University Changsha, 410007, China; ²Gannan Medical University, Ganzhou 341000, China

*To whom correspondence should be addressed.

E-mail: jxhlm@163.com

Aim: To observe the inhibition effect of high-efficiency immune modulators (HIS) on tumor growth in S₁₈₀-bearing mice and its regulation in immunologic function and improvement of tumour apoptosis. **Methods:** The second day of S₁₈₀-bearing mice model has established, mice were randomly divided into control group, model group, cyclophosphamide (CTX) group (50 mg/kg) and HIS group. After 10 days of different treatments, the mice's spleen index and inhibition rates of tumor were measured; the level of IL-6 was measured by Enzyme-linked immunosorbent assay (ELISA); The expression of IL-10 and Bcl-2 and Bax were detected by Western blot. **Results:** HIS can significantly inhibit the tumor growth with a 46.67% tumor inhibition rate. Compared with the model group, HIS group mice showed a decrease Bcl-2 (0.38±0.03, P<0.01) and an increase Bax (0.99±0.06, P<0.01) expression in S₁₈₀ sarcoma. HIS can also improve immunologic function by increasing the spleen index in S₁₈₀ sarcoma mice (P<0.01). The level of IL-6 in serum and the expression of IL-10 in S₁₈₀ sarcoma tissue were down-regulated in HIS treated mice. **Conclusion:** HIS had an anti-tumor effect on S₁₈₀-bearing mice, and its mechanisms probably result of inducing the apoptosis of S₁₈₀ sarcoma cells by the means of decreasing Bcl-2 and increasing Bax expression and improving the immunologic function via increasing the spleen index and downregulating the expression of IL-6 and IL-10.

Keywords: high-efficiency immune modulators; Bcl-2; Bax; IL-6; IL-10; tumor growth
Acknowledgements: This work was supported by grants from the Jiangxi Natural Science Foundation (2009GGY0014) and Bureau of Education (GJJ12557) of Jiangxi province.

S1.5**The role of microRNAs in tumor**

Shu-lan CHEN^{1,2}, Jun ZHONG², Li-juan LIU². ¹Medical School of Nanchang University, Nanchang 330006, China; ²Jiangxi Cancer Hospital, Nanchang 330006, China

*To whom correspondence should be addressed.

E-mail: liu_lujuanpharm@163.com

miRNA can function as either tumour suppressors or oncogenes in cancers by targeting genes associated with tumor, such as anti-apoptosis genes, cell proliferation genes, senescence genes and cell cycle factors, etc. These suggest miRNAs can be used as predictive, diagnosed and prognostic biomarker.

Moreover, based on the increasing number of studies demonstrate that the aberrant expression of miRNAs can predict the response to chemotherapeutic drugs, it is of great clinical value. miRNAs induced resistance to 5-fluorouracil in colorectal cancer by targeting some associated genes, including DNA, mismatch repair system, cell cycle factors and apoptosis factors, etc.

Keywords: microRNA; oncogene; tumour suppressors; drug resistance

Acknowledgements: This work was supported by Special Major Research Projects of Jiangxi Province (No 20114013)

S1.6

4-Acetylanthroquinol B isolated from *Antrodia cinnamomea* inhibited prostate cancer cells growth and metastasis *in vitro* and *in vivo*.

Ching-Hu CHUNG. Department of pharmacology, School of Medicine, Tzu Chi University, Hualien, Taiwan, China

Aim: The 4-acetylanthroquinol B (4AAQB) isolated from the mycelium of *Antrodia camphorata* could inhibit proliferation of hepatocellular carcinoma cells HepG2 with very low IC₅₀. However, the detail molecular mechanism underlying this anti-tumor drug is poorly understood. We are interesting to investigate the *in vitro* and *in vivo* anti-proliferation and anti-metastasis effects of 4AAQB in prostate cancer. **Methods:** We first used MTT assay to determine the effect of 4AAQB on PC3 cell viability. Further, we used LDH release assay to verify the inhibition effect either causing cytotoxicity or approving apoptosis. Then, cell scratch motility assay was used to evaluate whether the 4AAQB would inhibit motility of PC3 cell line. To investigate the possible mechanisms involved in the inhibitory effect of 4AAQB, the phosphorylation of Akt, ERK and IκB was analysis by Western blot. 4AAQB IP administration was used to examine the anti-tumor and anti-metastasis effect *in vivo* model. **Results:** 4AAQB does-dependently inhibited the PC3 cell viability and there were no cytotoxicity at doses less than 3 μmol/L. In non-toxic concentration, 4AAQB does-dependently inhibited the migration of PC3 cells *in vitro*. Furthermore, 4AAQB dose and time-dependently affected the Akt, ERK and IκB phosphorylation in PC3 cells. 4AAQB also significantly inhibited tumor growth and metastasis in PC3 *in vivo* model. **Conclusion:** Taken together, these findings suggested that 4AAQB could reduce the growth and metastasis of PC3 cells *in vitro* and *in vivo*. 4AAQB may be a promising therapeutic agent for prostate cancer. **Keywords:** 4AAQB; prostate cancer; proliferation; metastasis

S1.7

Anticancer effects of Gekko ethanol extract in human hepatocellular carcinoma HepG2 cells and in H22 bearing mice

Chao-chu CUI, Jian-gang WANG, Leng-xin DUAN, Xin-li XU. The Key Laboratory of Pharmacology and Medical Molecular Biology, Medical College, Henan University of Science and Technology, Luoyang 471003, China

Aim: To explore the anticancer effects and underlying mechanism of Gekko ethanol extract (GEE) *in vitro* and *in vivo*. **Methods:** MTT assay was employed to evaluate the anti-proliferative effect of GEE *in vitro*. Hoechst 33258 fluorescence staining was performed to observe morphologic changes nucleus of HepG2 cells; Antitumor properties of GEE *in vivo* were tested in H22 tumor bearing mice. Tumor weight of each mouse was evaluated at the end. The expression levels of IL-6 and TNF-α were analyzed by ELISA assay. Western blotting analysis was carried out to detect the expression levels of caspase-3 and VEGF in tumors. **Results:** GEE has anti-proliferative effect in a time-dose dependent manner, with inducing typical apoptotic morphological changes ability in HepG2 cells. GEE resulted in significant decrease in tumor weight. The expression levels of IL-6 and VEGF were down-regulated, whereas TNF-α and caspase-3 were increased after treatment of GEE *in vivo*. **Conclusion:** Taken together, these findings suggest GEE has significant anti-liver cancer activities both *in vitro* and *in vivo*, which may be involved in antiangiogenesis and induction of apoptosis.

Keywords: Gekko ethanol extract; apoptosis; antiangiogenesis; hepatocellular cancer

S1.8

Potentiating effect of citral on anticancer activities of doxorubicin, vincristine, and etoposide in human B-lymphoma cells

Darinee DANGKONG¹, Wacharee LIMPANASITHIKUL². ¹Interdisciplinary Program of Pharmacology, Graduate School, Chulalongkorn University, Bangkok, Thailand; ²Department of Pharmacology, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

Aim: Citral is a mixture of terpene aldehydes in essential oils of several plants used for cooking such as lemongrass. This study intended to evaluate the potentiating effect of citral on anticancer activities of doxorubicin, vincristine, or etoposide

against human B-lymphoma Ramos cells. **Methods:** Ramos cells were treated with each anticancer drug in the presence and the absence of citral. Cytotoxicity of these agents and the combination was determined by resazurin reduction assay. Proliferation and apoptosis of the treated cells were evaluated by an automated cell counter and AnnexinV/FITC and PI staining, respectively. The expression of Bcl-2 family proteins in the treated cells was performed by qPCR. **Results:** Citral had cytotoxicity on Ramos cells with IC₅₀ value 85.6 μmol/L. At citral 10-40 μmol/L significantly increased the cytotoxic effects of doxorubicin, vincristine, and etoposide on Ramos cells, leading to the decrease in the IC₅₀ value of each drug. It potentiated the apoptotic induction effects of these anticancer drugs. The combination of citral with doxorubicin significantly decreased the expression of anti-apoptotic Bcl-x_L when compared to doxorubicin alone. It did not change anti-proliferative effects of these drugs. **Conclusion:** Citral had potentiating effect on the Ramos cell cytotoxicity of doxorubicin, vincristine, and etoposide by increasing in apoptotic effects of these drugs. The results from this study strongly suggest that citral in several edible oils may have beneficial effect in patients with B cell lymphoma treated with chemotherapy.

Keywords: citral; anticancer; human B-lymphoma Ramos cells

S1.9

Differential protein expression of plumbagin treated CCA cell line

Veerachai EURSITTHICHAI¹, Bodin SROYBUDDA², Egkaluck NGAMLOON², Panita SOONTORNNON², Kanpriya PHEWNUAL², Tullayakorn PLENGSURIYAKARN¹, Kanawut KOTAWONG¹, Sitthirak ROYTRAKUL³, Naruemol PHAONAKROP³, Vithoon VIYANANT¹, Kesara NA-BANGCHANG¹. ¹International College of Medicine, Thammasat University, Pathumthani 12121, Thailand; ²Department of Medical Technology, Faculty of Allied Health Sciences, Thammasat University, Pathumthani 12121, Thailand; ³Center for Genetic Engineering and Biotechnology (BIOTEC), National Science and Technology Development Agency, Pathumthani 12120, Thailand

Aim: Medicinal plant extracts are alternative sources of promising anti-cancer compound such as *Plumbago* extracts. Therefore, this study aims to investigate the effect of plumbagin on protein expression in CCA cell line namely CL6. **Methods:** The CL6 cells were treated with plumbagin compound for 48 h. The cells were harvested at 0, 1, 3, 6, 12, 24, and 48 h, protein extracted, and subjected for LC-MS/MS analysis. **Results:** Differential protein expression has been observed. To identification of up- and down-regulated protein will be analyzed by using DeCyder, MASCOT, and Mev software. **Conclusion:** This protein profile will be useful for anti-CCA drug development.

Keywords: plumbagin; cholangiocarcinoma; protein expression; LC-MS/MS

S1.10

12-O-tetradecanoylphorbol-13-acetate increases sensitivity of cisplatin to human lung adenocarcinoma cells

Lei FAN¹, Yu-zhong GUO^{1,*}, Hong ZHANG², Zheng-tao HAN². ¹Department of Clinical Pharmacology, School of Medicine, Zhengzhou University, Zhengzhou 450052, China; ²Tumor hospital of Henan Province, Zhengzhou 450008, China

*To whom correspondence should be addressed:

E-mail: gyzh@zzu.edu.cn

Aim: To investigate whether 12-O-tetradecanoylphorbol-13-acetate (TPA) can increase the sensitivity of cisplatin on human lung adenocarcinoma cisplatin-sensitive cell line A549 and cisplatin-resistant cell line A549/CP. **Methods:** MTT assay was used to observe cell proliferation after treating by TPA alone or in combination with cisplatin. Intracellular cisplatin accumulation was measured by flameless atomic absorption spectroscopy. Specific activities of Na⁺, K⁺-ATPase were detected according to the procedure of Na⁺, K⁺-ATPase activity detection kit. Assesment of Na⁺, K⁺-ATPase α and β subunits was done by Western blotting. **Results:** TPA at 8.1 nmol/L or lower didn't affect the cells' proliferation. TPA decreased IC₅₀ of cisplatin by 77% in A549 cells and 69% in A549/CP cells at 8.1 nmol/L after 96 h respectively. Intracellular cisplatin accumulation was increased after pretreated with TPA for 72 h in both cell lines. However, pretreatment with Na⁺, K⁺-ATPase activity inhibitor ouabain and TPA markedly reduced intracellular cisplatin accumulation. Na⁺, K⁺-ATPase activity was enhanced with increasing concentrations of TPA in all the cells. Western blotting results showed that Na⁺, K⁺-ATPase β1 subunit was induced in both cell lines. **Conclusion:** These results suggest that pretreatment of A549 and A549/CP cells with TPA increases the sensitivity of cisplatin by intracellular cisplatin accumulation possibly correlated with Na⁺, K⁺-ATPase.

Keywords: 12-O-tetradecanoylphorbol-13-acetate (TPA); cisplatin; lung adenocarcinoma cells; Na⁺, K⁺-ATPase

S1.11**Chrysin enhances sensitivity of BEL-7402/ADM cell to doxorubicin by suppressing PI3K/Akt/Nrf2 and ERK/Nrf2 pathway**

Ai-mei GAO^{1,2}, Zun-ping KE², Fang SHI³, Hui CHEN^{1*}. ¹Department of Pharmacology, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China; ²Department of Pharmacy, the Fifth People's Hospital of Shanghai, Fudan University, Shanghai, China; ³Department of Cardiology, the Fifth People's Hospital of Shanghai, Fudan University, Shanghai, China; ⁴Laboratory of Chinese Herbal Pharmacology, Renmin Hospital, Hubei University of Medicine, Shiyan, China

*To whom correspondence should be addressed.

Aim: Nuclear factor-E2-related factor 2 (Nrf2) is a redox-sensitive transcription factor regulating expression of a number of cytoprotective genes. Recently, Nrf2 has emerged as an important contributor to chemoresistance in cancer therapy. Here, we investigated whether Nrf2 was associated with drug resistant in doxorubicin resistant BEL-7402 (BEL-7402/ADM) cells, and if chrysin could reverse drug resistance in BEL-7402/ADM cells. **Methods:** Reverse of drug resistance was assayed by MTT and Fluorometric analysis. Inhibition of Nrf2 signaling pathway by chrysin was detected through real-time quantitative PCR and western blot analysis. **Results:** We found that notable higher level of Nrf2 and its target proteins in BEL-7402/ADM cells compared to BEL-7402 cells. Similarly, intracellular Nrf2 protein level was significantly decreased in BEL-7402/ADM cells and ADM resistance was partially reversed by Nrf2 siRNA. Chrysin is a potent Nrf2 inhibitor which sensitizes BEL-7402/ADM cells to ADM and increases intracellular concentration of ADM. Mechanistically, chrysin significantly reduced Nrf2 expression at both the mRNA and protein levels through down-regulating PI3K-Akt and ERK pathway. Consequently, expression of Nrf2-downstream genes HO-1, AKR1B10, and MRP5 were reduced and the Nrf2-dependent chemoresistance was suppressed. **Conclusion:** The results suggested that activation of Nrf2 is associated with drug resistance in BEL-7402/ADM cells and chrysin may be an effective adjuvant sensitizer to reduce anticancer drug resistance by down-regulating Nrf2 signaling pathway.

Keywords: chrysin; Nrf2; hepatocellular carcinoma; sensitivity; chemoresistance

Acknowledgements: We thank Professor XB WANG (Laboratory of Chinese Herbal Pharmacology, Hubei University of Medicine) for the generous gift of BEL-7402 and BEL-7402/ADM cells.

S1.12**Anticancer activity of celastrol in hepatocellular carcinoma**

Wei HE, Sha-sha SONG, Jing-tao LU*, Wei WEI*. *Institute of Clinical Pharmacology, Anhui Medical University, key Laboratory of Anti-inflammatory and Immune Medicine (Anhui Medical University), Ministry of Education, Hefei 230032, China*

*To whom correspondence should be addressed.

E-mail: lujt18@126.com (Jing-tao LU); wwei@ahmu.edu.cn (Wei WEI)

Aim: Celastrol is an active ingredient of the traditional Chinese medicinal plant *Tripterygium Wilfordii*, which has anti-inflammatory and anticancer activities. The objective of the study was to evaluate the anticancer effects of celastrol in hepatocellular carcinoma. **Methods:** Sprague-Dawley rats were intragastrically administered with 0.2% (*w/v*) DEN every 5 d for 16 wk (10 mg/kg) and maintained for an additional 4 weeks without administration of DEN, and were orally fed with celastrol since 10th week for 10 weeks. In addition, Bel-7402 cells, HCC cell line, were used to further explore the molecular mechanisms related to anticancer of celastrol, significantly model group celastrol showed a marked reduction in the above condition and subsequently. **Results:** Results showed that celastrol significantly reduced the numbers and volume of live tumor foci in DEN-induced HCC rats, decreased obviously the levels of AST, ALT, ALP and AFP in rat serum elevated by DEN. Meanwhile, celastrol induced apoptosis in Bel-7402 cells in dose- and time-dependent manners. Celastrol increased the expression of p53, inhibited the activity of MDM2, and elevated the ratio of Bax/Bcl-2 in Bel-7402 cells and in DEN induced rat liver. Furthermore, Celastrol induced the release of *cytochrome c*, increased the expression of FAS, FasL, cleavage of caspase-3, caspase-8 and caspase-9 in Bel-7402 cells. **Conclusion:** This study indicates that celastrol could prevent the progression of HCC development and its anticancer mechanisms is related to the induction of apoptosis, ie, celastrol may be a potential chemotherapy agent for HCC.

Keywords: celastrol; hepatocellular carcinoma; diethylnitrosamine; apoptosis; p53

S1.13**Role of osteopontin in human osteosarcoma**

Ishan HSIEH¹, Houngchi LIOU¹, Rongsen YANG², Wenmei FU¹. ¹Department of

Pharmacology, College of Medicine, National Taiwan University, Taipei 100, Taiwan, China; ²Department of Orthopedic Surgery, National Taiwan University Hospital, Taipei 100, Taiwan, China

Aim: Osteopontin (OPN), an extracellular matrix protein, has a functional RGD domain for binding to integrin and has been reported to be associated with malignant transformation. OPN plays a crucial role in tumor progression, invasion and metastasis. Here we investigated the effect of osteopontin on the regulation of glucose transporters in osteosarcoma. **Methods:** Protein expression was assayed by Western blot. Cell viability was assayed by MTT. Glucose uptake was assayed by flow cytometry. Cell motility was assayed by trans well migration assay. **Results:** We found that OPN and glucose transporters expression were upregulated by hypoxia. The expression of Type I glucose transporters can be increased by osteopontin through $\alpha v \beta 3$ integrin and CD44 receptor. PI3K-Akt, p38, and JNK pathways are involved in the signaling of osteopontin. OPN can also enhance migration of osteosarcoma cells migration. **Conclusion:** Osteopontin can regulate the expression of glucose transporter and control the supply nutrient, which will influence the viability of osteosarcoma cells. Osteopontin seems a good drug target for osteosarcoma therapy.

Keywords: integrin; osteopontin; osteosarcoma

S1.14**Involvement of the prostaglandin E receptor, EP₂, in paeoniflorin-induced human hepatoma cells apoptosis**

Shan-shan HU, Di WANG, Juan JIN, Jing-jing WU, Jing-yu CHEN, Hua-xun WU, Qing-tong WANG, Wu-yi SUN*, Wei WEI*. *Institute of Clinical Pharmacology, Anhui Medical University, key Laboratory of Anti-inflammatory and Immune Medicine (Anhui Medical University), Ministry of Education, Hefei 230032, China*

*To whom correspondence should be addressed.

E-mail: sunwuyi51@yahoo.com.cn (Wu-yi SUN); wwei@ahmu.edu.cn (Wei WEI)

Aim: Prostaglandin E₂ (PGE₂) has been shown to play an important role in tumor development and progression. PGE₂ mediates its biological activity by binding any one of four prostanoid receptors (EP₁ through EP₄). The present study was designed to investigate the role of the EP₂ receptor during proliferation and apoptosis of human HepG2 and SMMC-7721 hepatoma cell lines and the effect of paeoniflorin, a monoterpene glucoside. **Methods:** The proliferation of HepG2 and SMMC-7721 cells was determined by methyl thiazolyl tetrazolium (MTT) after exposure to the selective EP₂ receptor agonists butaprost and paeoniflorin. Apoptosis of HepG2 and SMMC-7721 cells were also quantified by flow cytometry with annexin V-fluorescein isothiocyanate (FITC) and propidium iodide (PI) staining. The expression levels of Bcl-2 and Bax were quantified by western blotting and immunohistochemistry. The expression of the EP₂ receptor and cysteine-aspartic acid protease (caspase)-3 was determined by Western blotting. **Results:** Butaprost significantly increased proliferation in HepG2 and SMMC-7721 cells. Paeoniflorin significantly inhibited HepG2 and SMMC-7721 cells proliferation stimulated by butaprost at multiple time points (24, 48, and 72 h). Paeoniflorin induced apoptosis in HepG2 and SMMC-7721 cells that was quantified by annexin-V and PI staining. Our results indicate that the expression of the EP₂ receptor, Bcl-2 was significantly increased, Bax, cleaved caspase-3 was decreased in HepG2 and SMMC-7721 cells after stimulated by butaprost. Paeoniflorin significantly decreased the EP₂ receptor, Bcl-2 expression and increased Bax and caspase-3 activation in HepG2 and SMMC-7721 cells addition with butaprost. **Conclusion:** Our results suggest that the PGE₂ receptor subtype EP₂ may play a vital role in cell survival in both HepG2 and SMMC-7721 cells. Paeoniflorin, which may be a promising agent in the treatment of liver cancer, induced apoptosis in HCC cells by downregulating EP₂ expression and increased the Bax-to-Bcl-2 ratio, thus up-regulating the activation of caspase-3.

Keywords: paeoniflorin; butaprost; EP₂; hepatocellular carcinoma; apoptosis; caspase-3

Acknowledgements: This work was supported by grants from Specialized Research Fund for the Doctoral Program of Higher Education of China (No 20113420120002), Natural Science Foundation of the Higher Education Institutions of Anhui Province (No KJ2012A153), the National Natural Science Foundation of China (30973543, 81173075). The authors acknowledge the help of the staff members of the Institute of Clinical Pharmacology, Anhui Medical University in conducting the study.

S1.15**Anti-tumor effect of three kinds of traditional chinese medicine formula on S₁₈₀-bearing mice**

Zhi-ping HU, Xing-feng CHANG, Chun-hua SONG, Bao-ying LUO, Jie WANG, Li-ming HAN*. *Gannan Medical University, Ganzhou 341000, China*

*To whom correspondence should be addressed.

E-mail: jxhlm@163.com

Aim: To study the anti-tumor effect of three kinds of traditional Chinese medicine (TCM) and discuss the mechanisms about sarcoma cell apoptosis and angiogenesis. **Methods:** The second day of S_{180} -bearing mice model established, mice were randomly divided into control group, model group, cyclophosphamide (CTX) group (50 mg/kg) and three TCM groups. After 10 days of different treatments, the tumor inhibition rate were measured, and the expression of Bcl-2, Bax, caspase-3 and VEGF protein in the tumor tissues were investigated by Western blot. **Results:** Bie Jia Jian Wan (BJJW) and Liu Wei (LW), especially BJJW, can significantly affect the tumor growth with 35.9% and 22.0%, respectively. While Fu Zhen TCM has no anti-tumor effect in S_{180} -bearing mice. Compared with the model group, BJJW and LW group mice showed decreases Bcl-2 and VEGF, and increases Bax and caspase-3 expression in S_{180} sarcoma. **Conclusion:** BJJW and LW have anti-tumor effects in S_{180} -bearing mice. Its mechanisms probably related with the induction of the apoptosis of S_{180} sarcoma cells through downregulation of Bcl-2 and upregulation of Bax and caspase-3 expression, and disturb the tumor angiogenesis via downregulation of VEGF.

Keywords: traditional Chinese drugs; tumor-bearing mice; anti-tumor; apoptosis; angiogenesis

Acknowledgements: This work was supported by grants from the Jiangxi Natural Science Foundation (20122BAB205079).

S1.16

Anti-angiogenic mechanisms of PPemd 25, a novel anthraquinone derivative

Shiu-Wen HUANG¹, Jin-Cherng LIEN², Sheng-Chu KUO², Tur-Fu HUANG¹. ¹Graduate Institute of Pharmacology, College of Medicine, National Taiwan University, Taipei, Taiwan, China; ²Graduate Institute of Pharmaceutical Chemistry, China Medical University, Taichung, Taiwan, China

Aim: Angiogenesis contributes to ischemic, inflammatory, infectious and immune disorders and also occurs during numerous malignancies. It is a balanced process that involves proliferation, migration, differentiation and tube formation of endothelial cells. As a critical factor in inducing angiogenesis, vascular endothelial growth factor (VEGF) has become an attractive target for anti-angiogenesis treatment. **Methods:** In an effort to develop inhibitors to block VEGF signaling and angiogenesis, we selected PPemd25, an anthraquinone derivative, and investigated its inhibitory mechanisms using human umbilical endothelial cells (HUVECs). **Results:** PPemd25 concentration-dependently inhibited VEGF-induced proliferation, migration and tube formation of HUVECs. PPemd25 also suppressed *ex vivo* VEGF-induced microvessel sprouting from aortic rings. In addition, PPemd25 inhibited VEGF-induced phosphorylation of focal adhesion kinase (FAK), Akt and ERK. Using an implanted matrigel plugs angiogenesis model, PPemd25 attenuated MDAMB-231 breast cancer cells-induced angiogenesis. **Conclusion:** Together our findings suggest that PPemd25 inhibits VEGF-induced angiogenesis via downregulation of VEGFR2-mediated signaling. Results of this study suggest that PPemd25 may be a potential drug candidate in anti-cancer therapy, acting by suppressing tumor-induced angiogenesis.

Keywords: angiogenesis; endothelial cells; vascular endothelial growth factor (VEGF)

S1.17

WMJ-S-001, a novel HDAC inhibitor, induces HCT116 colorectal cancer cell death through AMPK/p38MAPK signaling cascade

Yu-Han HUANG¹, Pei-Ting CHIU¹, Wei-Jan HUANG², Ming-Jen HSU¹. ¹Department of Pharmacology, School of Medicine, College of Medicine, Taipei Medical University, Taipei, Taiwan, China; ²Graduate Institute of Pharmacognosy, College of Pharmacy, Taipei Medical University, Taipei, Taiwan, China; ³Department of Surgery, Landseed Hospital, Taoyuan, Taiwan, China

Aim: Histone deacetylases (HDACs) are often found over-expressed in human cancers, including colorectal cancer, and has been implicated in tumorigenesis. HDAC inhibition induces growth arrest and cell death in various transformed cell. However, the mechanisms by which this reduces cell viability in colorectal cancer cells remain to be elucidated. **Methods:** We explored the actions of a novel HDAC inhibitor, WMJ-S-001 in HCT116 colorectal cancer cells. **Results:** WMJ-S-001 dose-dependently decreased cell viability and proliferation in HCT116 cells. These results are associated with the modulation of survivin, Bax and p21. WMJ-S-001 activated AMP-activated protein kinase (AMPK) and p38 mitogen-activated protein kinase (p38MAPK), whereas an inhibitor of AMPK or p38MAPK signaling abrogated WMJ-S-001's effects of increasing Bax and p21 levels. In addition, WMJ-S-001 induced p53 phosphorylation in a time-dependent manner. Furthermore,

WMJ-S-001 also suppressed tumor growth in HCT116 colorectal tumor xenograft model. **Conclusion:** WMJ-S-001 may cause AMPK- and p38MAPK-mediated p53 activation, leading to cell death in HCT116 colorectal cancer.

Keywords: colorectal cancer; HDAC; AMPK; p38MAPK

S1.18

The study for apoptosis of human cervical cancer HeLa cell induced by celosin A from Celosiae Semen and its mechanisms

Zhi-qin HUANG, Qi-lai CHENG, Hong-liang LI, Cheng HUANG. College of Pharmacy, Gannan Medical University, Ganzhou 341000, China

Aim: To investigate the apoptosis of human cervical cancer HeLa cell induced by celosin A (CA) from the semen of *Celosia argentea* L and its mechanisms. **Methods:** The viability of HeLa cells was detected by the method CCK-8. The activity of LDH was measured by MTT assay. Immune colorimetry was used to test the proliferation of HeLa cells. The change of morphology of apoptosis HeLa cells was observed by DAPI fluorescence staining. Western blot was used to detect the expressions of caspase-3/7 and NF- κ B proteins of HeLa cells. **Results:** After treatment with CA, the leakage of LDH increased at dosage and time dependently. In the process of determination of BrdU and CCK-8, the inhibitory effect of CA to HeLa cell was stronger than that of the control, and showed clear inhibitory effect on the HeLa cell proliferation with the increasing of concentration and the acting time. By using the DAPI fluorescence dyes, HeLa cell cytoplasm concentrating, nucleolus dispersion and chromatin agglomeration were observed. The Western blot analysis showed that the expression of caspase-3/7 was significantly enhanced and the expression of NF- κ B proteins was down regulation. **Conclusion:** These results suggested that celosin A could induce the apoptosis of HeLa cells and its mechanisms for activating the activity of caspase-3/7 and down-regulating the expression of NF- κ B protein.

Keywords: *Celosia argentea* L; celosin A; HeLa cells; apoptosis; NF- κ B

Acknowledgements: This work was supported by the grant from Natural Science Foundation of Jiangxi Province (2012ZBAB205006)

S1.19

A betulinic acid derivative, SYK023, inhibits lung cancer formation

Jan-Jong HUNG¹, Wen-Chang CHANG². ¹Institute of Bioinformatics and Biosignal Transduction, College of Bioscience and Biotechnology, National Cheng-Kung University, Tainan 701, Taiwan, China; ²Graduate Institute of Medical Sciences, College of Medicine, Taipei Medical University, Taipei, Taiwan, China

Aim: Our previous study indicated that betulinic acid (BA) could inhibit lung cancer formation *via* inhibiting Sp1 level. Herein we try to screen BA derivative(s), which is more effective than BA itself. **Methods:** Various BA derivatives were used to treat the lung cancer cell lines at first to evaluate the therapeutic effect of these compounds, and then used the compound(s) effected in the cellular level to do the animal experiments including the xenograph and spontaneous induced lung cancer animal. Finally, the therapeutic molecular mechanism of the BA-derivative in lung cancer will be addressed. **Results:** After screen more than thirty BA derivatives in inhibiting the growth of lung cancer cells, several BA derivatives contain more cytotoxicity than BA itself in lung cancer cells. Moreover, cell viability by those BA derivatives was also decreased more than BA treatment by MTT assay. Subsequently, three more effective BA derivatives (SYK010, SYK019 and SYK023) were chose to study further. At first, that Sub-G₁ was found more in these BA derivatives treated cells than that by BA treatment. Second, these BA derivatives also affect the Sp1 level like BA treatment. Finally, the effects of these BA derivatives were then evaluated in animal experiments. We not only use the xerograph study, but also use the spontaneous induced lung cancer animal model, Kras^{G12D}-induced lung cancer mice, to address this issue. **Conclusion:** These results indicate that compound, SYK023, was more effective in preventing cancer formation than BA or SYK019 treatment, which might be benefit for the cancer therapy in the future.

Keywords: betulinic acid; SYK023; Sp1; lung cancer

S1.20

Cotargeting autophagy and hsp70 for breast cancer treatment

Xiao-xiao JIANG, Bao YONG, Yong-hua YANG*. School of Pharmacy Fudan University, Shanghai 201203, China

*To whom correspondence should be addressed.

Autophagy is a conserved catabolic pathway in which autophagosomes fuse with lysosomes for protein degradation and amino acid recycling. The initiation of autophagy is triggered by a lipid kinase (PI3KC3) signaling complex, consisting of

Vps34, Vps15, and beclin1. Treatment of breast cancer cells with anti-tumor drugs inducing the metabolic and protein-misfolding stress results in heat shock proteins (hsp) expression and induction of autophagy to avoid cell death.

Hsp70 is a stress-activated multifunctional chaperone containing cytoprotective function to inhibit cell apoptosis and cause resistance in breast and other cancers. Suppression of Hsp70 sensitizes cancer cells to chemotherapy. In our previous studies, treatment of breast cancer MCF-7 cells with HDAC inhibitor LBH589 induces autophagosome formation accompanied by hyperacetylation of hsp70 which enhances the stability and activity of Vps34, via kap-1 dependent sumoylation, leading to the formation of VPS34-Beclin1 complex.

Recently, Chloroquine, a widely used antimalarial agent, has been revealed to have promising effects on cancer treatment in clinical trials. Here, we report that co-treatment with LBH589 and chloroquine or hsp70 inhibitor Pifithrin- α (PFT α) significantly enhanced lethality of breast cancer MB-231, MB-468, and MCF-7 cells. We found that both chloroquine and PFT α inhibited autophagy activation and hsp70 expression following LBH589-induced autophagy protecting cell from apoptosis. Importantly, combinatorial treatment with LBH-589 with hsp70 inhibitor chloroquine or PFT α resulted in the abrogation of LBH589-induced interaction of hsp70 with VPS34, and the sumoylation and activity of Vps34. Both chloroquine and PFT α targeting Hsp70 significantly delayed breast cancer growth *in vivo*.

Our findings demonstrate that a new strategy using agent, such as chloroquine and PFT α , cotargeting hsp70 and Vps34 has the potential benefits to improve treatment outcomes in cancer.

S1.21

The anti-proliferation effect of FSEO on human colorectal cell line HCT116 *in vivo* and *in vitro*

Zhong-liang JIANG¹, Fu-qiang REN², Si-si WANG¹, Bao XU¹, Ling ZHU*. ¹Department of Pharmacology, School of Preclinical and Forensic Medicine, Sichuan University, Sichuan 610041, China; ²Department of thoracic, Number Three Hospital of Chengdu, Chengdu 610031, China

*To whom correspondence should be addressed.

E-mail: zhuling529@163.com

Aim: To investigate the inhibitory effect and the anti-tumor mechanism of FSEO on human colorectal HCT116 cancer cell line *in vitro* and *in vivo*. **Methods:** Their morphological and cellular characteristics were observed by Wright Giemsa staining and HE staining. Growth inhibition was analyzed with MTT assay. Establish the xenograft tumor model of HCT116 cell line in naked mice. The expression level of mRNA or protein *in vitro* and *in vivo* (tumor tissue) was measured by RT-PCR or Western blotting. siRNA was used to silence the mRNA expression of p53 gene in HCT116 cell line *in vitro*. After p53 gene been silenced, cells were treated by high dose of FSEO for 24h. Then the mRNA expression was tested via RT-PCR method. **Results:** After treated with FSEO for 24 to 48 h, the proliferation of different cell lines were inhibited, and there was obvious dose-effect relationship of the inhibitory effect, IC₅₀ of FSEO to HCT116 cells was 115.17±14.42 mg/mL. Apoptosis phenomenon in HCT116 cell line or in tumor tissue sections was observed after treated with different concentrations of FSEO. The *in vivo* anti-tumor trial illustrated all the tested doses of FSEO (1, 2, and 4 g/kg) had inhibitory effects on xenografted HCT116 tumor in naked mice. The inhibitory rates of each dose group are: 21.86% ($P<0.05$), 33.16% ($P<0.05$), 47.31% ($P<0.05$). There was a reduction of Bcl-2 and Cyclin d1 expression, up-regulation of p53 and Bax expression, down turn the Bcl-2/Bax protein ratio in FSEO groups under both *in vivo* and *in vitro* conditions ($P<0.05$). RNA interference trial results showed FSEO may via p53 dependent pathway to regulate the expression of Bcl-2 and Bax to induce the apoptosis. **Conclusion:** FSEO had antiproliferative activity against colorectal I cancer *in vitro* and *in vivo*. Its mechanism may be involved in apoptosis of cells induced by down-regulation of oncogenes and up-regulation of apoptotic genes.

Keywords: FSEO; HCT116 cell line; Apoptosis

S1.22

Effect of resveratrol on RUNX3 gene in B16F10 murine melanoma cell

Si-si KANG, Xin-xia LIU, Chang MA, Wen-juan HE, Hui CHEN*. Department of Pharmacology, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, China

*To whom correspondence should be addressed.

Aim: To explore the effect and mechanism of RES on RUNX3 gene in B16F10 cell line. **Methods:** The B16F10 cell line were treated with DMSO, different concentration of RES (20, 40, and 80 $\mu\text{mol/L}$) and positive compound [(5-Aza-CdR),

10 $\mu\text{mol/L}$] for 48 h. The difference of mRNA expression and protein expression and the promoter region methylation status of the RUNX3 gene in B16F10 cell were measured by using RT-PCR, Western-Blotting and MSP, respectively. **Results:** Loss expression of RUNX3 mRNA and protein were found in the B16F10 cell. The methylation inhibitor 5-Aza-CdR up-regulates the expression of RUNX3 mRNA and protein in the B16F10 cell. After the B16F10 cell treated by different concentrations of RES, expression of RUNX3 mRNA and protein were significantly up-regulated in a dose-dependent manner. RUNX3 gene promoter was hypermethylated in untreated B16F10 cell. After treated by RES, the promoter of RUNX3 gene was demethylated partly and such effect was concentration-dependent. **Conclusion:** RES could increase the expression of RUNX3 in melanoma B16F10 cells, suggesting that one of the potential mechanisms may be related with Demethylation of RES.

Keywords: Resveratrol; melanoma; RUNX3; Methylation

S1.23

Anticancer activity of cyclooxygenase-2 inhibitor on breast cancer *in vitro*

Jiraporn KUESAP¹, Thitiya TRAICHOKGUL¹, Atitaya CHOMPBUCHAN¹, Wanna CHAIJAROENKUL², Kesara NA-BANGCHANG². ¹Graduate Program in Biomedical Sciences, Faculty of Allied Health Sciences, Thammasat University, Thailand; ²International College of Medicine, Thammasat University, Thailand

Aim: Cyclooxygenase-2 (COX-2) plays a key role in breast cancer progression and metastasis by promotes carcinogenesis, tumor proliferation, angiogenesis, prevention of apoptosis, and immunosuppression. This study investigated the antitumor effects of celecoxib, the selective COX-2 inhibitor, and 5-fluorouracil on breast cancer cell *in vitro*. **Methods:** Human breast cancer HCC1937 cell (ATCC cat.no. CRL-2336) was tested with eight different concentrations of celecoxib or 5FU. The chemosensitizer effect of celecoxib and 5-fluorouracil was performed at fixed celecoxib 50 $\mu\text{mol/L}$ with different concentrations of 5FU. The MTT assay was used to determine the growth inhibition of breast cancer cells *in vitro*. **Results:** The proliferation of HCC1937 cell was inhibited by celecoxib with dose dependent manner. The concentration that inhibited cell growth by 50% (IC₅₀) of 5FU and celecoxib were 5642 and 259 $\mu\text{mol/L}$, respectively. However, celecoxib does not reverse 5-fluorouracil sensitivity. **Conclusion:** Result suggests that celecoxib may be a potential candidate for breast cancer chemotherapy but could not use with 5FU.

Keywords: cyclooxygenase-2 inhibitor; breast cancer; celecoxib

Acknowledgements: The study was supported by Thammasat University, The Commission on Higher Education, Ministry of Education of Thailand, The National Research University Project of Thailand (NRU).

S1.24

Anticancer activity of the novel histone deacetylase inhibitor, MPT0G028, in human colorectal cancer cells *in vitro* and *in vivo*

Chia-Hua LEE¹, Chun-Han CHEN¹, Shioh-Lin PAN², Che-Ming TENG¹. ¹Pharmacological Institute, College of Medicine, National Taiwan University, Taipei, Taiwan, China; ²The PhD program for Cancer Biology and Drug Discovery, College of Medical Science and Technology, Taipei Medical University, Taipei, Taiwan, China

Aim: Overexpression of histone deacetylases (HDACs) were observed in many types of cancers, including colorectal cancer (CRC). Therefore, small molecules targeting HDACs may have benefits in developing new anticancer drugs for CRC patients. So, we investigated the anticancer activity of a novel synthetic HDAC inhibitor, MPT0G028, in CRC *in vivo* and *in vitro*. **Methods:** Anticancer activity was evaluated by SRB and MTT in HCT-116 and HT-29 CRC cell lines. Cell cycle distribution was accessed by flow cytometry and protein expression was examined by Western blotting. HDAC activities were measured by fluorogenic HDAC assay kit. HCT-116 xenograft model was used to examine the antitumor activity *in vivo*. **Results:** MPT0G028 showed a better anti-proliferation effect than vorinostat in human CRC and exhibited significant inhibitory effects against class I HDACs by using *in vitro* biochemical assay and repressed total HDAC enzymatic activity in HCT-116 cells. Moreover, MPT0G028 can induce apoptotic cell death in a time- and concentration-dependent manner. The data also showed that the repression of Wnt/ β -catenin pathway might involve in MPT0G028-induced apoptosis. Finally, MPT0G028 has a potent inhibitory effect on the tumor growth in the HCT-116 xenograft model. **Conclusion:** MPT0G028 is a novel HDAC inhibitor with a better potency than the FDA-approved HDAC inhibitor, vorinostat, and induces dramatic apoptosis in CRC cells. We hope that this study will have contributions to the new drug development and clinical benefits for CRC patients.

Keywords: histone deacetylases (HDACs); MPT0G028; vorinostat; Wnt/ β -catenin; colorectal cancer

S1.25**Study on the effect and its mechanism of Xihuang pill reversing tumor immune escape**

Wen-bo LIANG¹, Jie MA², Shuo GUAN³, Wei YANG³, Jun-xia HU³, Wen-bin GAO⁴, Chang-qian ZENG¹. ¹Department of pharmacology, Medical college of Dalian University, Dalian 116622, China; ²Department of pharmacology, Zunyi Medical College, Zunyi 563003, China; ³Department of Oncology, Medical college of Dalian University, Dalian 116622, China; ⁴Department of Oncology, Affiliated Zhongshan Hospital of Dalian University, Dalian 116001, China

Aim: To investigate the effect and mechanism of Xihuang Pill reversing tumor immune escape of tumor-bearing mice tumor immune escape. **Methods:** 4T-1 tumor cell were inoculated into right armpit of BALB/c mouse and randomly divided into 6 groups: normal control group, model control group, positive control group (matrine) and Xihuang pill low dose group, mid dose group, high dose group. After the drug administration in 14 consecutive days, the inhibitory rate of tumor was observed. The tumor tissue was Stripping and weighing, the tumor-inhibitory rate was calculated. The level of IL-2, IL-6, IL-10, TGF- β , VEGF, IFN- γ in Serum were determined by enzyme-linked immunosorbent assay (ELISA). Adhesion molecule B7-1 (CD80) and Treg, MDSC of peripheral blood were determined by flow cytometry (FCM). **Results:** The tumor-inhibitory rate of Xihuang pill at high dose group was 36.54 percent. Compared with the model group, the average level of IL-2, IFN- γ and the number of B7-1 of the Xihuang pill large dose group significantly up-regulated, however the number of IL-6, IL-10, TGF- β , VEGF, and Treg, MDSC significantly decreased ($P < 0.05$). **Conclusion:** Xihuang pill can significantly inhibits tumor growth by interfere the tumor immune escape, the mechanism relate to the form of tumor immunosuppressive microenvironment by reducing the level of immunosuppressive factors and enhancing immune clearance.

Keywords: Xihuang pill; 4T1 cell; BALB/c mice; tumor microenvironment; immune escape

S1.26**Anti-cancer effects of *Chaenomeles speciosa* flavonoids in murine sarcoma 180 cells**

Chao-qi LIU¹, Xiao-lin QIN¹, Wei WU², Xiao-hua ZOU², Ding YUAN²*. ¹Institute of Molecular Biology of Three Gorges University, Yichang 443002, China; ²Medical College of Three Gorges University, Yichang 443002, China

*To whom correspondence should be addressed.

Aim: In recent years, the use of natural plant products has intensively attracted attention among clinicians and the public for the prevention and treatment of various chronic diseases. *Chaenomeles speciosa* (Sweet) Nakai is one of the natural plants with high medicinal and dietary value, and comprised of amounts of various effective ingredients such as the total flavonoids, organic acid, and so forth, is originally from Anhui, Hubei and Sichuan of China. This investigation evaluates anti-tumor activity and its mechanism of total flavonoids from *Chaenomeles speciosa* (Sweet) Nakai (FLC). **Methods:** The sarcoma 180 (S180)-bearing mice were used to study potential anti-cancer activity of FLC. The tumor size, survival rate and immune responses were measured in S180 solid and ascites tumor. Molecular screening model of PD-1/PD-L1 was established and the FLC was used to investigate the inhibite activity. Lymphocyte proliferation was also assayed for the extract FLC. **Results:** Our data showed that the FLC could significantly inhibit the tumor growth and prolong the survival time in the S180 tumor bearing mice. The lymphocyte proliferation assay showed that the FLC could improve the specific lymphocyte proliferation. In the S180 ascites, the FLC enhanced tumor infiltrating lymphocytes (TIL) and inhibited programmed death-1-ligand 1 (PD-L1) expression. Using molecular binding assay FLC could inhibit the binding of PD-L1 to PD-1. **Conclusion:** These results suggest that FLC exhibited anti-cancer and immunomodulatory activities, which improved natural and adaptive immune functions, such as interfering PD-1/PD-L1 pathway.

Keywords: *Chaenomeles speciosa* (Sweet) Nakai; flavonoids; anti-cancer activity; immune regulation

S1.27**Effect of S003 on sulfur mustard-induced cutaneous injuries in the mouse ear vesicant model and human HaCaT cells**

Feng LIU, Ning JIANG, Zhi-yong XIAO, Xin-bo ZHOU, Jun-ping CHENG, Yi-zhou MEI, Xiaorui ZHANG, Pan ZHENG, Wen-xia ZHOU*, Yong-xiang ZHANG*. Beijing Institute of Pharmacology and Toxicology, Beijing 100850, China

*To whom correspondence should be addressed.

E-mail: zhouwx@bmi.ac.cn (Wen-xia ZHOU); zhangyx@bmi.ac.cn (Yong-xiang ZHANG)

Aim: Sulfur mustard (SM) can cause DNA damage and the activation of poly(ADP-

ribose)polymerase (PARP). PARP inhibitor has been reported some protective effects in SM skin injury *in vivo* and *in vitro*. S003 is a PARP inhibitor synthesized by our institute. The aim of our study is to observe the effect of S003 in SM treated HaCaT cells and in SM induced mouse ear vesicant model (MEVM). **Methods:** HaCaT cells were exposed to SM 30 min and then cultivated in SM free medium with or without S003 for up to 6 h. Cell viability, NAD⁺ and PAR contents, the percentage of apoptotic and necrotic cells, and caspase 3 activity were assayed. SM in propylene glycol (0.16 and 0.64 mg/ear) was applied to the right ear of KM mouse, S003 was ip at dose of 200 mg/kg. **Results:** The amount of intracellular PAR were decreased to 69% after exposed to 1000 mol/L SM about 30 min, and it was reduced to 11% by combined treatment with S003. NAD⁺ levels were diminished to 11% and cell viability were reduced to 45% after 6 h exposure of SM. While, treatment with S003 increased the NAD⁺ levels to 83% and cell viability to 78%. Meanwhile, the recuction of S003 on apoptosis and necrosis was also observed. While after 6 h exposure of 100 mol/L SM, S003 did not affect the cell viability, but significantly increased percentage of apoptotic and necrotic cells, and caspase 3 activation. In the MEVM, S003 significantly reduced relative ear weight about 30% in the 0.64 mg/ear SM injury, but showed no protective effect in the 0.16 mg/ear SM injury. **Conclusion:** S003 may have protective potentials in SM injury because of the protection of cell viability, the inhibition of PAR synthesis and intracellular NAD⁺ depletion, and recuction on apoptosis and necrosis. S003 can reduce the ear edema of SM exposure, but showed different actions under different levels of SM injury.

Keywords: Sulfur mustard; PARP; S003; apoptosis; ear edema

Acknowledgements: This work was supported by major scientific and technological specialized project for the significant formulation of new drugs (2009ZXJ09002-012, 2013ZX09J13103-01B)

S1.28**3-bromopyruvate induces necroptosis and autophagy in breast cancer cell lines**

Hao LIU¹, Yuan-yuan ZHANG¹, Qian-wen ZHANG¹, Zhen-hua CHAO¹, Pei ZHANG¹, Chen-chen JIANG², Zhi-wen JIANG¹*. ¹Faculty of pharmacy, Bengbu Medical College, Bengbu 233000, China; ²School of Medicine and Public Health, Faculty of Health, University of Newcastle, NSW, Australia

*To whom correspondence should be addressed.

Aim: 3-bromopyruvate (3-BrPA) has antineoplastic activity on malignant cells. In this study, we investigated the toxicity of 3-BrPA on two breast cancer cell lines.

Methods: The intracellular ATP level was measured by the ATP Assay Kit. Cell viability was measured using MTT assay. Cell death was assessed by flow cytometry with propidium iodide (PI) staining or PI/Annexin V staining. The ultrastructural details of two breast cancer cells treated with vehicle DMSO, 3-BrPA, or 3-BrPA plus caspase inhibitor (z-VAD-fmk) were analyzed by electron microscopy (EM). Small interfering RNA mediated RIP1 and RIP3 knockdown, and the expression of RIP1 and RIP3 were analyzed by Western blot. **Results:** 320 μ mol/L 3-BrPA had significant toxicity on MDA-MB-231 cells, whereas no obvious effect was observed in MDA-MB-435 cells. In MDA-MB-231 cells, z-VAD-fmk switches the apoptotic response to necroptosis; meanwhile, knockdown of RIP1 or RIP3 block this form of cell death. Morphologically, autophagosome in insensitive MDA-MB-435 cells were observed. MDA-MB-231 cells treated with 3-BrPA plus z-VAD-fmk exhibit the features of necrosis. **Conclusion:** 3-BrPA can induce necroptosis in MDA-MB-231 cells and the signaling pathway requires the involvement of RIP1 and RIP3. MDA-MB-435 cells which are insensitive to relatively low-level concentrations of 3-BrPA may be related to activating the autophagy pathway.

Keywords: 3-bromopyruvate (3-BrPA); breast cancer cells; necroptosis; autophagy

Acknowledgements: This work was supported by the National Natural Science Foundation of China (81000992, 81072207) and the Natural Science Foundation of Anhui Province (090413135), Education Department of Anhui Natural Science Research Key Project China (KJ2012A202).

S1.29**The tumor suppression pharmacodynamics research of new drug MNF-211 in tumor-burdened (PC-3) nude mice**

Xiang-yun LIU, Yue SHAO, Yi CHENG, Qi PAN, Jian-yan YAN, Lei LI, Li XU, Zu-yue SUN*. Department of Pharmacology and Toxicology, Shanghai Institute of Planned Parenthood Research, Shanghai 200032, China

*To whom correspondence should be addressed.

Aim: To observe the tumor suppression effects of MNF-211 on tumor-burdened (PC-3) nude mice by gastric gavage. **Methods:** We first establish prostate cancer model *in vivo* by subcutaneous injection of PC-3 cells into nude mice. Then, the

nude mice were randomly divided into 5 groups, including the negative control group, positive control group and MNF-211 high, medium and low dose group, each group of 11, and then respectively give solvent, bicalutamide 35 mg/kg and MNF-211 70, 35, and 17.5 mg/kg, by gastric gavage for 28 consecutive days. **Results:** Each dose group had different degrees of weight loss, but no significant difference among the groups. After D17 gastric gavage, relative tumor volume (RTV) of bicalutamide group and MNF-211 high, medium, low dose groups reduced obviously ($P < 0.05$). At d 28, compared with negative control group, bicalutamide group and MNF-211 high, medium dose groups were significant reduction ($P < 0.05$). The bicalutamide group and MNF-211 high, medium, low dose groups, all the relative tumor proliferation rate less than 60%, was 58.92%, 37.51%, and 53.75%, respectively ($P < 0.05$). Compared with negative control group, the proliferating cell nuclear antigen in bicalutamide group and MNF-211 high, medium, low dose groups decreased obviously ($P < 0.05$). **Conclusion:** New drug MNF-211 (35 and 70 mg/kg) and bicalutamide (35 mg/kg) has inhibitory effect on prostate cancer cells by subcutaneous inoculation into nude mice.

Keywords: prostate cancer; MNF-211; relative tumor proliferation rate; nude mouse; Proliferating cell nuclear antigen

Acknowledgements: This work was supported by New drug Major Program of China (No 2011ZX09301-005); National Natural Science Foundation of China (No 11101077); and Shanghai "Science and Technology Innovation Action Plan" experimental animal research project (No 11140901300, 11140901302).

S1.30

Downregulation of PLK1 inhibits tumor cell metastasis through degradation of β -catenin in esophageal squamous cell carcinoma

Xiao-ying LIU, Wen-jing YU, Li-mei CHEN, Zhi-qin GAO, Wei-guo FENG, Chang-qing DU, Chun-ling ZHAO*. *Department Of Cell Biology, Weifang Medical University, Weifang 261042, China*

*To whom correspondence should be addressed.

E-mail: zhaochunlingbj@163.com

Aim: PLK1 is a serine/threonine kinase that regulates cell cycle and is overexpressed in many tumors. Our study aimed to investigate the biological functions and mechanisms of PLK1 in metastasis of esophageal squamous cell carcinoma (ESCC). **Methods:** The ESCC cell overexpressed PLK1 was screened by RT-PCR and Western blot. The effective PLK1siRNA fragment was screened by Western blot. Effects of PLK1siRNA on migration and invasion of ESCC cells *in vitro* were studied by wound-healing assay and Matrigel chemoinvasion assay. The effective interference fragment was constructed into lentiviral vector, which infected ESCC cells. Interference efficiency was detected by RT-PCR and Western blot. The recombinant lentiviral vector on invasion and metastasis of ESCC cells *in vivo* were studied by pulmonary metastasis model in nude mice and metastasis model in embryonic zebrafish. Related molecules in Wnt/ β -catenin signaling transduction pathway which induce ESCC cells epithelial-mesenchymal transition (EMT) were detected by Western Blot and IP. **Results:** PLK1siRNA could obviously inhibit migration and invasion of ESCC cells *in vitro*. Lentivirus mediated RNA interference targeting PLK1 could significantly inhibit metastasis of ESCC cells *in vivo*. PLK1 inhibited expression of E-cadherin by upregulating Snai, thereby induced ESCC cells EMT. In ESCC cells overexpressed PLK1, β -catenin from the cell and the nucleus were upregulated. In the polymerized compound that degraded β -catenin, the expression of APC and Axin decreased in immune precipitate of GSK-3, and the expression of APC and GSK-3 β also significantly decreased in immune precipitate of Axin. **Conclusion:** RNA interference targeting PLK1 could obviously inhibit invasion and metastasis of ESCC cells. PLK1 induced EMT to promote the metastasis of ESCC by regulating stability of β -catenin.

Keywords: PLK; Esophageal squamous cell carcinoma; RNA interference; EMT; β -catenin

Acknowledgements: This work was supported by the National Natural Science Foundation of China (30901779).

S1.31

Paeoniflorin inhibited the apoptosis in human hepatocellular carcinoma cells via suppressing hedgehog/Gli signaling pathway

Jing-tao LU, Wei HE, Sha-sha SONG, Ping-fan YUAN, Wei WEI*. *Institute of Clinical Pharmacology, Anhui Medical University, Key Laboratory of Anti-inflammatory and Immune Medicine (Anhui Medical University), Ministry of Education, Hefei 230032, China*

*To whom correspondence should be addressed.

E-mail: wwei@ahmu.edu.cn

Aim: To investigate the effect of paeoniflorin (Pae) on the apoptosis in human hepatocellular carcinoma (HCC) cells and its mechanism. **Methods:** Cells apoptosis was determined by flow cytometric analysis. The protein expression of B-cell CLL/lymphoma 2 (Bcl-2), p53, caspase-3 and shh, Gli-1 of hedgehog/Gli signaling pathway in HepG₂ and Bel-7402 was examined by Western blot analysis. **Results:** Pae (12.5, 25, and 50 μ mol/L) significantly induced earlier period apoptosis of HCC HepG₂ and Bel-7402 cells ($P < 0.01$); Pae (100 μ mol/L) simultaneously led to later period apoptosis of HepG₂ and Bel-7402 cells; Pae (12.5, 25, and 50 μ mol/L) significantly decreased the expression of shh and Gli-1 ($P < 0.01$), while significantly decreased the expression of anti-apoptotic Bcl-2, increased the expression of caspase-3 and p53 ($P < 0.01$). **Conclusion:** Pae can induce apoptosis of HCC cells, which maybe correlated with down-regulating hedgehog/Gli signaling pathway by Pae.

Keywords: paeoniflorin; human hepatocellular carcinoma; apoptosis; hedgehog/Gli signaling pathway

S1.32

The apoptosis of human breast cancer MDR cell line MCF-7/Adr induced by novel compound GA and realted mechanism

Wei LU¹, Jin-bo CHEN^{1,2}, Ya-juan CHEN¹, Hong-bin ZHANG^{2,*}, Chen QING^{1,*}. ¹The School of Pharmaceutical Science & Yunnan Key Laboratory of Pharmacology for Natural Products, Kunming Medical University, kunming, Yunnan, 650500, China; ²Key Laboratory of Medicinal Chemistry for Natural Resource, Ministry of Education, School of Chemical Science and Technology, Yunnan University, Kunming 650091, China

*To whom correspondence should be addressed.

E-mail: qingchenhh@yeah.net

Aim: To investigate the apoptosis of human tumor multidrug resistance (MDR) cell induced by a novel tetracyclic diterpenoid compound GA13315 and realted mechanism. **Methods:** This study was under taken using human breast cancer cell line MCF-7/Adr with typical multi-drug resistant (MDR) properties induced by doxorubicin (ADM) as model. Cells apoptosis induced by compound GA alone or in combination with ADM were measured by flow cytometry (FCM) combined Annexin V/PI staining. Effects of compound GA alone or in combination with ADM on transcription of MDR related genes: multidrug resistance-related protein (MRP), lung resistance protein (LRP), mutant p53, bax, bcl-2, and caspase-3 were detected by real-time fluorescence quantitative PCR (RQ-PCR). The proteins expression of MDR related genes MRP, LRP, mutant p53, bax, bcl-2, and caspase-3 were measured by Western blot. **Results:** Apoptosis of MCF-7/Adr was significantly induced by compound GA alone or in combination with ADM in a dose-dependent and time-dependent manner. After exposure of compound GA alone at concentration of 8, 16, 32, and 64 μ mol/L to MCF-7/Adr for 24 h and 48 h, the apoptosis rate of cells were 8.92% \pm 2.40%, 30.96% \pm 1.37%, 41.89% \pm 6.97%, 87.66% \pm 3.74% and 18.35% \pm 0.97%, 23.51% \pm 4.24%, 63.92% \pm 3.88%, 78.16% \pm 2.25%, respectively. The treatment of the above same concentrations of GA combined separately with concentration of 18.4 μ mol/L ADM exposure to MCF-7/Adr for 24 h and 48 h, the apoptosis rate of cells were 21.05% \pm 5.74%, 26.45% \pm 6.77%, 35.69% \pm 6.34%, 53.98% \pm 4.77%, and 23.60% \pm 7.44%, 35.27% \pm 4.58%, 54.28% \pm 6.06%, 91.62% \pm 4.12%, respectively. RQ-PCR results showed that after exposure of compound GA alone or in combination with ADM to MCF-7/Adr, the expression of mRNA of Bcl-2, mutant p53 were down-regulated; the expression of mRNA of bax and caspase-3 were up-regulated, but effects on the mRNA expression of MRP and LRP were unclear. Western blot results showed that GA alone and in combination with ADM could down-regulate the protein expression of Bcl-2, mutant p53 and LRP, and up-regulated the protein expression of bax at the same time, but treatment of GA alone and in conjunction with ADM had no obvious effect on the protein expression of MRP of MCF-7/Adr. **Conclusion:** Either treatment of MCF-7/Adr with GA alone or GA in combined with ADM, can significantly induced apoptosis of MCF-7/Adr cells. But GA does not always enhance the effect of ADM on cell apoptosis, the reason is not unclear. The mechanism of apoptosis of MCF-7/Adr induced by GA alone or GA in combined with ADM is involved in down-regulation of the expression of bcl-2, mutant p53 and LRP, up-regulation of the expression of bax. But whether related to the effect on the expression of MRP and caspase-3 remain still uncertain.

Keywords: compound GA; tumor; multi-drug resistance; apoptosis

Acknowledgements: This work was supported by grants from the National Natural Science Foundation of China (81160405 and 21062029).

S1.33

Pulsatilla chinensis (Bunge) Regel saponins suppresses the proliferation of tumor

cells and induces apoptosis and autophagy via inhibiting PI3k/Akt signalling pathway

Ying-ying LUO^{1,2}, Lan-ying CHEN^{1,*}, Ya-ru CUI¹, Xin-xu XIE¹, Mu-lan WANG¹, Qin GONG¹, Jian-sheng GUO^{2,*}. ¹Jiangxi university of TCM, Nangchang 330006, China; ²Hunan university of Chinese Medicine, Hunan 410208, China

*To whom correspondence should be addressed.

Aims: To investigate the *in vitro* and *in vivo* anti-tumor effect of Pulsatilla saponins, the major components extracted from *Pulsatilla chinensis* (Bunge) Regel, and its underlying mechanisms. **Methods:** Human tumor cells were treated with Pulsatilla saponins, cell viability was assessed by MTT assay; cell cycle and apoptosis were assessed by flow cytometry; cleaved-caspase3, LC3, PI3k, p-PI3k, Akt, and p-Akt were measured by Western Blot assay. Tumor xenograft model was used to further evaluate the anti-tumor effect. The T/C(%) value and tumor inhibition rate were calculated. **Results:** Pulsatilla saponins could inhibit the proliferation of tumor cells, it induced tumor cells S/G₁ phase arresting and apoptosis by increasing cleaved caspase3. It also induced the expression of LC3-II, which is a marker for cellular autophagy. Further study showed Pulsatilla saponins decreased the expression of p-PI3k and p-Akt. Pulsatilla saponins presented anti-tumor effect with 56.74% of tumor inhibition rate in A₅₄₉ xenograft model, without reducing leukocyte number. **Conclusion:** Pulsatilla saponins induced tumor cells apoptosis and autophagy via inhibiting PI3k/Akt pathway, it inhibited tumor growth without reducing leukocyte number, may be a novel effective candidate as chemotherapeutic agent.

Keywords: Pulsatilla saponins; anti-tumor; apoptosis; autophagy; PI3k/Akt pathway

Acknowledgements: This research is supported by projects from National Science and Technology Major Projects for "Major New Drugs Innovation and Development" (No 2011ZX11102-001-19)

S1.34

Progress in mechanism of apoptosis induced by Chinese herbs and the relationship between autophagy in breast cancer cell

Cong MA^{1,2}, Xiao-min WANG^{1,*}, Ming-wei YU¹, Gan-lin ZHANG¹. ¹Department of Oncology, Affiliated Beijing Hospital of Traditional Chinese Medicine, Capital Medical University, Beijing 100010, China; ²Beijing University of Chinese Medical, Beijing 100029, China

*To whom correspondence should be addressed.

E-mail: ntxm100@sina.com

Defects in programmed cell death (PCD) regulation contribute to tumor. Apoptosis as a PCD, is a barrier to cancer development. The autophagic program could degrade some cellular organelle to maintain homeostasis. Metabolites support survival in the stressed environments. Induction of autophagy also may operate independently of or in concert with apoptosis. Nutrient starvation and certain cytotoxic drugs can induce elevated levels of autophagy that are apparently protective for cancer cells. Autophagy seems to have conflicting effects on tumor cells. Recent studies suggested that many Chinese herbs are potential anti-tumor drugs. Herbs extract have multi anticancer effects, some could active apoptotic cell death and induce autophagy. There is increasing interest regarding interconnections between apoptotic and autophagic. Although there has been some progress in studying the apoptotic mechanism of herbs induction and the relationship of autophagy, little is known about the role autophagy plays in breast cancer cell. In this review, we stated the mechanism of apoptosis induced by Chinese herbs and the relationship between autophagy in breast cancer cell.

Keywords: herb; breast cancer; apoptosis; autophagy

Acknowledgements: This work was supported by National Natural Science Foundation 81173239 and 81202689), Beijing Natural Science Foundation (7132104), and Basic-Clinic Cooperation Fund of Capital Medical University (11JL62)

S1.35

Study of the antineoplastic role and its immune mechanism of Xihuang Pill Ethanol extract

Jie MA¹, Wei YANG², Shuo GUAN², Jun-xia HU², Chang-qian ZENG⁴, Wen-bin GAO³, Weng-bo LIANG^{4,*}. ¹Department of pharmacology, Zunyi Medical College, Zunyi 563000, China; ²Department of Oncology, Medical college of Dalian university, Dalian 116622, China; ³Department of Oncology, Affiliated Zhongshan Hospital of Dalian University, Dalian 116000, China; ⁴Department of pharmacology, Medical college of Dalian University, Dalian 116622, China

*To whom correspondence should be addressed.

Aims: To investigate the antitumor activity and immunological mechanisms of Xihuang pill Ethanol extract on tumor-bearing rats. **Methods:** Walker256 tumor-

bearing rats were randomly divided into 7 groups ($n=10$): normal control group, model control group, Xihuang pill Ethanol extract low dose group, midst dose group, high dose group, and lentinan group and Xihuang pill high dose group. After the drug administration in 14 consecutive days, the inhibitory rate of tumor was observed. FCM method was used to detect the number of CD4⁺, CD8⁺, B7-1. ELISA was used to detect the level of IL-2, IL-6, IL-10, TGF- β , IFN- γ in serum. The comparison was made among groups. **Results:** The Xihuang pill Ethanol extract can effectively inhibit the growth of W256. The high dose group inhibitory rate of tumor was 36.8%. Compare with model group the serum level of IL-2, IFN- γ and the number of B7-1 CD4⁺ CD8⁺ of the Xihuang pill Ethanol extract group significantly up-regulated, however the number of IL-6, IL-10, TGF- β , significantly decreased ($P<0.05$). The Xihuang pill ethanol extract indicator was better than Xihuang pill in equivalents dose. **Conclusion:** Xihuang pill ethanol extract improve the Xihuang pill bioavailability, and Xihuang pill Ethanol extract has obvious antitumor activity. The mechanism is relate to the enhancement of immune function and reducing the level of immunosuppressive factors.

Keywords: Xihuang pill; ethanol extract; Walker 256 cell of rats mammary cancer; rats; immunological mechanism

S1.36

Bio-activity of novel Histone Lysine specific demethylase1inhibitorin human gastric cancer cell line MGC-803

Jin-lian MA, Yi-chao ZHENG, Wen ZHAO, Wen-Lei LV, Ang YANG, Zhi-Ru WANG, Jin-Feng LI, Hong-min LIU*. School of Pharmaceutical Sciences, New Drug Research & Development Center, Zhengzhou University, Zhengzhou 450001, China

*To whom correspondence should be addressed.

Aims: Histone epigenetic modifications including acetylation, methylation, phosphorylation and ubiquitination play important roles in structural changes of chromatin. Lysine specific demethylase1(LSD1) belongs to the flavin adenine dinucleotide (FAD)-dependent amine oxidase family which is a significant enzyme for post-translation modification and regulates the activation and silencing of gene transcription. Studies have shown that LSD1is highly expressed in a variety cancers. Moreover, it can suppress translation of tumor suppressor gene. In this study, modulation for activity of LSD1by its inhibitor 3e in MGC-803 was investigated. **Methods:** The novel LSD1 inhibitor 3e was screened and obtained by LSD1 recombinant based screen model. The bioactivity of 3e in cell level was investigated in human gastric cancer cell line MGC-803. After 48 h treatment of compound 3e, total lysates and total histone were extracted, denatured, and separated in SDS-PAGE. Then the results were analysed by Western-blot. In this study, LSD1, H3K4me1, H3K4me2, H3K4me3, and H3K9me2 were analysed by Western blot. LSD1 and snail were analysed by Crosslink Co-Immunoprecipitation. **Results:** The small molecule inhibitor specifically inhibites activity of LSD1 but does not influence expression LSD1 in MGC-803 cells. Meanwhile, it can interrupt the complex formation between LSD1 and snail, which results in the up regulation of E-cadherin. **Conclusion:** As expected, Crosslink Co-Immunoprecipitation of LSD1 revealed the association with Snail, but the interaction of Snail with LSD1 can be blocked by LSD1 inhibitor 3e. It might regulated the processes of epithelial-mesenchymal transition (EMT), and further research is still going on.

S1.37

ATM-mediated Chk1/2 and Cdc2 phosphorylation is involved in Jaridinin-induced cell cycle arrest in human esophageal cancer cells

Yong-cheng MA¹, Xiao-jing SHI¹, Yu KE¹, Xiao-lin ZI², Mi-yang DAI¹, Wen ZHAO¹, Hong-min LIU^{1,*}. ¹School of Pharmaceutical Sciences, Zhengzhou University, Zhengzhou 450001, China; ²Departments of Urology, Pharmacology and Pharmaceutical Sciences, University of California, Irvine, CA 92697, USA

*To whom correspondence should be addressed.

Aims: Jaridinin, a novel diterpenoid from *Rabdosia rubescens* has been shown previously to inhibit proliferation of esophageal squamous cancer cells (ESCC) through G₂/M phase cell cycle arrest. However, the involved mechanism is not fully understood. **Methods:** Cell cycle was analysed by flow cytometry. The related protein expressions were determined by Western blot. **Results:** In this study, we found that the cell cycle arrest by Jaridinin was associated with the increased expression of ATM and phosphorylation of Cdc2 at Tyr15. Jaridinin also resulted in enhanced phosphorylation of Cdc25C via activation of checkpoint kinases Chk1 and Chk2, as well as in increased phospho-H2A.X (Ser139). We also found that the increased activation of ATM is related to the DNA damage caused by Jaridinin. Furthermore, Jaridinin-mediated alterations in cell cycle arrest were significantly attenuated in the presence of NAC, implicating the involvement of ROS in

Jaridonin's effects. On the other hand, addition of ATM inhibitor caffeine reversed Jaridonin-related activation of ATM and Chk1/2 as well as phosphorylation of Cdc25C, Cdc2 and H2A.X and G₂/M phase arrest. **Conclusion:** These findings identified that Jaridonin-induced cell cycle arrest in human esophageal cancer cells is associated with ROS-mediated activation of ATM-Chk1/2-Cdc25C pathway.

Keywords: Jaridonin; diterpenoid; *Rabdosia rubescens*; ROS; ATM; cell cycle arrest; Chk1/2; Cdc2; esophageal cancer cells

Acknowledgements: This work was supported by National Natural Science Foundation of China (No 81172937 to Dr Hong-min LIU and No 81270270 to Dr Wen ZHAO).

S1.38

Progress of studies on anti-tumor action of *Caulis Spatholobi*

Nan NAN, Xiao-min WANG*, Gan-lin ZHANG, Ming-wei YU. *Beijing Hospital of Traditional Chinese Medicine, Capital Medical University, Beijing 100010, China*

*To whom correspondence should be addressed.

Caulis Spatholobi is a kind of herb with invigorating the circulation of blood and replenishing blood, which contains flavone, condensed tannin, sterides and so on. According to modern pharmacology, *Caulis Spatholobi* has anti-coagulants, improves hematopoiesis, antiviral and anti-tumor activities by improving cell death and inhibiting cell proliferation. This paper has reviewed anti-tumor components in *Caulis Spatholobi* and their pharmacological mechanism.

Keywords: *Caulis Spatholobi*; anti-tumor; herb

Acknowledgements: This work was supported by Beijing Natural Science Foundation (7122083).

S1.39

Gamma-linolenic acid trigger apoptosis in glioblastoma cells by regulating miR-22 and its corresponding gene Sirt-1 expression

Yi-ran NI¹, Fan-fan CHEN². ¹Department of Pathology, Medicine school, Three Gorges University, Yi Chang 443002, China; ²Medicine school, Three Gorges University, Yi Chang 443002, China

Aim: Gamma-linolenic acid (GLA) was reported had cytotoxic effect on glioblastoma cells. In this study, we investigated the direct cytotoxic action of GLA against glioblastoma cells and the underlying mechanisms. **Methods:** Glioblastoma cells (GBM2) were treated with 25, 75, and 150 μmol/L GLA for 24 h and the proliferation activity of cells were detected by MTT assay. The apoptosis effect and the change of cell cycle were determined by flow cytometry (FCM) technology. The Expression of microRNA-22 and mRNA levels of its corresponding tumor suppressor gene sirt-1 were analyzed by using QRT-PCR. **Results:** The apoptosis of glioblastoma cells was markedly enhanced when the concentration of GLA increased ($P < 0.05$ between each group), and there was an inverse correlation between the concentration of GLA used and miR-22 expression tested by QRT-PCR. We also found an increased expression of Sirt-1 in the same treated cells. **Conclusion:** We conclude that GLA can trigger apoptosis in glioblastoma cells. The mechanisms might be in part due to regulating miR-22 and its corresponding tumor suppressor gene Sirt-1 expression. Our study highlights the potential application of GLA on the anti-cancer therapy of glioblastoma.

Keywords: gamma-linolenic acid; apoptosis; glioblastoma; miRNA-22; Sirt-1

S1.40

Anti-tumor effect of annexinA2 monoclonal antibody *in vitro*

Ji-qin NIE, Jian-lin YANG, Wei ZHANG, Chao-qi LIU*. *Institute of Molecular Biology of Three Gorges University, Yichang 443002, China*

*To whom correspondence should be addressed.

Aim: Annexins are a structurally related family of calcium and phospholipid-binding proteins that are involved in the regulation of a range of molecular and cellular processes. To explore the anti-tumor effect of annexinA2 (ANXA2) and its related mechanism, which provide the experimental basis for the tumor treatment *in vivo*. **Methods:** pET28a(+)-ANXA2 prokaryotic expressing vector was constructed and the expressed protein was purified, and then it was identified by SDS-PAGE and Western blot with His-tag antibody. Balb/c mice were immunized with the recombinant protein. Through cell fusion and selective culture, stable hybridoma cell strains were obtained. The antibody specificity and sensitivity were detected by indirect ELISA, Western Blot and immunoprecipitation. After ANXA2 antibody treated HepG2 cells, MTT detected the cell proliferation, FCM detected cell cycle, RT-PCR and real-time PCR detected cell cycle-related genes; Western blot detected cell cycle-related protein CDK6. Migration of HepG2 cells were demonstrated by Transwell method. **Results:** pET28a(+)-ANXA2 prokaryotic expressing vector

was successfully constructed, and ANXA2 protein obtained and identified. Four hybridoma cell strains which secreted ANXA2 monoclonal antibody were obtained. A monoclonal antibody specifically recognizes the PC3 (human prostate cancer), ECV (human vascular endothelial cells), T24 (human bladder cancer), HepG2 (human hepatoma cells), PANC1 (human pancreatic cancer cells) and cancer tissue expressed ANXA2 protein was identified by Western blot. MTT showed that the ANXA2 monoclonal antibody could inhibit the growth of HepG2 cells. FCM displayed that the antibody inhibited the growth of HepG2 cells by extended G₁ phase, and shortened G₂+M phase time. RT-PCR results identified that the antibody inhibited cell cycle related genes hCCD1, hE2F1, CDK6 expression. Western blot results showed that the antibody decreased HepG2 cells cycle protein CDK6 expression. Transwell method results confirmed that the antibody treatment inhibited HepG2 cells migration. **Conclusion:** In this study, we obtained ANXA2 prokaryotic expression protein and high titers of specific monoclonal antibodies. The ANXA2 monoclonal antibody inhibited tumor cells growth and migration. These results hinted that the hinte expression has certain correlation with tumor occurrence and development and its antibodies may be used for tumor diagnosis and therapy.

Keywords: annexinA2; monoclonal antibody; cell cycle cell migration

S1.41

The novel tubulin-targeting agent MT-4 suppresses human ovarian cancer growth *in vitro* and *in vivo*

Hui-Chen PAI¹, Shioh-Lin PAN², Che-Ming TENG¹. ¹Pharmacological Institute, College of Medicine, National Taiwan University, No. 1, Jen-Ai Road, Sec. 1, Taipei, Taiwan, China; ²Department of Pharmacology, Taipei Medical University, Taipei, 11031, Taiwan, China

Aims: In this study, the anti-cancer mechanisms of moscatilin(MT-4), have been examined in cisplatin-sensitive ovarian cancer cells (A2780) and multidrug resistance ovarian cancer cells (NCI-ADR-RES). **Methods:** MTT, flow cytometry, tubulin binding assay, and Western blot assay were used. **Results:** MT-4 inhibited cell viability in both ovarian cancer cells. Drug resistance ratios were also calculated as the ratio of the IC₅₀ in the resistant cell line (NCI-ADR-RES cells) to the IC₅₀ in the sensitive cell line (A2780 cells). The data demonstrated that, compared with A2780 cells, the NCI-ADR-RES cells displayed a 271.3-fold relative resistance to paclitaxel. In contrast, the A2780 cells and NCI-ADR-RES cells did not display any difference in their response to MT-4 (relative resistance ratio=3.7). We also demonstrated that MT-4 caused cell population at G₂/M phase arrest and then increased sub G₁ phase in both ovarian cancer cells. The data showed that MT-4 could disrupt microtubule dynamics and induce phosphorylation of G₂/M protein. Treatment of MT-4 induced apoptosis through catalytic cleavage of caspase-3, -7, -8, -9, and poly (ADP-ribose) polymerase (PARP). In addition, we also found that inhibition of p38 MAPK pathway and could reverse MT-4-induced apoptosis. We demonstrated that MT-4 (20 mg/kg, qd, iv) significantly inhibited tumor growth in xenograft mouse model. **Conclusion:** The MT-4 inhibited cell viability *in vitro* and *in vivo*. These finding indicated that MT-4 can be a potential anticancer agent for treatment of cancer.

Keywords: moscatilin (MT-4); resisrance; apoptosis; ovarian cancer cells; anti-cancer

S1.42

The inhibitory effect of celecoxib on Lewis lung carcinoma cell line by regulation of cyclooxygenase-2/cytosolic phospholipase A2

Yue PAN, Xuan-ang XIAO, Li-jing LI*. *Department of Pharmacology, Norman Bethune Medical College, Jilin University, Changchun 130021, China*

*To whom correspondence should be addressed.

Aim: Celecoxib is a potent nonsteroid anti-inflammatory drug (NSAID) that has demonstrated great promise in cancer chemoprevention and treatment. The goal of this study was to determine the inhibitory effect and mechanism of celecoxib on Lewis lung carcinoma. **Methods:** The effect of celecoxib on viability of Lewis lung carcinoma cells was assessed with methyl thiazolyl tetrazolium (MTT) assay. Apoptosis and the mitochondrial membrane potential were detected by flow cytometric assay. The protein expression of cytosolic phospholipase A2 (cPLA2), cyclooxygenase-2 (COX-2), and peroxisome proliferator-activated receptor gamma (PPARc) were determined by Western blot analysis. **Results:** Celecoxib inhibited the proliferation of Lewis lung carcinoma and induced apoptosis in a dose-dependent manner by breakdown of mitochondrial membrane potential. The protein expressions of cPLA2 and PPARc were upregulated, but COX-2 protein expression was downregulated in the Lewis lung carcinoma cells exposed to celecoxib. **Conclusion:** The major findings in this study are that celecoxib could inhibit the viability of Lewis lung carcinoma cells by interference of the arachidonic

acid pathway and upregulation of PPARc simultaneously, which are novel and important since the molecular mechanisms of celecoxib underlying the anti-neoplastic effects remain unclear.

Keywords: celecoxib; cyclooxygenase-2; arachidonic acid; prostaglandin E₂

S1.43

Anticancer activity of melanoxoin in human non-small cell lung cancer cells and in nude mice

Chiehyu PENG^{1,2}, Shoufang WU³, Hsiuyi HUNG¹, Yangchang WU^{1,2}. ¹Natural Medicinal Products Research Center and Center for Molecular Medicine, China Medical University Hospital, Taichung 404, Taiwan, China; ²School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung 404, Taiwan, China; ³Graduate Institute of Natural Products, Kaohsiung Medical University, Kaohsiung 807, Taiwan, China

Aim: There are various anticancer drugs, which derived from natural products. In this study, we screened a serious of compounds isolated from *Pterocarpus santalinus* to identify their cell cytotoxicity and antitumor mechanisms. **Methods:** We performed the MTT assay to evaluate the cytotoxic effects of test compounds against several human cancer cell lines. We used Human OneArray® v5 to comprehensive analysis the DNA transcription of H1299 cells. Cell cycle distributions were analysed by flow cytometry. We performed the *in vitro* tubulin polymerization assay to examine whether the test compound affects microtubule organization. We used comet assay to measuring DNA strand breaks. We examined *in vivo* antitumor activity of melanoxoin in nude mice bearing human lung tumor xenografts. **Results:** Among the serious of compounds isolated from *Pterocarpus santalinus*, melanoxoin showed the highest cytotoxic effects with a IC₅₀ of 1.98 µg/mL in the human non-small cell lung cancer H1299 cells. It showed that melanoxoin regulated the transcription of several cell cycle regulators in the H1299 cells. Melanoxoin inductions of tubulin depolymerization suggest that melanoxoin causes G₂/M arrest and then increased tumor cell apoptosis. The neutral comet assay revealed that the treatment with melanoxoin induced significant DNA damage in H1299 cells. Our mice xenograft models show the *in vivo* efficacy of melanoxoin. **Conclusion:** We are interested in the development the melanoxoin as a new source of anti-cancer drug.

Keywords: *Pterocarpus santalinus*; melanoxoin; apoptosis

S1.44

Compound *Astragalus* and *Salvia Miltiorrhiza* extract retards hepatocellular carcinoma progression by inhibiting fibrosis and PAI-1 transcription activity

Wen-juan RUI, Lei XIE, Shu-fang HE, Chao WU, Yan YANG*. Department of Pharmacology and Institute of Natural Medicine, Anhui Medical University, Hefei 230032, China

*To whom correspondence should be addressed.

E-mail: yangyan276866@sohu.com

Aims: To investigate the effects of Compound *Astragalus* and *Salvia Miltiorrhiza* Extract (CASE) on the hepatocarcinogenesis in rats. **Methods:** Rats were randomly divided into control group, DEN (0.2%, ig ×14 weeks) group, and three dosages groups of CASE (60, 120, or 240 mg/kg, ig ×16 weeks). The rats were sacrificed at 12th week or 16th week after the start of DEN administration. Blood samples were collected for determination liver function. The extent of fibrosis of liver was determined by histopathologic examination. The expression of α-smooth muscle actin (α-SMA) and Glutathione S-transferase protein (GST-P) were detected by immunohistochemistry. PAI-1 transcription activity was analyzed by Real-time RT-PCR. **Results:** At 12th week, CASE significantly ameliorated the serum parameters of liver function, and decreased the extent of fibrosis and the overexpression of α-SMA. At 16th week, the elevated incidence and multiplicity of hepatoma, and GST-P expression were reduced by CASE treatment. CASE inhibited PAI-1 transcriptional activity in HepG2 cell induced by TGF-β₁ in a dose-dependent manner. **Conclusion:** These results suggested that CASE might retard DEN-induced hepatocarcinogenesis by inhibiting fibrosis and PAI-1 transcription activity.

Keywords: CASE; diethylnitrosamine; hepatic fibrosis; hepatocellular carcinoma; plasminogen activator inhibitor

Acknowledgements: This study was financially supported by the National Natural Science Foundation of China (No 81073098) and the Academic and Technique Foregoer Foundation of Anhui province, China (2012381).

S1.45

Suppression of signal transducer and activator of transcription-3 signaling cascade by zerumbone inhibits the proliferation and induces apoptosis in renal cell carcinoma xenograft mouse model

Muthu K SHANMUGAM¹, Rajendran PERAMAIYAN¹, Feng LI¹, Siveen KODAPPULLY SIVARAMAN¹, Kwang Seok AHN², Alan Prem KUMAR^{1,3,4,5} and Gautam SETHI^{1,3,*}.

¹Department of Pharmacology, Yong Loo Lin School of Medicine, National University of Singapore, 117597 Singapore; ²College of Oriental Medicine, Kyung Hee University, Seoul 130-701, Republic of Korea; ³Cancer Science Institute of Singapore, Centre for Translational Medicine, 14 Medical Drive, #11-01M, 117599 Singapore; ⁴School of Biomedical Sciences, Faculty of Health Sciences, Curtin University, Western Australia; ⁵Department of Biological Sciences, University of North Texas, Denton, Texas, USA.

*To whom correspondence should be addressed.

Aim: Signal transducer and activators of transcription 3 (STAT3) has been linked to the deregulated proliferation, survival, and angiogenesis in renal cell carcinoma (RCC). In this study we investigated whether zerumbone, a sesquiterpene, obtained from the plant *Zingiber zerumbet* Smith exerts its anticancer effect through the modulation of STAT3 activation pathway. **Methods:** The pharmacological effects of zerumbone on STAT3 activation, associated protein kinases and phosphatase, STAT3-regulated gene products and apoptosis was investigated using both RCC cell lines and xenograft mouse model. **Results:** We observed that zerumbone suppressed STAT3 activation in a dose- and time-dependent manner in RCC cells and was mediated through the inhibition of activation of upstream kinases c-Src, JAK1, and JAK2 and induced apoptosis in RCC cells. We found that zerumbone induced the expression of tyrosine phosphatase SHP-1 that correlated with the down-regulation of constitutive STAT3 activation. Deletion of SHP-1 gene by siRNA abolished the ability of zerumbone to inhibit STAT3 activation. The inhibition of STAT3 activation by zerumbone also caused the suppression of the gene products involved in proliferation, survival, and angiogenesis. Finally, when administered ip, zerumbone inhibited STAT3 activation in tumor tissues and the growth of human RCC xenograft tumors in athymic nu/nu mice. **Conclusion:** Overall, our results suggest that zerumbone is a novel blocker of STAT3 signaling cascade with potential for the treatment of RCC and other solid tumors.

Keywords: STAT3; JAK2; zerumbone; renal cell carcinoma; apoptosis.

S1.46

The anti-tumor activities of OP46, a novel ent-kaurene diterpenoid compound from *Isodonrubescens*, and its involving molecular mechanisms

Xiao-jing SHI, Yong-cheng MA, Dong CHEN, Ran WANG, Xu-yao ZHANG, Wen ZHAO, Hong-min LIU*. School of Pharmaceutical Sciences, Zhengzhou University, Zhengzhou 450001, China

*To whom correspondence should be addressed.

Aim: This current study is designed to determine the cytotoxic effects of OP46, a new ent-kaurene diterpenoid isolated from *Isodonrubescens* on human cancer cells and the underlying mechanisms. **Methods:** MTT assay and Trypan Blue exclusion method were used to evaluate the cell viability. Cell morphology, apoptosis, autophagy and cell cycle distribution were observed or analyzed by fluorescence microscopy or flow cytometry. **Results:** Compared with controls, OP46 caused strong anti-proliferative in human cancer cell lines MCF-7, PC-3, MGC-803, EC109 cells. OP46 also resulted in significant apoptotic, autophagy and G₂/M phase cell cycle arrest in human liver cancer cell SMMC-7721, but not in normal human liver cell L-02. These cytotoxic effects of OP46 on SMMC-7721 were associated with marked accumulation of reactive oxygen species (ROS), whereas a ROS scavenger, N-acetyl-L-cysteine (L-NAC), significantly attenuated these effects. **Conclusion:** OP46 could selectively inhibit the growth of cancer cells in vitro. The GSH depletion by OP46, associated with the generation of ROS, maybe a central mechanism of OP46-induced cytotoxicity in cancer cells.

Keywords: OP46; ent-kaurene diterpenoid; anticancer activity; GSH; ROS

Acknowledgements: This work was supported by National Natural Science Foundation of China (No 81172937 to Dr Hong-min LIU and No 81270270 to Dr Wen ZHAO).

S1.47

Effects of *Astragalus polysaccharide* on enteric mucosal immunity in S180 tumor bearing mice

Shu-yu SUN¹, Xiao-juan HE², Chang-yuan YU¹, Ai-ping LU². ¹College of Life Science and Technology, Beijing University of Chemical Technology, Beijing 100029, China; ²Institute of Basic Research in Clinical Medicine, China Academy of Chinese Medical Sciences, Beijing 100700, China

Aim: *Astragalus polysaccharide* (APS) is a main immunocompetent constituent of *Radix Astragali*. In this study, the effects of APS on enteric mucosal immunity in S180 tumor bearing mice were investigated. **Methods:** Mice were subcutaneously implanted with S180 tumor cells in the right axillary region. Then these mice

were randomly divided into model group, diamminedichloroplatinum(DDP) group, APS low dose group, and APS high dose group. Another ten normal mice were used as normal control. The therapeutic effects of APS were determined by examining tumor size. The production of IgA in enteric mucosa was detected by Immunohistochemistry (IHC) assay. The number of Peyer's pathes(PPs) was detected by counting assay. **Results:** After 14 days of orally administration, APS could inhibit S180 tumor growth *in vivo*. Compared with model group, the number of PPs increased in APS group. In addition, production of IgA in the enteric mucosa was also improved in APS group, there was a significant difference compared with model group. **Conclusion:** The results indicated that APS might improve enteric mucosal immunity by regulating concentrations of IgA and PPs in S180 tumor bearing mice.

Keywords: *Astragalus polysaccharide*; S180 tumor; IgA; enteric mucosal immunity

Acknowledgements: This research was supported in part by the projects from National Natural Science Foundation of China (No 81001676).

S1.48

Research advances in Bacillus Calmette-Guerin anticancer action

Han-wen TIAN¹, Chen-yi YE^{2,*}. ¹Department of Pharmacology, West China School of Preclinical and Forensic Medicine, Sichuan University, 3-17 Renmin South Road, Chengdu, Sichuan 610041, China; ²Clinical Medicine, West China School of Medicine, SiChuan University, Chengdu 610041, China

*To whom correspondence should be addressed.

E-mail: yechenyi577@163.com

Bacillus Calmette-Guerin (BCG) is the only available vaccine against tuberculosis, it is an attenuated live vaccine prepared from *Mycobacterium bovis*. Now researches show that BCG has antineoplastic activity. This paper reviews applications and mechanisms of BCG anticancer action. BCG can be used to treat malignant tumors, including pediatric leukaemia, melanoma, cancer of colon, hepatoma, pulmonary carcinoma, renal cancer, and bladder cancer, by exclusive use of BCG or combined therapy to produce local antitumor action and general anticancer action. The mechanisms of BCG underlying antitumor therapy are involved in the receptor responsible for the attachment of BCG to fibronectin, directly killing tumor cells, enhancement of antigen-presenting dendritic cells ability, cytotoxic T cells activated by CD4⁺ and CD8⁺, enhancing the antitumor activity of NK cells.

Keywords: BCG; cancer; anticancer; antitumortherapy

S1.49

Corilagin nanoparticle-induced apoptosis in human gastric cancer SGC-7901 cells via the mitochondrial pathway

Bao-qing WANG. College of Pharmacy, Harbin University of Commerce, Harbin 150076, China

Aim: Corilagin is a novel member of polyphenol family which has been discovered from the medicinal plant *Geranium* in northeast China. Corilagin nanoparticle is a new physical dosage-form. This study was designed to investigate the effect of corilagin nanoparticle on the apoptosis of SGC-7901 cells. **Methods:** The cytotoxicity was assessed with SRB assay. The morphological changes were observed by transmission electron microscopy (TEM). The levels of mitochondrial transmembrane potential and ROS were analyzed by Laser confocal scanning microscopy (LCSM). The apoptotic rate was detected by flow cytometry (FCM). The expression of Bcl-2, Bax and Cytochrome c protein were detected by Western blotting, and caspase-3 activity was measured by a microplate reader. **Results:** The results showed that after treatment with corilagin nanoparticle, the expression of Bcl-2 was down-regulated and Bax was up-regulated compared to that in the control. Corilagin nanoparticle treatment reduced the mitochondrial transmembrane potential, increased the release of cytochrome c and induced caspase 3 activation. **Conclusion:** Corilagin nanoparticle can inhibit the proliferation of SGC-7901 cells and induce apoptosis via mitochondrial pathway.

Keywords: corilagin nanoparticle; human gastric cancer; apoptosis

S1.50

A novel apoptosis inducer WF-210 activates procaspase-3 through sequestering inhibitory zinc ions

Fang-yang WANG^{1,2}, Li-hui WANG^{1,2}, Shuang WANG^{1,2}, Yang-fang ZHAO^{1,2}, Ping GONG^{1,2}, Jing-yu YANG^{1,2,*}, Chun-fu WU^{1,2,*}. ¹Department of Pharmacology, Shenyang Pharmaceutical University, 103 Wenhua Road, 110016 Shenyang, China; ²Benxi Medicine Institute, Shenyang Pharmaceutical University, Benxi 177005, China

*To whom correspondence should be addressed.

E-mail: yangjingyu2006@gmail.com (Jing-yu YANG); wucf@syphu.edu.cn (Chun-fu WU)

Aim: The direct induction of apoptosis has emerged as a powerful anticancer strategy, and small molecules that either inhibit or activate certain proteins in the apoptotic pathway have great potential as novel anticancer drugs. Increased expression of procaspase-3 had been found in numerous tumors. Described herein is the investigation of the effect and mechanism of a novel small compound WF-210. **Methods:** The procaspase-3 activate activity was detected using a modified molecular docking and enzymatic based *in vitro* method. The MTT method and Annexin-V/PI co-stain method were used to detect cytotoxic and apoptosis of U-937 cells. Western blot analysis was performed for detecting the expression of apoptosis-related proteins. To further confirm that the apoptosis induced by WF-210 was caspase-3-dependent, chemically synthesized siRNAs was used to silence caspase-3 in HL60 cells with high expression of procaspase-3. Transfection of HL60 cells with siRNAs reduced caspase-3 expression by 60%–70%. Eukaryotic expression plasmid encoding procaspase-3 was transiently transfected into MCF-7 cells which was procaspase-3 deficient. Further study of the procaspase-3 activate mechanism was assayed using recombinant human procaspase-3 protein in buffer with or without zinc. **Results:** In a modified molecular docking and enzymatic based assay, WF-210 markedly bound and activated procaspase-3 in a concentration-dependent manner with EC₅₀ (0.85 μmol/L). WF-210 effectively induced apoptosis of U937 cells in a concentration-dependent manner. Cotreatment with different caspase inhibitor and WF-210 partly blocked cell apoptosis. The block rates of van-caspase and caspase-3 inhibitors were higher than caspase-8 and caspase-9 inhibitors. After treatment with WF-210, caspase-3, caspase-8, and caspase-9 were activated in HL60 and U937 cells, and procaspase-3 was activated at first. Silence procaspase-3 gene expression in HL60 could block apoptosis induced by WF-210. Transient transfection of caspase-3 into MCF-7 cells could restore apoptosis induced by WF-210. In further study, WF-210 markedly activated procaspase-3 in a concentration-dependent manner in buffer with zinc, showed no activity in buffer without zinc. **Conclusion:** Taken together, our results suggest that WF-210 activates procaspase-3 *in vitro* by sequestering inhibitory zinc ions, and inducing cell apoptosis in a caspase-3 dependent manner.

Keywords: Anticancer; Procaspase-3; Apoptosis; Zinc

Acknowledgements: This research was supported by the National High Technology Research and Development Program of China (2012AA020305), National Natural Science Foundation of China (No 81102470) and Liaoning Science and Technology Program (No 2011412004-3).

S1.51

Flavokawain B, a kava chalcone, induces apoptosis of non-small cell lung cancer by through Bax-initiated mitochondrial pathway

Jing WANG, Qi-yu ZHU, Zhe PEI, Zhong-qiu ZENG, Ya-xiong TANG. Chengdu Institute of Biology, Chinese Academy of Sciences, Chengdu 610041, China

Aim: Flavokawain B, a chalcone from kava root extracts, possesses strong anti-neoplastic activity against many cancer cells. Here the antitumor activity and molecular mechanisms in lung cancer cells were investigated. **Methods:** The anticancer potential of Flavokawain B in lung cancer H460 and A549 cells were tested *in vitro*. **Results:** FKB significantly inhibited cell proliferation and caused arrest of the cell cycle G₂-M in a dose-dependent manner. FKB also induced apoptosis, which was associated with cytochrome c release, caspase-7 and caspase-9 activation and Bcl-x_L/Bax dys-regulation. FKB significantly down-regulated survivin and XIAP, and the inhibitory effect induced by FKB was greatly attenuated by through overexpression of survivin or Bax^{-/-} MEFs. Furthermore, FKB activated the mitogen-activated protein kinases and the JNK inhibitor SP600125 significantly decreased the growth-inhibitory and apoptotic effects of FKB. **Conclusion:** These results suggest the anti-lung cancer potential of Flavokawain B for the prevention and treatment of lung cancer.

Keywords: apoptosis; Bax; Flavokawain B; JNK; lung cancer; survivin

Acknowledgements: This work was financially supported by the National Natural Science Foundation of China #81173095 (Ya-xiong TANG) and by the Major Research plan of the National Natural Science Foundation of China #91013006 (Jian SUN).

S1.52

Polyamine analogues tetrabutyl propanediamine induced apoptosis of human leukemia HL-60 cells

Qing WANG^{1,2}, Yan-lin WANG^{1,2}, Chun-yu CAO^{1,2,*}. ¹Medical College, Three Gorges University, 8 Daxue Road, Yichang 443002, China; ²Institute of Molecular Biology, Three Gorges University, 8 Daxue Road, Yichang 443002, China

*To whom correspondence should be addressed.

E-mail: ccy_ctgu@yahoo.cn

Aim: High concentration of polyamines are required for quick proliferation of tumor cells, and an exhaustion in intracellular polyamines will inhibit tumor cell growth and induce apoptosis. Polyamine analogues can promote polyamine degradation and thus to induce apoptosis in tumor cells. Therefore, it is necessary to design and develop new polyamine analogues for clinical trial or mechanism research which target polyamine metabolism. In the present study, the impact of a new polyamine analogue, tetrabutyl propanediamine (TBP), on the proliferation and apoptosis of human leukemia HL-60 cells were observed. **Methods:** HL-60 cells were treated with 50 $\mu\text{mol/L}$ TBP for 24, 48, 72 h. Proliferation rate of HL-60 cells was assayed by MTT. The flow cytometry was performed to identify cell cycle. The DNA fragmentation analysis was used to evaluate apoptosis. Western blot was used to identify the expression of apoptosis-related protein. **Results:** TBP dramatically inhibited proliferation of HL-60 cells and the inhibitory effect was in a time and dose dependent manner. The results from flow cytometry analysis indicated that TBP induced cell cycle arrest in G_0/G_1 . After TBP treatment, typical DNA fragmentation for apoptotic cells was observed by agarose gel electrophoresis. Furthermore, the contents of proapoptotic protein Bax and cytochrome *c* in the cytosol were significantly increased. **Conclusion:** TBP can markedly inhibit the proliferation of human leukemia HL60 cells by arresting cell cycle and inducing apoptosis. These results first showed that TBP, as an asymmetric analogue of putrescine, has a potential value in clinical treatment of leukemia.

Keywords: polyamines; tetrabutyl propanediamine; leukemia; apoptosis

S1.53

A standardized extract from *Paeonia lactiflora* and *Astragalus membranaceus* inhibits proliferation and induces apoptosis in human hepatoma cell lines

Jing-jing WU, Shan-shan HU, Sen ZHANG, Wu-yi SUN*, Wei WEI*. *Institute of Clinical Pharmacology, Anhui Medical University, Key Laboratory of Anti-inflammatory and Immune Medicine (Anhui Medical University), Ministry of Education, Hefei 230032, China*
*To whom correspondence should be addressed.

Aims: *Paeonia lactiflora* and *Astragalus membranaceus* are two traditional Chinese medicines, commonly used in Chinese herb prescription to treat liver disease. Protective effects of the extract prepared from the roots of *Paeonia lactiflora* and *Astragalus membranaceus* (PAE) on liver fibrosis were demonstrated in previous investigations. The present study was designed to investigate whether PAE effect in the growth-inhibition and apoptosis-induction of human hepatoma cell lines HepG2 and SMCC-7721. **Methods:** After treatment with PAE, the proliferation of HepG2 cells and SMCC-7721 cells was determined by methyl thiazolyl tetrazolium (MTT) and the apoptosis of these two human hepatoma cell lines was measured by flow cytometry with propidium iodide (PI) technology. The expression of Bcl-2, Bax and cleaved caspase-3 was detected by Western blotting, while the expression of Bcl-2 and Bax was also detected by immunohistochemistry. **Results:** Treatment with PAE (10-200 mg/L) caused an inhibitory effect on cell proliferation in hepatoma cell lines HepG2 and SMCC-7721. Meanwhile, PAE induced HepG2 cells and SMCC-7721 cells to undergo apoptosis, which was demonstrated by PI staining. Furthermore, Western blotting and immunohistochemistry revealed that the expression of Bcl-2 was significantly decreased while Bax and cleaved caspase-3 was increased in HepG2 cells and SMCC-7721 cells after treatment by PAE (10-200 mg/L). **Conclusion:** Our results suggested that PAE inhibits hepatoma cell proliferation and induces cell apoptosis. PAE may induce apoptosis in HCC cells by regulating Bcl-2 families.

Keywords: PAE; hepatocellular carcinoma; apoptosis; HepG2; SMCC-7721

Acknowledgements: This work was supported by grants from Specialized Research Fund for the Doctoral Program of Higher Education of China (No 20113420120002), Natural Science Foundation of the Higher Education Institutions of Anhui Province (NoKJ2012A153), the National Natural Science Foundation of China (30973543, 81173075). The authors acknowledge the help of the staff members of the Institute of Clinical Pharmacology, Anhui Medical University in conducting the study.

S1.54

Cinobufacini inhibits proliferation of human cervical cancer Caski cells by inducing apoptosis

Yan XIA^{1,2}, Jian-lin YANG^{1,2}, Chun-yu CAO^{1,2,*}, Yu HAN^{1,2}, Yan-lin WANG^{1,2,*}. ¹Medical College, Three Gorges University, 8 Daxue Road, Yichang 443002, China; ²Institute of Molecular Biology, Three Gorges University, Yichang 443002, China

*To whom correspondence should be addressed.

E-mail: fzswangyl@ctgu.edu.cn (Yan-lin WANG); ccy_ctgu@yahoo.cn (Chunyu CAO)

Aim: To investigate the effects of cinobufacini on proliferation and apoptosis in human cervical cancer Caski cells. **Methods:** MTT assay was used to assay

cell proliferation; the flow cytometry assay was used to determine cell cycle; DNA fragmentation assays were used to evaluate apoptosis; Western blot was used to analyze cytochrome *c* content in the cytosol; the level of reactive oxygen species (ROS) was determined by fluorescence histochemistry; the mitochondrial membrane potential was determined by Dioc6-staining method. **Results:** Cinobufacini significantly inhibited the proliferation of Caski cells in a time- and dose-dependent manner. The concentration of median inhibition (IC_{50}) at 48 h is 1.68 $\mu\text{g/mL}$. Cinobufacini treatment resulted in DNA fragmentation, elevated ROS levels, increased release of cytochrome *c* from mitochondria to the cytosol and a decrease in mitochondrial membrane potential. Meanwhile, cinobufacini inhibited cell division and led to cell cycle arrest in G_2/M phase. **Conclusion:** Cinobufacini significantly inhibits the proliferation of Caski cells. The underlying mechanism is associated with the increase of ROS generation, which induced endogenous apoptosis and inhibited cell division. The results of this study suggest that cinobufacini has a potential value for the clinical treatment of cervical cancer.

Keywords: cervical cancer cell; apoptosis; cinobufacini

S1.55

Mitogen-activated protein kinases inhibitors exerts anti-fibrosis by mediating TGF- β /Smad signaling in HepG2 cells and HSC cells

Lei XIE^{1,2}, Yan YANG¹, Wen-juan RUI¹, Jie-mei JIANG¹, Xiang-peng HU¹, Chao WU¹. ¹Department of Pharmacology and Institute of Natural Medicine, Anhui Medical University, Hefei 230032, China; ²Department of Anesthesiology, Maternal and Child Care Hospital of Anhui province, Hefei 230001, China

Aims: To define further the mechanisms of TGF- β /Smad signal transduction mediated by mitogen-activated protein kinases (MAPKs), including the extracellular signal-regulated protein kinase (ERK), c-Jun N-terminal kinase (JNK) and p38 pathways, in HepG2 cells and HSC cells. **Methods:** In the absence or presence of three MAPK (ERK, JNK and p38)-specific inhibitors, HepG2 cells and HSC cells were stimulated with exogenous TGF- β to activate Smad signalling. plasminogen activator inhibitor (PAI)-1 transcriptional activity was measured by real-time reverse transcriptase-polymerase chain reaction analysis. **Results:** TGF- β induced PAI-1 mRNA expression in HepG2 cells and HSC cells; in addition, the three MAPK (ERK, JNK and p38)-specific inhibitors decreased PAI-1 mRNA expression. **Conclusion:** These results suggested that the ERK, JNK and p38 pathways mediate TGF- β /Smad signal transduction and might be considered as specific targets of drug therapy for hepatocellular carcinoma.

Keywords: HepG2 cells; HSC cells; mitogen-activated protein kinase; plasminogen activator inhibitor 1; Smad; TGF- β

S1.56

The effects of extracts from *Astragalus membranaceus* Bge (AMB) on HepG2 cell

Shui-xiang XIE¹, Li-chun WEN¹, Wen-feng HUANG², Jun YUAN², Xiao-li WANG¹, Qiong-jun XIE¹, Lin-bo CHEN^{1,*}. ¹Gannan Medical University, Ganzhou 341000, China; ²Department of digestion and ambury, the first affiliated hospital of Gannan Medical University, Ganzhou 341000, China

*To whom correspondence should be addressed.

Aim: To observe the effect of extracts from *Astragalus membranaceus* Bge (AMB) on HepG2 cell and its molecular mechanism. **Methods:** HepG2 cells were cultured in 2 groups: control group and AMB group; MTT assay and RT-PCR were used to test the variation of the two groups. **Results:** We observed that the cell population and parapodium of HepG2 decreased transparently in AMB group compared with the control group, and some cell floated. The expression of p53 increased and Bcl-xL decreased. **Conclusion:** The extracts of *Astragalus membranaceus* Bge (AMB) can restrain the growth of HepG2 cell, its mechanism may be the activation of p53 and the suppression of Bcl-xL.

Keywords: *Astragalus membranaceus* Bge; HepG2; Bcl-xL; p53

S1.57

Investigation of protective effect of *Luobuma* (*Apocynum venetum*) extract and its metabolites against tert-butyl hydroperoxide (t-BHP) induced human hepatoma HepG2 cell injury

Wen-yan XIE, Xiao-ying ZHANG*. *College of Veterinary Medicine, Northwest A&F University, Yangling 712100, China*

*To whom correspondence should be addressed.

Aims: According to previous publications, *Luobuma* (*Apocynum venetum*) extract showed protective effect for liver damage in *in vitro* and *in vivo* models, and the bioactive compounds were considered to be flavonoids that included in *Luobuma* extract. However, after oral administration, flavonoids are extensively metabolized

during absorption, which means only their metabolites instead of the mother flavonoids can reach liver and represent activities. In this work, we investigate the protective effect of Luobuma extract, as well as p-hydroxyphenylacetic acid (p-HPAA) and 3,4-dihydroxyphenylacetic acid (DOPAC) that are two metabolites of its main compounds (hyperoside and quercetin), against tert-Butyl hydroperoxide (t-BHP) induced human hepatoma HepG2 cell injury. **Methods:** HepG2 cells were pre-incubated with Luobuma extract, p-HPAA or DOPAC for 2 or 24 h, and then treated with t-BHP for 3 h. Subsequently, cell viability and alanine aminotransferase (ALT) release were measured by MTT assay and fluorometric assay, respectively. **Results:** After 2 h pre-incubation, 40–80 µg/mL of Luobuma extract increased 25% cell proliferation and inhibited ALT release by almost 40%, but p-HPAA and DOPAC at tested concentrations showed no effect compared to the control. After 24 h pre-incubation, the protective effect of Luobuma extract was decreased, while DOPAC at concentrations of 0.5–5 µg/mL showed significant stronger beneficial effect compared to the cells treated t-BHP, no obvious change was observed in p-HPAA treated cells. **Conclusion:** Our results suggest that after biotransformation, Luobuma extract can protect hepatocytes from oxidant attack via its metabolites, which probably act through regulation of specific gene expression since the prolonged protective effect of the metabolites.

Keywords: *Apocynum venetum*; HepG2 cells; ethanol; paracetamol; t-BHP

Acknowledgements: This work was supported by the Ministry of Education and State Administration of Foreign Experts Affairs “overseas teacher” project (No MS2011XBNL057) and the Open Project Program of State Key Laboratory of Natural Medicines, China Pharmaceutical University (No SKLNMKF201221).

S1.58

Caffeic acid 3,4-dihydroxyphenethyl ester arrests colorectal cancer cells growth by upregulating phosphorylation of p38 MAPK

Wen-xiu XIN¹, Yan-hua ZHANG¹, Xin XIE¹, Ying-hui TONG¹, Zhi-zhen ZHANG²*, Xiao-yuan LIAN¹*, ¹College of Pharmaceutical Sciences, Zhejiang University, Hangzhou 310058, China; ²Ocean College, Zhejiang University, Hangzhou 310058, China

*To whom correspondence should be addressed.

E-mail: xlyian@zju.edu.cn (Xiao-yuan LIAN); zzhhang88@zju.edu.cn (Zhi-zhen ZHANG)

Aims: Caffeic acid 3,4-dihydroxyphenethyl ester (CADPE) has been reported to possess a broad spectrum of antitumor activity; however, its antitumor mechanism is not well understood. In this study, we investigated the influence of CADPE on cell cycle and its underlying mechanisms using human colorectal cancer cell lines.

Methods: First, the anticancer activity of CADPE in human colon cancer cells was determined by measuring the cell growth and the colony formation in the presence or absence of CADPE. Second, the effects of CADPE on cell cycle and the related cell cycle-regulating proteins were analyzed using flow cytometry and Western blot, respectively. Finally, a potential role of p38 MAPK in the anticancer activity of CADPE was explored by using chemical inhibitors of three components of MAPK signaling pathways and determining whether any of the inhibitors was able to abolish or attenuate the CADPE-induced alterations of above parameters. **Results:** CADPE inhibited cell growth in a variety of human colorectal cancer cell lines with IC₅₀ values ranging from 4.06 to 17.60 µmol/L, and markedly reduced the colony formation of cancer cell lines. Consistently, CADPE significantly arrested cell cycle progression at G₀/G₁ phase in measured cell lines with a corresponding and robust decrease in the protein levels of Cdc25A, cyclin D1, cdk2, cdk4, and p-Rb (Ser780), suggesting a crucial role of inhibition of Cdc25A-Cdk2/4-Rb pathway in CADPE-induced cell cycle arrest. Furthermore, CADPE substantially activated p38 MAPK pathway as indicated by a considerable and sustained increase in the level of p-p38 MAPK (Thr180/Tyr182) and the inhibition of p38 MAPK with SB239063 substantially suppressed CADPE-induced arrests of colon cancer cell growth and cell cycle progression as well as CADPE-induced inhibition of Cdc25A-Cdk2/4-Rb pathway, whereas inhibition of ERK1/2 or JNK had no these effects. **Conclusion:** These results demonstrate that activation of p38 MAPK that leads to inhibition of Cdc25A-Cdk2/4-Rb pathway plays a key role in CADPE-induced cell cycle arrest as well as in the anticancer activity of CADPE.

Keywords: CADPE; human colon cancer cells; cell cycle; Cdc25A-Cdk2/4-Rb pathway; p38 MAPK

S1.59

Tangeretin enhances the therapeutic effect of 5-Fu against colon cancer *in vivo* and *in vitro* by inhibiting miR-21 at low concentration

Yao XIONG, Li-hong WAN, Li-ming ZHOU*. Department of Pharmacology, West China School of Preclinical and Forensic Medicine, Sichuan University, Chengdu 610041, China

*To whom correspondence should be addressed.

Aim: Tangeretin exhibits anti-proliferative, anti-invasive, anti-metastatic and antioxidant activities under several cancer models *in vivo*. In this study, we evaluate the effects of tangeretin alone or in combination with 5-FU on cell viability and apoptosis in colon cancer cell lines. **Methods:** Effects of tangeretin and 5-FU combination or alone on the cell proliferation were tested in colon cancer cells HCT-116 and SW-480 by MTT assay. Iso-biographic analysis was employed to evaluate the combination effect of tangeretin and 5-FU. Apoptosis in cells were examined by DAPI-staining. The mRNA level of miR-21 (a signature of tumor aggressiveness) was determined by real-time PCR. Two different xenograft studies also used to evaluate the effect of co-administration of tangeretin and 5-FU. **Results:** Combination of 5-FU and tangeretin has a synergistic effect. Tangeretin at low concentration (25 µg/mL) could markedly enhance the therapeutic effect of 5-FU compared to 5-FU groups ($P < 0.05$). Low concentrations of tangeretin and 5-FU promote apoptosis and decreased mRNA level of miR-21 ($P < 0.01$). *In vivo* mouse xenograft studies showed that coadministration of tangeretin and 5-FU reduced tumor volumes without bodyweight losing. **Conclusion:** Our data indicate that combination of tangeretin with 5-FU could improve the efficacy of chemotherapy for colon tumors.

Keywords: tangeretin; 5-FU; colon cancer; synergistic effect; miR-21

S1.60

Recent advances in anticancer research of homoharringtonine

Bin XU. Shanghai Institute of Materia Medica, Chinese Academy of sciences, Shanghai, 200031, China

Homoharringtonine (HHT) was studied in our Institute for many years. In collaboration with other institutions it was proved as an effective antileukemic drug in China in 1978. Afterwards, In our Country HHT was manufactured in several pharmaceutical companies and used clinically as one of basic antitumor drugs approved by Chinese Ministry of Health in 1993. Meanwhile and later, more and more papers were published regarding its action mechanism of antitumor activity and clinical efficacy for treatment of different kinds of leukemias. In recent years in USA, several laboratories and clinical hospitals conducted systematic study on this drug and found some new uses for different malignancies. In particular, HHT is demonstrated as a good therapeutic drug for chronic myelocytic leukemia (CML) including patients who have developed resistance to targeting agents such as imatinib or dasatinib. It attracts much attention from medical circles. In October 2012 FDA approved this drug for treatment of CML patients. Semisynthetic preparation of HHT called omacetaxine mepesuccinate (ceflatonin, CGX-653, Myelostate) has been produced and recommended for clinical application in USA and other countries. When the drug is injected by subcutaneous (SC) route, the bioavailability is excellent. It has been reported that HHT was given as a single agent to 71 patients with late CML-CP (chronic phase of CML) patients at a daily dose of 2.5 mg/m² for 14 d during the remission induction phase, and for 7 d monthly during the maintenance phase. A complete hematologic response was obtained by 72%, and cytogenetic response was achieved by 31%. The major toxicity was myelosuppression. The mechanism of action of omacetaxine is related to the induction of apoptosis in a Bax-independent fashion through mitochondrial disruption and release of cytochrome *c*, leading to caspase-9 and caspase-3 activation in leukemic cells. Gene expression analysis showed that the transforming growth factor-beta (TGF-β) and the tumor necrosis factor (TNF) signaling pathways appear to play an important role in induction of apoptosis by omacetaxine.

Keywords: homoharringtonine; chronic myelocytic leukemia; FDA approval

S1.61

Anti-cancer effects of triterpenoids isolated from *Alismatis Rhizoma* on HepG2 cells

Wen XU¹*, Ting LI²*, Ming-qing HUANG¹*, Xiu-ping CHEN², Jin-jian LU²*. ¹College of Pharmacy, Fujian University of Traditional Chinese Medicine, Fuzhou 350108, China, ²State Key Laboratory of Quality Research in Chinese Medicine, Institute of Chinese Medical Sciences, University of Macau, Macao, China

*These authors contributed equally to this research

*To whom correspondence should be addressed.

E-mail: hmql1115@126.com; jinjianlu@umac.mo

Aim: This study aims (1) to isolate different types of compounds from *Alismatis Rhizoma*, a well-known traditional Chinese medicine, (2) to compare the anti-cancer activities *in vitro*, and (3) to study the structure-function relationship. **Methods:** Silica gel column chromatography and preparative HPLC were used to isolate the compounds and spectroscopic techniques including UV, MS, ¹H-NMR, and ¹³C-NMR were employed to elucidate the structures. MTT assay and observation

of morphological changes were performed to evaluate the cell proliferation. Flow cytometry was used to detect cell cycle distribution in different phases. **Results:** Thirteen protostane triterpenoids, including alisol A, alisol A 23-acetate, alisol A 24-acetate, 16-di-hydroxy-alisol A 23-acetate, 16-di-hydroxy-alisol A 24-acetate, alisol B, alisol B 23-acetate, 11-deoxyalisol B, alisol C, alisol C 23-acetate, alisol F, alisol G, and alisol L, together with sitosterol, were obtained from the rhizome of *Alismatis Rhizoma*. The anti-cancer activities of alisol A, alisol A 24-acetate, alisol B, alisol B 23-acetate, alisol C, alisol C 23-acetate, alisol L, alisol F and sitosterol were screened, and alisol B showed the most effective anti-cancer activity as confirmed by the results of proliferation inhibition and cell numbers in sub-G1 section. Base on structure-activity relationship, C-16 hydrogen is critical to the anti-cancer activity of these alisol derivatives. **Conclusion:** The anti-cancer potential of alisol B is deserved further evaluation and C-16 hydrogen plays an important role for receiving their anti-cancer effects.

Keywords: *Alismatis Rhizoma*; triterpenoid; alisol B, cancer; structure-activity relationship.

Acknowledgements: This study was supported by the Provincial Natural Science Foundation of Fujian (2012J05152) and the Research Fund of University of Macau (SRG026-ICMS13-LJJ and MRG008/CXP/2013/ICMS).

S1.62

Furanodiene induces G1 cell cycle arrest and activates the signaling molecules regulating apoptosis and autophagy, and slightly inhibits key steps of cell metastasis in 95-D lung cancer cells

Wen-shan XU, Ting LI, Guo-sheng WU, Yuan-ye DANG, Xiu-ping CHEN, Jin-jian LU*, Yi-tao WANG*. State Key Laboratory of Quality Research in Chinese Medicine, Institute of Chinese Medical Sciences, University of Macau, Macao, China

*To whom correspondence should be addressed.

E-mail: jinjianlu@umac.mo; ytwang@umac.mo

Aim: This study aims to systematically investigate the effects of furanodiene (FUR), a natural terpenoid isolated from *Rhizoma Curcumae*, on cell viability and significant processes of tumor progression in 95-D lung cancer cells. **Methods:** MTT assay and flow cytometry were used to detect the anti-proliferative effects and cell cycle distribution, respectively. Western blot assay was performed to detect the expression of proteins related to the different cellular signal pathways. Cell adhesion was tested by cell attachments to matrigel and fibronectin using a microscope. Wound healing and transwell assays were employed to investigate cell migration and invasion, respectively. **Results:** FUR concentration-dependently inhibited cell proliferation and blocked the cell cycle progressions in G₁ phase via down-regulating the protein levels of cyclin D1 and CDK6, and up-regulating those of p21 and p27. FUR also affected the signaling molecules that regulate apoptosis revealed by the down-regulation of the protein levels of PARP, caspase-7, survivin, and the expression ratio of Bcl-2/Bax. Further studies showed that FUR induced expression of light chain 3-II in the protein level, indicating autophagy is involved in this process. However, the adhesive ability of 95-D cells to matrigel and fibronectin was only slightly inhibited after FUR treatment for 1 h. FUR also exhibited slight suppression on cell migration and invasion in 95-D cells. **Conclusion:** FUR induced G1 cell cycle arrest and activated the signaling molecules regulating apoptosis and autophagy, while slightly inhibited cell metastasis in 95-D lung cancer cells.

Keywords: furanodiene; cell cycle; apoptosis; autophagy; metastasis

Acknowledgements: This study was supported by the Research Fund of University of Macau (UL016/09Y4/CMS/WYT01/ICMS, MYRG208 (Y2-L4)-ICMS11-WYT and SRG026-ICMS13-LJJ) and the Science and Technology Development Fund of Macau Special Administrative Region (045/2011/A, 077/2011/A3, 074/2012/A3).

S1.63

Study on the purification and identification for anti-tumor activity of DL1106 and its analogue

Xiao-na XU, Lian-hua FANG^{1,2,*}, Guan-hua DU^{1,2,*}. ¹Beijing Key Laboratory of Drug Targets Identification and Drug Screening; ²State Key Laboratory of Bioactive Substances and Functions of Natural Medicines, Institute of Materia Medica, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing 100050, China

*To whom correspondence should be addressed.

E-mail: fanglh@imm.ac.cn (Lian-hua FANG); dugh@imm.ac.cn (Guan-hua DU)

Aim: To get enough quantity and quality of DL1106 and its analogue to evaluate their anti-tumor activity, the fermentation conditions of marine actinomycetes REL-3 were optimized. **Methods:** Estimating the effects of fermentation time

and the pH value and salt concentration of fermentation broth on the yield, the fermentation conditions of REL-3 were optimized. After extracting the collection of the fermentation liquid with equal volume of acetic acid ethyl three times, the relative content of the DL1106 and its analogue in the concentrated extracts was detected by HPLC. The crude DL1106 and its analogue were purified by preparative HPLC. The MTT assay was used to evaluate the inhibitory ability of DL1106 and its analogue on the growth of tumor cells, such as A549, HepG2, Skov3 and SY5Y. **Results:** We got the most objective production until the sixth day in the following fermentation conditions: the salt concentration of fermentation is 2.8% and the pH value is 7.2. We got DL1106 and its analogue with higher purity. DL1106 and its analogue have strong anti-tumor effect indicated by significant cytotoxicity in A549, HepG2, Skov3, and SY5Y cells. **Conclusion:** We got the optimal fermentation conditions of REL-3 and obtained almost pure DL1106 and its analogue. The MTT assay showed that DL1106 and its analogue have a favourable anti-tumor activity *in vitro*.

Keywords: REL-3; DL1106; analogue; HPLC; anti-tumor

Acknowledgements: This study was supported by the national scientific & technological major special projects "significant creation of new drugs" (2013ZX09103001-008) and special foundation on scientific and technological basic work (2007FY130100).

S1.64

Gecko Peptides Mixture induce apoptosis in human esophageal squamous carcinoma EC109 cell line *in vitro*

Xin-li XU, Jian-gang WANG, Rui-fang LI, Leng-xin DUAN, Chao-chu CUI, Wan-li ZHANG. The Key Laboratory of Pharmacology and Medical Molecular Biology, Medical College, Henan University of Science and Technology, Luoyang 471003, China

Aims: To investigate the effect of Gecko Peptides Mixture (GPM) on apoptosis of human esophageal squamous carcinoma EC109 cell line and the molecular mechanisms. **Methods:** EC109 cells were cultured *in vitro* and the inhibitory effect of GPM on their proliferation was measured by MTT assay. The apoptosis of EC109 cells was determined by Hoechst 33258 fluorescence staining. The effects of GPM on Bcl-2 and caspase-3 protein expression of EC109 cells were investigated by using immunohistochemical method. **Results:** GPM significantly inhibited proliferation of EC109 cells in a dose-dependent and time-dependent manner, the values of the IC₅₀ for 24, 48, and 72 h were 0.317 mg/mL, 0.227 mg/mL, and 0.183 mg/mL, respectively. EC109 cells treated with 0.1 and 0.2 mg/mL GPM for 24 h induced typical apoptotic morphological features, decreased Bcl-2 protein expression while increased caspase-3 expression in EC109 cells. **Conclusion:** GCPs exerts anti-tumor effects *in vitro* by inducing apoptosis, decreasing the bcl-2 expression and increasing caspase-3 expression.

Keywords: gecko peptides mixture; human esophageal squamous carcinoma; apoptosis; Bcl-2; caspase-3

S1.65

Growth inhibition and apoptosis induction effects on human tumor cell lines by actinomycin D and its analog from *Streptomyces parvus* REL-3

Sheng-qian YANG, Yu YAN, Huan-li XU, Yang-yang HE, Yong-jie LI, Li LI, Guan-hua DU. ¹Beijing Key Laboratory of Drug Targets Identification and Drug Screening, ²State Key Laboratory of Bioactive Substances and Functions of Natural Medicines, Institute of Materia Medica, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing 100050, China

Aims: As an important natural product resource, the diversity and chemical complexity of marine microorganism are treasury for development of new antitumor drugs. In the present study, growth inhibitory and pro-apoptosis effects of actinomycin D (Act D) and its analog (DL1106-2) from *Streptomyces parvus* REL-3 were investigated *in vitro*. **Methods:** Growth inhibitory effect of actinomycin D and its analog in seven human tumor cell lines, including promyelocytic leukemia cell (HL-60), oral epithelial carcinoma cell (KB), cervical cancer cell (HeLa), hepatocellular carcinoma cell (HepG2, Bel-7402), colorectal carcinoma cell (HCT-8) and lung adenocarcinoma cell (A549), was measured by using the microculture tetrazolium (MTT) assay after 72 h of treatment. Apoptosis of cells were evaluated by Hoechst 33342 staining, DNA agarose gel electrophoresis and flow cytometry. **Results:** Following exposure of tumor cells to Act D or DL1106-2 for 72 h, Act D or DL1106-2 at the concentration of 0.3 µg/mL completely inhibited the growth of tumor cell lines mentioned above. The IC₅₀ of DL1106-2 against cells were 4.3, 14.8, 3.3, 13.6, 13.4, 11.4, and 9.3 ng/mL, while the IC₅₀ of Act D were 7.8, 29.0, 9.5, 35.8, 74.7, 36.1, and 27.2 ng/mL, respectively. These results showed that the growth inhibitory effect of DL1106-2 in these tumor cells was stronger than that of

Act D. Results from Hoechst 33342 staining, DNA agarose gel electrophoresis, and flow cytometry demonstrated that DL1106-2 concentration-dependently induced apoptosis in the tumor cells, and also had more remarkable apoptosis-inducing activity than Act D. **Conclusion:** DL1106-2 can inhibit the growth of many human tumor cell lines by inducing apoptosis, which is similar to Act D. These data suggest that DL1106-2 has the potential to be used as a new cytotoxic anticancer drug.

Keywords: actinomycin D; cancer; growth inhibition; apoptosis; *Streptomyces parvus*

S1.66

Drug release characteristics and antitumor activity of doxorubicin-hyd-PEG-folate conjugate

Wei-liang YE, Si-yuan ZHOU*. *Department of Pharmaceutics, School of Pharmacy, Fourth Military Medical University, Xi'an 710032, China*

*To whom correspondence should be addressed.

E-mail: zhouys@fmmu.edu.cn

Aim: Doxorubicin (DOX) is a highly potent chemotherapeutic agent that is widely used to treat various types of cancers, including breast and urothelial cancer. However, its severe side effects, such as cardiotoxicity, remain as a major problem. To overcome these limits, more effective and active targeting system is needed to enhance intracellular uptake of drug in cancerous cells at the tumor sites. The aim of this paper is to investigate the drug release characteristics and antitumor activity of doxorubicin-hyd-PEG-folate conjugate. **Methods:** A PEG based folate mediated active tumor targeting delivery systems for DOX was designed and synthesized. In this system, DOX was conjugated with PEG by hydrazone bond (DOX-hyd-PEG-FA) or by amide bond (DOX-ami-PEG-FA). The drug loading of the conjugates was detected by HPLC method, and the drugs release characteristic was performed in different pH media *in vitro*. The KB cells (folate receptor over-expression) and A549 cells (folate receptor low-expression) were selected to study the cytotoxicity of the conjugates, meanwhile the effect of exogenous folic acid on the cytotoxicity of the conjugates was also observed. The uptake of DOX and its conjugates by tumor cells were analyzed by flow cytometry and fluorescence microscope. The intracellular distribution of conjugates was studied by using laser scanning confocal microscope. **Results:** The rate and amount of DOX release from the DOX-hyd-PEG-FA were strongly dependent on the pH of the medium. DOX-hyd-PEG-FA showed a much faster DOX release at pH 5.0 than at pH 7.4. The rate and amount of DOX release from the DOX-ami-PEG-FA was independent on the pH of the medium. Compared with DOX-hyd-PEG, DOX-hyd-PEG-FA could increase the intracellular accumulation of DOX in short time and showed a DOX translocation from lysosomes to nucleus in KB cells. DOX-hyd-PEG-FA could deliver DOX to nucleus in KB cells in a time dependent manner. The cytotoxicity of DOX-hyd-PEG-FA on KB cells was much higher than that of DOX-ami-PEG-FA and DOX-hyd-PEG. Meanwhile, the cytotoxicity of DOX-hyd-PEG-FA on KB cells was much higher than that on A549 cells. The cytotoxicity and intracellular uptake of DOX-hyd-PEG-FA were inhibited by exogenous folic acid. These results indicate that the DOX-hyd-PEG-FA can provide excellent stability in blood circulation. FA presented in the DOX-hyd-PEG-FA plays an important role in enhancing the cytotoxic effect. Both passive and active tumor targeting abilities and acid sensitivity of DOX-hyd-PEG-FA were likely to act synergistically. **Conclusion:** DOX-hyd-PEG-FA can be selectively transported into the tumor cells that folate receptors are highly expressed via folate receptor-mediated endocytosis, and then the DOX is released and exert anti-tumor effect. Based on the results obtained in this study, the *in vivo* antitumor activity of DOX-hyd-PEG-FA is currently further evaluated on athymic nude mice bearing KB cells tumors.

Keywords: poly(ethylene glycol) (PEG); pH-sensitive; tumor-targeted; doxorubicin; folate receptor; cellular uptake; cytotoxicity

Acknowledgements: This work was supported by Shaanxi Science and Technology Innovation Project (2012KTCL03-18).

S1.67

Attenuation of tumor formation by *Ardisia crispa* root in murine skin carcinogenesis model

Looi Ting YEONG¹, Abdul Hamid ROSLIDA¹, Saiful Yazan LATIFAH^{1,2}, Khaza'ai HUZWAH¹.

¹Department of Biomedical Science, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, 43400 Selangor Darul Ehsan, Malaysia; ²Laboratory of Molecular Biomedicine, Institute of Bioscience, Universiti Putra Malaysia, 43400 Selangor Darul Ehsan, Malaysia

Aim: *Ardisia crispa* has been traditionally used to treat inflammation-related diseases. In this study, chemopreventive effect of its triterpene- and quinone-

rich fraction (TQRF) was investigated. **Methods:** Mice (group I-IV; *n*=10) were initiated with single dose of 7,12-dimethylbenz(a)anthracene (DMBA, 100 µg/100 µL) followed by repeated promotion of croton oil (1%) twice weekly (20 weeks). Group I (anti-initiation) received TQRF (30 mg/kg) 7 d before and after DMBA; group II (anti-promotion) received TQRF 30 min before each croton oil application; group III (anti-initiation/promotion) was treated as combination of group I and II. Two other groups served as vehicle control (group V) and treated control (group VI). **Results:** Group IV (carcinogen control) showed highest tumor volume (8.79±5.44) and tumor burden (3.60±1.17). Group I revealed marked increment of tumor incidence (40%) with prompted latency period of tumor formation (week 7). Group II showed significant reduction of tumor volume (3.11), tumor burden (3.00) and tumor incidence (11.11%), along with prominent delayed of latency period of tumor formation (week 12). Group III revealed only 20% of tumor incidence, tumor burden (3.00±1.00) and tumor volume (2.40±1.12), which was significantly different from group IV. No tumor formation was observed in groups V and VI. **Conclusion:** This study evidences the inhibitory effects of TQRF during promotion period which further suggests the plausibility of each compound as chemopreventive agent.

Keywords: *Ardisia crispa*; skin cancer; chemoprevention; triterpene; quinone

S1.68

Molecular basis of eIF3a gene in therapeutic response of lung cancer patients to platinum-based chemotherapy

Ji-ye YIN, Xiao-jing XU, Hong-hao ZHOU, Zhao-qian LIU. *Institute of Clinical Pharmacology, Hunan Key Laboratory of Pharmacogenetics, Central South University, Changsha 410078, China*

Drug resistance is one of the major obstacles in successful chemotherapy of lung cancers. However, the mechanism of resistance is not yet clearly understood. Recently, eIF3a that plays an important role in translational control and regulation of gene expression was found to correlate negatively with prognosis in human cervical and esophageal cancers. We, thus, hypothesize that eIF3a may be an important factor in lung cancer response to chemotherapy by regulating the expression DNA repair proteins. Examination of 211 lung cancer tissue samples revealed that eIF3a expression negatively correlates with responses of the patients to cisplatin treatment. Ectopic over-expression or knockdown of eIF3a reduced or increased cellular resistance, respectively, to cisplatin, doxorubicin, and etoposide (VP-16) in cell lines as determined using MTT assay. Altering eIF3a expression also changed DNA repair activity as well as the expression of DNA repair proteins. We conclude that eIF3a possibly suppresses the synthesis of DNA repair proteins via translational control, which in turn contributes to the increased sensitivity to DNA-damaging anticancer drugs.

S1.69

2-Methoxyjuglone induces apoptosis in HepG2 human hepatocellular carcinoma cells and exhibits *in vivo* antitumor activity in a H22 mouse hepatocellular carcinoma model

Heng-yi YU¹, Xiao-qiong ZHANG², Xue Li¹, Fan-bo ZENG¹, Han-li RUAN¹. ¹Hubei Key Laboratory of Natural Medicinal Chemistry and Resource Evaluation, Faculty of Pharmacy, Tongji Medical College of Huazhong University of Science and Technology, Wuhan 430000, China; ²Hubei Institute for Food and Drug Control, Wuhan 430000, China

In order to discover anticancer agents from natural sources, an ethanol-soluble extract of the root bark of *Juglans cathayensis* was investigated and showed cytotoxic effects against various human cancer cell lines. A subsequent phytochemical study on the EtOAc-soluble fraction determined 2-methoxyjuglone (1) as one of the main active constituents. Compound 1 was shown to be cytotoxic against HepG2 cells. Morphological features of apoptosis were observed in 1-treated HepG2 cells, including cell shrinkage, membrane blebbing, nuclear condensation, and apoptotic body formation. Cell cycle analysis with propidium iodide staining showed that 1 induced cell cycle arrest at the S phase in HepG2 cells. Flow cytometric analysis with annexin V and propidium iodide staining demonstrated that 1 induced HepG2 cell apoptotic events in a dose-dependent manner (0-8 µg/mL). Western blot analysis of apoptosis-related proteins revealed that 1 induces HepG2 cell apoptosis through mitochondrial cytochrome c dependent activation of the caspase-9 and caspase-3 cascade pathway (intrinsic pathway). An *in vivo* experiment using tumor-bearing mice showed that treatment with 1 at 0.5 and 1.0 mg/kg per day decreased the tumor mass by 56% and 67%, respectively.

Keywords: *Juglans cathayensis*; 2-methoxyjuglone; apoptosis; intrinsic pathway; *in vivo* antitumor effect

Acknowledgements: Financial support from the National Natural Science

Foundation of China (No 30873361, 81072547 and 31270394), the Ministry of Science and Technology of the People's Republic of China (International Cooperative Project, grant No 2010DFA32430), and Wuhan Program for Science and Technology (No2013060501010158) is gratefully acknowledged.

S1.70

Src mediates extracellular signal-regulated kinase 1/2 pathway and autophagic cell death induced by cardiac glycosides in human non-small cell lung cancer cells

Yue-chen ZHAN, Yan WANG, Rong-guang SHAO, Dian-dong LI, Zhen WANG. *Institute of Medicinal Biotechnology, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing 100050, China.*

Aim: Our previous work demonstrated that extracellular signal-regulated kinase 1/2 (ERK1/2) pathway regulates autophagic cell death induced by representative CGs digoxin or ouabain in human non-small-cell lung cancer (NSCLC) cells. In this study, the role of Src in ERK1/2 signaling pathway as well as autophagy induced by the CGs was examined in A549 and H460 cells. **Methods:** Src inhibitor PP2 or siRNA were used to knock down phosphorylation or mRNA level of Src, respectively, along with the CGs treatment. Cellular viability was performed with MTT assay. Level of intracellular reactive oxygen species (ROS) was detected with H₂DCFDA staining, followed by flow cytometry. **Results:** Src activation was significantly observed simultaneously with mitogen-activated protein kinase kinase 1/2 (MEK1/2) and ERK1/2 activation upon the CGs treatment. Src inhibitor PP2 or siRNA knockdown could block either drug-induced MEK1/2 and ERK1/2 activation, and autophagic cell death in the cells. In addition, the level of ROS was markedly increased by CGs in a time-dependent manner, and this increase was significantly attenuated with PP2 pretreatment. **Conclusion:** CGs induces autophagy in human NSCLC cells through activation of Src-mediates ERK1/2 pathway. Besides, Src also plays an important role in CGs-induced ROS generation.

Keywords: cardiac glycosides; autophagic cell death; Src; lung cancer; reactive oxygen species

S1.71

Aspirin enhanced ABT-737-induced apoptosis via regulation of PI3K pathway in human cancer cells

Chong ZHANG¹, Lin-yi FENG¹, Xiao-dan SHUAI¹, Ya-si XU², Dan ZHANG², Da-yong ZHANG², Bao-ming WANG¹, Li-huang ZHANG¹, Jun-bo WANG¹, Jian-ping PAN¹, Hua ZHOU¹, Si-cong WANG¹, Neng-ming LIN². ¹School of medicine, Zhejiang university city college, Hangzhou 310015, China ; ²Laboratory of clinical pharmacy, Zhejiang cancer hospital, Hangzhou 310022, China

Aim: Observational studies show that regular use of aspirin after diagnosis was associated with longer survival among patients with mutated-PIK3CA colorectal cancer, but not among patients with PIK3CA wild-type cancer. ABT-737, a potent small-molecule inhibitor of antiapoptotic members of the Bcl-2 family, is a promising therapeutic agent for multiple malignancies. We examined the anti-cancer activity of 2 chemotherapeutic agents (aspirin and ABT-737) in several PIK3CA wild-type cancer cells. **Methods:** We analyzed the anti-tumor efficacy of aspirin and ABT-737 in human cancer cells *in vitro* by SRB cytotoxicity assay. Apoptosis and cell cycle were determined using flow cytometry analysis. Western blotting analysis was carried out to examine the expression of related molecules. AO staining was used to visualize Acidic autophagic vacuole. **Results:** In this study, we showed that combination treatment with clinically achievable concentrations of aspirin and ABT-737 could induce a synergistic growth arrest and apoptosis in several human PIK3CA wild-type cancer cells. In addition, we found that the combination of aspirin and ABT-737 resulted in a blockade of the PI3K signaling pathway. Furthermore, our data showed increased autophagy correlated with the resistance to aspirin or ABT-737 as single-agents, the enhanced autophagy induced by aspirin plus ABT-737 switched from a cytoprotective signal to a death-promoting signal. **Conclusion:** Our study indicated that the combination of aspirin and ABT-737 exerted synergistic anti-proliferative effects against PIK3CA wild-type cancer cells *in vitro*. We hope that this synergy may contribute to failure of aspirin cancer therapy and ultimately lead to efficacious regimens for cancer therapy.

Keywords: aspirin; ABT-737; solid tumors; apoptosis; autophagy

Acknowledgements: The authors gratefully acknowledge financial support from National Natural Science Foundation of China (81272473, 30872325), Zhejiang Provincial Foundation of National Science (Y2100682, LQ12H31001), Science Research Foundation of Zhejiang Health Bureau (2012KYA068, 2012KYB066), Zhejiang Provincial Program for the Cultivation of High-level Innovative Health talents (2010-190-4), Student Research Fund of Zhejiang University City College

(XZ2013562105, X2013562120), Scientific Research Fund of Zhejiang Provincial Education Department (Y201120633), Major Science Research Foundation of Zhejiang Health Bureau (WKJ2012-2-022), and Science and Technology Bureau of Hangzhou, Zhejiang Province, China (20120633B30).

S1.72

Heparan sulfate proteoglycans and cancer

Gan-lin ZHANG, Jin-ping LI, Yi ZHANG, Ming-wei YU, Guo-wang YANG, Xiao-min WANG*. *Beijing Hospital of Traditional Chinese Medicine, affiliated to Capital Medical University Beijing 100010, China*

*To whom correspondence should be addressed.

E-mail ntxm100@sina.com

Heparan sulfate proteoglycans (HSPGs) are widely distributed in extracellular matrix (ECM) of mammalian tissues and involved in a number of processes related to malignancy. They are composed of a core protein to which attached chains of the glycosaminoglycan, heparan sulfate (HS). HSPGs interact with many proteins including growth factors, chemokines and structural proteins of ECM to influence cell growth, differentiation, migration, and the cellular response to the environment. Recently, HSPGs are reported to have close relationship to cancer, especially tumor cell migration. This review describes current understanding of HSPGs in cancer and highlights new possibilities for therapeutic control.

Keywords: heparan sulfate proteoglycan; tumor; metastasis; growth factors

Acknowledgements: This paper was supported by National Natural Science Foundation of China 81202840 and Capital Medical University project: 12JL76.

S1.73

Temozolomide analogs overcome tumor resistance and induce DNA damage

Ji-hong ZHANG¹, Tracey BRADSHAW², Malcolm STEVENS², Yin-xian FAN¹. ¹Faculty of Medicine, Kunming University of Science and Technology, Kunming 650500, China; ²School of Pharmacy, University of Nottingham, Nottingham NG7 2RD, United Kingdom

Aim: Based on the molecular and resistance mechanism of alkylating agent temozolomide (TMZ), TMZ analogs's mechanism of action was investigated by *in vitro* anti-tumor activity assay, cell cycle and apoptosis analysis and DNA damage assay in glioma cells. **Methods:** the anti-tumor activity of temozolomide analogs was investigated by MTT assay with SNB19, U373 cells and the paired O6-methylguanine-methyltransferase (MGMT) transfected cells lines SNB19M and U373M; also the corresponding acquired resistance glioma cell lines SNB19VR and U373VR and the intrinsic TMZ resistance cell line HCT116 (mismatch repair gene *hMLH1* loss) were used as the screening models; the cell cycle analysis of TMZ analog 14112 were performed by flow cytometry in the SNB19V and SNB19M glioma cell lines; DNA double strand breaks (DSBs) induced by 14112 was detected by Western blot and immunofluorescence with SNB19V and SNB19M cells. In addition, DNA single strand breaks and DNA repair kinetics were analyzed by comet assay. Furthermore, the apoptosis caused by 14112 was investigated by PARP antibody in HCT116 cells.

Results: TMZ showed highly resistant to MGMT overexpressed cells, however, TMZ analogs were very active in TMZ resistance cell lines, showing equi-activity in the low and high MGMT expressed cells. Analog 14112 showed the most activity in the tested cell lines. Furthermore, TMZ analogs showed anti-tumor activity in the TMZ acquired resistant cell lines SNB19VR and U373VR. Analog 14112 caused cells arrest at G₂/M and apoptosis. Moreover, DNA strand breaks was observed with treatment time increased, the excision repair of 14112 was the similar to that of TMZ. The double strand breaks induced by 14112 were earlier than that of TMZ indicated by γ H2AX expression. **Conclusion:** The TMZ analogs can overcome TMZ resistance, the DNA damage induced by analog is not repaired by MGMT.

Keywords: glioma; resistance; temozolomide; analogs; DNA damage

S1.74

Jaridonin, a novel anticancer drug, induced human gastric cancer cell line MGC-803 autophagy and its possible mechanisms

Xu-yao ZHANG, Xiao-jing SHI, Dong CHEN, Ran WANG, Hong-min LIU*. *School of Pharmaceutical Sciences, New Drug Research & Development Center, Zhengzhou University, Zhengzhou 450001, China*

*To whom correspondence should be addressed.

Aim: Jaridonin, a novel ent-kaurane diterpenoid compound from *Isodon Rubescens* in our laboratory. We designed the current study to determine the effects of Jaridonin on human gastric cancer line MGC-803 autophagy and its possible mechanisms.

Methods: After treatment by Jaridonin for 24 h, the effect of Jaridonin on the MGC-803 cell viability was examined using MTT assay. Used the fluorescence microscope to observe the autophagosome and then used the flow cytometry to

determine the effects of Jaridonin on the ratio of the autophagy. Finally we used Western blot to detect the expressions of the proteins related with cell autophagy. **Results:** Jaridonin caused strong antiproliferative on MGC-803 cells in a dose- and time-dependent manner. Jaridonin also induced the produce of autophagosome in the MGC-803 cells. After the treatment of Jaridonin with MGC-803 cells 24 h, the expression of LC3I, LC3II were up-regulating, and the rate of LC3I/LC3II was increasing too, but the expression of PI3K, Akt and the phosphorylation of Akt were down-regulating. **Conclusion:** Jaridonin can significantly inhibit the growth of MGC-803 cells, and induce autophagy. The antitumor activity of Jaridonin might be attributed partly to the upregulation of LC3I, LC3II and the downregulation of PI3K, Akt, and the phosphorylation of Akt.

Keywords: ent-kaurene diterpenoid compound; anticancer; autophagy; molecular mechanism

S1.75

Rules of treatment and prescription-herbs on lung cancer in chinese medicine literatures

Yi ZHANG¹, Xiao-min WANG¹, Ming-wei YU¹, Wen-jie XU², Nan NAN¹, Cong MA¹.
¹Department of Oncology, Beijing Hospital of Traditional Chinese Medicine Affiliated to Capital Medical University, Beijing 100010, China; ²Beijing Institute of TCM, Beijing 100010, China

Aim: To explore the herbal administration rules, the syndrome and pathogenesis in TCM literatures. **Methods:** Collecting the prescriptions and herbs during and before the period of the Republic of China, analyzing data by frequency analysis. **Results:** Phlegm-dispelling and cough-suppressing medicinal, tonifying and replenishing medicinal, toxic herbs, qi-regulating medicinal, interior-warming medicinal, dampness-draining diuretic medicinal, dampness-resolving medicinal, exterior-releasing medicinal and blood-activating and stasis-resolving medicinal are the main types of herbs, qi and yin-tonifying medicinal are more commonly used. The herbs mainly belong to lung, spleen and stomach channels and characterized by the qi and flavor of pungency, sweetness, bitterness and warmth-heat. **Conclusion:** Reinforcing the healthy qi and restoring deficiency, dispelling phlegm and eliminating dampness, regulating qi and activating blood circulation and treating with poisonous agents are the main methods of treatment in TCM literatures. The qi and flavor of herbs is mainly pungency, sweetness, bitterness and warmth-heat. The channels of herbs is mainly lung, spleen, stomach. For further analysis, we preliminarily infer that lung cancer is the disease of root deficiency with insufficiency of qi and yin and branch excess with phlegm, dampness, blood stasis and toxin.

Keywords: TCM literatures; lung cancer; prescription-herbs; frequency analysis

Acknowledgements: The study was supported by the Capital Medical Development Foundation (SF-2007-1-08) and the Capital Public Health Project (Z111107067311044)

S1.76

Antiproliferation activity of Triterpenoid Saponins AG4 from *Ardisia Gigantifolia Stapf* on MCF-7 cells

Xiao-li ZHENG^{1,2}, Xian-zhe DONG¹, Li-hua MU¹, Hong-bo LIAO^{1*}, Ping LIU¹. ¹Department of Clinical Pharmacology, General Hospital of PLA, Beijing 100853, China; ²Tianjin University of Traditional Chinese Medicine, Tianjin 300193, China

*To whom correspondence should be addressed.

Aim: AG4 is a anti-tumor compound derived from *Ardisia gigantifolia Stapf*. The aim of this study was to investigate the influence of AG4 on different kinds of tumor cells and the possible mechanisms. **Methods:** We exposed 9 kinds of tumor cells to different concentrations of AG4 in order to investigate the effect of AG4 on cell viability by using the 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reduction assay. The appearance of MCF-7 cells treated by AG4 was measured by fluorescence microscopy. The apoptosis and cell cycle were detected by the flow cytometry. Activities of caspase-3 and caspase-9 were measured using activity quantitative detection kit. **Results:** AG4 showed proliferation inhibition in A549, BeL-7402, BGC-823, C6, EJ, HeLa, HepG2, MCF-7, and LS180 cells, and significantly in MCF-7 cells. Compared with control, the appearance has been changed and apoptosis apparently in MCF-7. Significant production of MDA was observed, the results also showed that AG4 down-regulated SOD activity and GSH level, and the activities of caspase-3 and caspase-9 had significant raise.

Simultaneously, AG4 significantly blocked the cell cycle in S phase. **Conclusion:** These results indicated that the inhibition on proliferation of MCF-7 cells was possibly through reactive oxygen species evoked, cycle blocking, and mitochondrial path way activation by AG4.

Keywords: *Ardisia gigantifolia Stapf*; cancer cells; apoptosis; cell cycle; caspase; redox system

S1.77

The anti-tumor effect of Sedum Lineare Thunb alcohol extract on S₁₈₀-bearing mice and its mechanism

Qing ZHOU, Li-zhen WU, Li-mei ZENG, Rong-yan SUI, Xing-ling CAO, Zhi-hua HUANG. Gannan Medical University, Ganzhou 341000, China

Aim: To observe the effects of Sedum Lineare Thunb (SLT) alcohol extract on tumor growth and to study its antioxidant ability and regulation of immune function. **Methods:** The second day of S₁₈₀-bearing mice model established, mice were randomly divided into control group, model group, cyclophosphamide (CTX) group (50 mg/kg), SLT low and high dose groups (4 and 8 g/kg), 10 per group, and continuously administrated for 14 d. The index of the inhibition rate of tumor weight was used to observe the anti-tumor activity of SLT. The levels of serum interleukin-6 (IL-6) and interleukin-10 (IL-10) were detected by Enzyme-linked immunosorbent assay (ELISA). Biochemical method was used to measure the activity of superoxide dismutase (SOD) and the content of malondialdehyde (MDA). Flow cytometer was used to survey the ratio of neutrophil and lymphocyte percentage in the white blood cell, the percentage of CD3⁺ positive cells in the lymphocytes and the percentage of CD4⁺ or CD8⁺ cells in the T-Lymphocytes, and then calculate the ratio of CD4⁺/CD8⁺. **Results:** The inhibition rate of the tumor growth in the SLT low and high dose groups (4 and 8 g/kg) are 84.26% and 62.24%, obviously higher than CTX group 29.28%. After treated with SLT, the thymus index was up-regulated, while the spleen index was down-regulated; the levels of serum IL-6 ($P < 0.05$) and IL-10 ($P < 0.001$) were increased significantly; the activity of SOD in serum was enhanced, but the content of MDA was reduced; the percentage of neutrophils was inhibited yet the percentage of lymphocytes and CD3⁺ lymphocyte were improved significantly; CD4⁺ lymphocyte T was obviously higher than that of the model group, however CD8⁺ T lymphocyte was lower; and CD4⁺/CD8⁺ was increased markedly. **Conclusion:** SLT alcohol extract had an anti-tumor effect on S₁₈₀-bearing mice, and its mechanism probably result of improving the antioxidant ability and immune function.

Keywords: Sedum Lineare Thunb; S₁₈₀-bearing mice; cytokine; antioxidation; immune function

Acknowledgements: This work was supported by grants from the health of Jiangxi province (2006B37).

S1.78

Synthesis and in vitro anti-cancer activity of four metal - fluorouracil complexes

Yi-ping ZHOU^{1*}, Yun ZHOU², Ying-jie CHEN³, Min LUO¹, Zhong-zheng SHI³, Wen-yuan ZHONG^{4*}. ¹School of Pharmaceutical Science, Kunming Medical University, Kunming 650500, China; ²Department of Chemistry, Yunnan Radio and TV University, Kunming 650000, China; ³School of Basic Medical Science, Kunming Medical University, Kunming 650500, China; ⁴Department of Chemistry, Kunming College, Kunming 650000, China

*To whom correspondence should be addressed.

Aim: In order to synthesize new efficient and low toxic metal-fluorouracil (5-FU) complexes anticancer drugs. **Methods:** Copper, iron, zinc salts and fluorouracil were used to synthesize copper, iron and zinc-fluorouracil complexes. Preliminary chemical structures of the 4 complexes were confirmed by elemental analysis and mass spectrometry. Their inhibitory activity on human cancer cells was measured by MTT colorimetric assay. **Results:** [Cu(5-FU)₂Cl₂], [Cu(5-FU)₂(NO₃)₂], [Fe(5-FU)₃SO₄], and [Zn(5-FU)₂Cl₂] were successfully synthesized. The IC₅₀ values of [Zn(5-FU)₂Cl₂] and [Fe(5-FU)₃SO₄] on HCT-116 cells were 9.2 and 18.1 μg/mL, lower than those of 5-FU and their metal salts counterparts. **Conclusion:** [Zn(5-FU)₂Cl₂] and [Fe(5-FU)₃SO₄] exhibited a synergic inhibitory effect with fluorouracil on cancer cell proliferation, their anti-cancer activity is worthy of further study.

Keywords: copper; iron; zinc-fluorouracil complexes; synthesis; anti-cancer activity