

**S5.1****Bone marrow mesenchymal stem cells reversed potassium channels remodeling in post-infarcted myocardium via regulating calcineurin pathway**

Ben-zhi CAI<sup>1</sup>, Nan CHEN<sup>1</sup>, Gang WANG<sup>1</sup>, Yan-ju LIU<sup>1</sup>, Fan YANG<sup>1</sup>, Yang WANG<sup>1</sup>, Zhen-wei PAN<sup>1</sup>, Bao-feng YANG<sup>1,2</sup>, Yan-jie LU<sup>1,2</sup>. <sup>1</sup>Department of Pharmacology, State-Province Key Laboratories of Biomedicine-Pharmaceutics of China, Harbin Medical University, Harbin 150081, China; <sup>2</sup>Institute of Cardiovascular Research, Harbin Medical University, Harbin 150081, China

**Aim:** Bone marrow mesenchymal stem cells (BMSCs) emerge as a new promising approach for treating heart diseases. However, the effects of BMSCs-based therapy on cardiac electrophysiological remodeling following myocardial infarction were largely unclear. This study was aimed to investigate whether BMSCs transplantation reverse cardiac potassium channels remodeling in post-infarcted hearts. **Methods:** Myocardial infarction was established in male SD rats, and BMSCs were then intramyocardially transplanted into the infarcted hearts after 3 d. Cardiac electrophysiological remodeling in the border zone was evaluated by Western blotting and whole-cell patch clamp technique after 2 weeks. **Results:** We found that BMSCs transplantation significantly ameliorated the increased heart weight index, the impaired left ventricular functions, and the enhanced cardiac fibrosis in infarcted hearts. The survival ratio of infarcted rats was also improved after BMSCs transplantation. Importantly, electrical stimulation-induced arrhythmias were seldom observed in BMSCs-transplanted infarcted rats compared with rats without BMSCs treatment. Furthermore, BMSCs transplantation effectively inhibited the prolongation of action potential duration and the reduction of transient and sustained outward potassium currents in ventricular myocytes from post-infarcted rats. Consistently, BMSCs-transplanted infarcted hearts exhibited the more expression of K<sub>v</sub>4.2, K<sub>v</sub>4.3, K<sub>v</sub>1.5, and K<sub>v</sub>2.1 proteins than infarcted hearts. Moreover, intracellular free calcium level, calcineurin and nuclear NFATc3 protein expression was shown activated in infarcted hearts, which was inhibited after BMSCs transplantation. **Conclusion:** Collectively, BMSCs transplantation prevented ventricular arrhythmias and reversed cardiac potassium channels remodeling in post-infarcted hearts via attenuating intracellular calcium overload and calcineurin/NFATc3 signal pathway, which help to advance our understanding of mesenchymal stem cells pharmacology and experimental therapeutics.

**Keywords:** bone marrow mesenchymal stem cells; myocardial infarction; arrhythmias; potassium channels; calcineurin

**S5.2****Expression and function of B cell-activating factor of the TNF family in T cell-mediated autoimmune arthritis**

Yan CHANG, Qiong QIN, Xiao-jing SUN, Xiao-yi JIA, Yu-jing WU, Ling-ling ZHANG, Wei WEI\*. Institute of Clinical Pharmacology, Anhui Medical University, Key Laboratory of Anti-inflammatory and Immune Medicine (Anhui Medical University), Ministry of Education, Hefei 230032, China

\*To whom correspondence should be addressed.

E-mail: wwei@ahmu.edu.cn.

**Aim:** To investigate the expression and function of B cell-activating factor (BAFF) and its receptors (BCMA, TACI, and BAFF-R) in T cell-mediated inflammatory immune responses of animal model for rheumatoid arthritis. **Methods:** The model of rat adjuvant-induced (AA) was induced by a single intradermal injection of 0.1 mL of CFA into the right hind metatarsal footpad of rat. The rats with experimental arthritis were randomly separated into different groups and then treated with TACI-Ig (0.7, 2.1, and 6.3 mg/kg) respectively once per day from d 16 to 34 after immunization. The degree of arthritis was evaluated by radiographic and histological examination. Levels of BAFF were assessed by enzyme-linked immunosorbent assay. Cluster of differentiation (CD)3, CD20, CD68, CD21, CD138, BAFF, and PNA<sub>d</sub> expression were detected by immunohistochemical analysis. (CD3<sup>+</sup>CD4<sup>+</sup>), (CD4<sup>+</sup>CD25<sup>+</sup>), (CD4<sup>+</sup>CD44<sup>+</sup>), and (CD4<sup>+</sup>CD62L<sup>+</sup>) markers on CD4<sup>+</sup> T cells were detected by flow cytometry. Double-labeled immunofluorescent staining was used to identify the expression of CD68 and BAFF receptor. Expression of BAFF and its receptors were analysed by immunohistochemical analysis, quantitative real-time PCR and Western blot analysis. **Results:** Compare with the control group, AA rats developed severe arthritis. This was characterized by marked ankle joint radiographic and histological manifestation. BAFF level in macrophage and joint homogenate elevated significantly in AA group, and the expression of BAFF in spleen and synovial tissue increased significantly by immunohistochemical analysis and quantitative real-time PCR. Correlation analysis showed joint histopathological manifestation significantly correlated

with the level of joint BAFF. The expression of total and activated CD4<sup>+</sup> T cells significantly decreased in blood and spleen in AA group compared with those in control group. The expression of CD21, CD138, and PNA<sub>d</sub> in synovial tissue and expression of CD3, CD20, and CD68 in spleen increased significantly in AA group. Immunofluorescent staining showed the number of CD68<sup>+</sup> macrophage coexpressed TACI, BCMA, BAFF-R in control and AA group. TACI, BCMA, BAFF-R mRNA expression in spleen increased apparently in AA group. The expression of TACI and BCMA in synovial tissue increased significantly, while the expression of BAFF-R decreased significantly in AA group. Subcutaneous administration of TACI-Ig significantly decreased BAFF level and attenuated progression of experimental arthritis, with reductions in inflammatory response and bone and joint destruction. **Conclusion:** Local production of BAFF may foster survival and/or expansion of B cells that produce pathogenic autoantibodies and/or promote local T cell activation and consequent joint destruction. Together, BAFF-receptors interactions are important not only for B cell function but for T cell-mediated immune responses.

**Keywords:** BAFF; receptor; arthritis; autoimmune; T cell

**Acknowledgements:** The authors deeply thank Wen-di ZHAO for her excellent assistance in histology and Li GUI for her essential contribution to the implementation of FACS. The study was supported by the National Natural Science Foundation of China (No 31200675 and No 81173075), by the Anhui Province Natural Science Foundation (No 1208085QH158 and No 11040606M195), by the Young Talents Foundation of the Education Department of Anhui Province (No 2010SQRL067), and the Financial Assistance Project of Scientific Research Funds of Doctor (XJ200808).

**S5.3****PPAR $\delta$  mediates the vascular benefits of metformin in diet-induced obese mice**

Wai San CHEANG<sup>1</sup>, Xiao Yu YIAN<sup>1</sup>, Wing Tak WONG<sup>2</sup>, Yu HUANG<sup>1</sup>. <sup>1</sup>Institute of Vascular Medicine and Li Ka Shing Institute of Health Sciences, The Chinese University of Hong Kong, Hong Kong, China; <sup>2</sup>Division of Cardiovascular Medicine, Department of Medicine, Stanford University School of Medicine, CA, USA

**Aim:** 5' Adenosine monophosphate-activated protein kinase (AMPK) was reported to form transcriptional complex with peroxisome proliferator-activated receptor delta (PPAR $\delta$ ) to induce gene expression synergistically. This study investigated whether PPAR $\delta$  mediates the effects of anti-diabetic drug metformin (AMPK activator) in ameliorating ER stress and endothelial dysfunction in high-fat diet-induced obese (DIO) mice. **Methods:** Endothelium-dependent relaxation (EDR) and protein expressions in aortae were measured by wire myograph and Western blotting, respectively. Fluorescence imaging determined the levels of reactive oxygen species (ROS) and nitric oxide (NO) under confocal microscopy. **Results:** DIO mice showed impaired EDR and elevated levels of ER stress markers and ROS in aortae. Chronic metformin treatment reversed the above-described effects in DIO PPAR $\delta$  wild-type littermates but not in knockout mice. Metformin and PPAR $\delta$  agonist GW1516 alleviated the tunicamycin (ER stress inducer)-induced impairment of EDR, ER stress and oxidative stress in mouse aortae as well as NO production in endothelial cells. Effects of metformin were abolished by the co-treatment of GSK0660 (PPAR $\delta$  antagonist) whilst those of GW1516 were not affected by compound C (AMPK inhibitor). **Conclusion:** The present study supports that metformin curtails ER stress and oxidative stress and increases NO bioavailability upon activation of AMPK/PPAR $\delta$  pathway, and subsequently combats against vasculopathy in obese diabetic mice.

**Keywords:** metformin; PPAR $\delta$ ; endothelial dysfunction; ER stress; obesity

**S5.4****Anti-hyperlipidemic effect of ZBH, a novel compound constructed by incorporating the pharmacophore of fibrates into resveratrol, in hyperlipidemic hamster**

Wei CHEN<sup>1</sup>, Shi-yong FAN<sup>1</sup>, Ni-na XUE<sup>1</sup>, Xin-ni XIE<sup>1</sup>, Xue-yuan JIN<sup>2</sup>, Li-li WANG<sup>1</sup>. <sup>1</sup>Beijing Institute of Pharmacology and Toxicology, Beijing 100850, China; <sup>2</sup>International Center for Liver Disease Treatment, Beijing 302 Hospital, Beijing 100039, China

**Aim:** Hyperlipidaemia, a prevalent disease, is a high risk factor of cardiovascular disease. However, effective and safe pharmacological option for its treatment remains limitation. In order to acquire more excellent anti-hyperlipidemic drug, a novel structural compound ZBH was constructed by incorporating the key pharmacophore of fibrates into natural scaffold of resveratrol. In present study, the hypolipidemic action of ZBH was systematically evaluated in hyperlipidemic hamster and its mechanism of action was further investigated. **Methods:** The action of ZBH at PPAR was characterized by using transactivation assay. It's binding to human PPAR $\alpha$  was corroborated by surface plasmon resonance (SPR) biosensor

technology. Hypolipidemic action of ZBH was evaluated in high fat diet induced hyperlipidemic hamsters after 5 weeks treatment. Tissue-based gene expression assays were performed after 3-d and 5-week ZBH treatment. **Results:** ZBH showed stronger PPAR $\alpha$  agonism *in vitro* relative to PPAR $\delta$  and PPAR $\gamma$ , its activation potency to PPAR $\alpha$  is 5.5 fold over than that of bezafibrate. Biacore SPR revealed that ZBH bind to the ligand binding domain (LBD) of PPAR $\alpha$  with high affinity. ZBH significantly lowered triglycerides, TC, FFA, proatherogenic lipoproteins LDL-C, hyperinsulinemia and improved insulin sensitivity, which is considerably stronger than bezafibrate. Gene expression disclosed that ZBH markedly decreased endogenous fatty acid synthesis in liver and increased fatty acid uptake and oxidation in liver and skeletal muscle, besides up-regulated SIRT1 expression. **Conclusion:** ZBH could significantly ameliorated dyslipidemia and insulin resistant of the hyperlipidemic hamsters by activating PPAR $\alpha$  through interactions with the LBD of the receptor and promoting the expression of SIRT1.

**Keywords:** PPAR $\alpha$ ; SIRT1; dyslipidemia; insulin resistance

### S5.5

#### Cucurbitacin B induced ATM-mediated DNA damage causes G<sub>2</sub>/M cell cycle arrest in a ROS-dependent manner

Jia-jie GUO, Guo-sheng WU, Jiao-lin BAO, Jin-jian LU, Xiu-ping CHEN\*. State Key Laboratory of Quality Research in Chinese Medicine, Institute of Chinese Medical Sciences, University of Macau, Macao, China

\*To whom correspondence should be addressed.

E-mail: chenxiu0725@yeah.net

**Aim:** Cucurbitacin B (Cuc B), a natural triterpenoid with potent anti-cancer activities both *in vitro* and *in vivo*, induces cancer cells cycle arrest at S or G<sub>2</sub>/M phases while the detailed mechanisms remain to be clear. This study was designed to precisely dissect mechanisms responsible for Cuc B-induced cell cycle arrest in human lung adenocarcinoma epithelial A549 cells. **Methods:** The effect of Cuc B on A549 cell cycle distribution was analyzed with flow cytometry. The effect of Cuc B on DNA damage was determined with Comet assay and protein expression of  $\gamma$ H<sub>2</sub>AX. The intracellular reactive oxygen species (ROS) was determined with flow cytometry using fluorescence probe DCFH<sub>2</sub>-DA. The role of ATM, Chk1 was evaluated with specific siRNA. Related protein expressions were determined with Western blotting. **Results:** Cuc B dramatically induced G<sub>2</sub>/M phase arrest. Cuc B treatment caused DNA damage without affecting the signal transducer and activator of transcription 3 (STAT3). Cuc B triggered ATM-activated Chk1-Cdc25C-Cdk1 pathway, which could be reversed by both ATM siRNA and Chk1siRNA treatment. Cuc B also triggered ATM-activated p53-14-3-3- $\sigma$  pathway, which could be reversed by ATM siRNA. Cuc B treatment increased intracellular ROS formation, which was inhibited by N-acetyl-L-cysteine (NAC) pretreatment. Furthermore, NAC pretreatment inhibited Cuc B induced DNA damage and G<sub>2</sub>/M phase arrest. **Conclusion:** Taken together, these results suggested that Cuc B induced DNA damage in A549 cells mediated by increasing intracellular ROS formation, which lead to G<sub>2</sub>/M cell phase arrest through ATM-activated Chk1-Cdc25C-Cdk1 and p53-14-3-3- $\sigma$  parallel branches. These observations provide novel mechanisms and potential targets for better understanding of the anti-cancer mechanisms of cucurbitacins.

**Keywords:** Cucurbitacin B; ROS; DNA damage; G<sub>2</sub>/M arrest

**Acknowledgements:** This study was supported by the Science and Technology Development Fund of Macau Special Administrative Region (045/2011/A) and the Research Fund of University of Macau (MYRG161(Y3-L2)-ICMS11-CXP and MRG008/CXP/2013/ICMS).

### S5.6

#### Novel monofunctional platinum (II) complex Mono-Pt induces apoptosis-independent autophagic cell death in human ovarian carcinoma cells distinct from cisplatin

Wen-jie GUO<sup>1, \*</sup>, Yang-miao ZHANG<sup>1, 2, \*</sup>, Li ZHANG<sup>3</sup>, Bin HUANG<sup>2</sup>, Fei-fei TAO<sup>1</sup>, Wei CHEN<sup>1</sup>, Zi-jian GUO<sup>2, \*</sup>, Qiang XU<sup>1, \*</sup>, Yang SUN<sup>1, \*</sup>. <sup>1</sup>State Key Laboratory of Pharmaceutical Biotechnology, School of Life Sciences, Nanjing University, Nanjing 210093, China; <sup>2</sup>State Key Laboratory of Coordination Chemistry, Coordination Chemistry Institute, Nanjing University, Nanjing 210093, China; <sup>3</sup>Key Laboratory of Nuclear Medicine, Ministry of Health, Jiangsu Key Laboratory of Molecular Nuclear Medicine, Jiangsu Institute of Nuclear Medicine, Wuxi 214063, China

\*These authors contributed equally to this work.

\*To whom correspondence should be addressed.

**Aim:** Failure to engage apoptosis appears to be a leading mechanism of resistance to traditional platinum drugs in patients with ovarian cancer. Therefore, an alternative strategy to induce cell death is needed for the chemotherapy of this

apoptosis-resistant cancer. Here we report that autophagic cell death, distinct from cisplatin-induced apoptosis, is triggered by a novel monofunctional platinum (II) complex named Mono-Pt in human ovarian carcinoma cells. **Methods:** Cytotoxic effects were assessed by MTT and typan blue assay; Cell cycle was detected with PI staining; Autophagic and apoptotic cell death were confirmed by MDC staining, Western blot and EM analysis. **Results:** Mono-Pt-induced cell death has the following features: cytoplasmic vacuolation, caspase-independent, no nuclear fragmentation or chromatin condensation, and no apoptotic bodies. These characteristics integrally indicated that Mono-Pt, rather than cisplatin, initiated a nonapoptotic cell death in caov-3 ovarian carcinoma cells. Furthermore, incubation of the cells with Mono-Pt but not with cisplatin produced an increasing punctate distribution of microtubule-associated protein 1 light chain 3 (LC3), and an increasing ratio of LC3-II to LC3-I. Mono-Pt also caused the formation of autophagic vacuoles as revealed by monodansylcadaverine staining and transmission electron microscopy. In addition, Mono-Pt-induced cell death was significantly inhibited by the knockdown of either Beclin-1 or ATG7 gene expression, or by autophagy inhibitors 3-methyladenine, chloroquine and bafilomycin A1. Moreover, the effect of Mono-Pt involved the AKT1-MTOR-RP s6KB1 pathway and ERK1/2 signaling, since the TOR inhibitor rapamycin increased, while the ERK1/2 inhibitor U0126 decreased Mono-Pt-induced autophagic cell death. **Conclusion:** Taken together, our results suggest that Mono-Pt exerts anticancer effect via autophagic cell death in apoptosis-resistant ovarian cancer. These findings lead to increased options for anticancer platinum drugs to induce cell death in cancer.

**Keywords:** autophagy; monofunctional platinum complex; cisplatin; AKT; ERK

### S5.7

#### Compound Astragalus and Salvia Miltiorrhiza Extract suppresses cell migration of keloid fibroblasts: involvement of MAPK signaling pathway

Shu-fang HE<sup>1, 2, \*</sup>, Yan YANG<sup>2, \*</sup>, Jie-mei JIANG, Sen YANG<sup>3</sup>, Xue-jun ZHANG<sup>3</sup>. <sup>1</sup>Department of Anesthesiology, the Second Affiliated Hospital of Anhui Medical University, Hefei 230032, China; <sup>2</sup>Department of Pharmacology and Institute of Natural Medicine, Anhui Medical University, Hefei 230032, China; <sup>3</sup>Institute of Dermatology and Department of Dermatology, Anhui Medical University, Hefei 230032, China

\*To whom correspondence should be addressed.

E-mail: yangyan276866@sohu.com

**Aim:** To investigate the effects of Compound *Astragalus* and *Salvia Miltiorrhiza* Extract (CASE) on cell migration of keloid fibroblasts (KFs) and its mechanisms involving mitogen-activated protein kinase (MAPK) pathways including extracellular signal-regulated kinase (ERK), c-Jun N-terminal kinase (JNK) and p38 kinase. **Methods:** KFs were cultured and then treated with transforming growth factor beta 1 (TGF- $\beta$ <sub>1</sub>) with or without CASE (7.5, 15, and 30  $\mu$ g/mL) treatment. *In vitro* scratch-wound assays were performed to observe cell migration; Phosphorylation of ERK, JNK, or p38 and their intracellular localization in KFs were detected by Western blot and immunofluorescence assay, respectively. **Results:** Cell migration induced by TGF- $\beta$ <sub>1</sub> in KFs was significantly inhibited by CASE in a dose-dependent manner. ERK, JNK and p38 were moderately phosphorylated and mainly localized in the cytoplasm in KFs under control conditions. TGF- $\beta$ <sub>1</sub> enhanced the phosphorylation of ERK, JNK and p38, and promoted their nuclear translocation and which were significantly suppressed by CASE treatment. **Conclusion:** CASE significantly suppressed cell migration induced by TGF- $\beta$ <sub>1</sub> in keloid fibroblasts indicating its anti-keloid effects, the mechanisms might involve the inhibition on MAPK signaling pathway.

**Keywords:** keloid; compound *Astragalus* and *Salvia miltiorrhiza* extract; cell migration; transforming growth factor beta 1; mitogen activated protein kinase

**Acknowledgements:** This study was financially supported by the Academic and Technique Foregoer Foundation of Anhui province, China (2012381).

### S5.8

**Synthesis and antibacterial activity evaluation of three novel biscoumarin derivatives** Zheng HOU<sup>1</sup>, Jing LI<sup>2</sup>, Bao-hui LIU<sup>1</sup>, Ying ZHOU<sup>1</sup>, Xiao-yan XUE<sup>1</sup>, Ming-kai LI<sup>1, \*</sup>, Xiao-xing LUO<sup>1, \*</sup>. <sup>1</sup>Department of Pharmacology, School of Pharmacy, the Fourth Military Medical University, Xi-an, China; <sup>2</sup>School of Chemistry and Chemical Engineering, Xi-an University of Arts and Sciences, Xi-an, China

\*To whom correspondence should be addressed.

**Aim:** The treatment failure of methicillin resistant *Staphylococcus aureus* (MRSA) infections and emergence of vancomycin-resistant *Staphylococcus aureus* (*S. aureus*) urgently requires developing new antimicrobials. **Methods:** Three new biscoumarin derivatives, 3,3'-(4-nitrobenzylidene)-bis-(4-hydroxycoumarin) (NBH), 3,3'-(4-methoxybenzylidene)-bis-(4-hydroxycoumarin) (MBH), and

3,3'-(4-chloromethylbenzylidene)-bis-(4-hydroxycoumarin) (CBH), were synthesized and characterized by FT-IR, <sup>1</sup>H NMR, HRMS and single crystal X-ray crystallography. The minimum inhibitory concentration and time-kill curves were observed for the three title compounds and antibiotics. **Results:** Our results showed that two classical asymmetrical intramolecular O-H...O hydrogen bonds were formed in the three compounds. The compound NBH exerted potent bactericidal effects against almost all *S aureus* tested, including LAC, a highly virulent and wide-spread clinical isolate responsible for the recent epidemic of MRSA infections. NBH exhibited almost completely growth inhibition on all *S aureus* tested at 128 µg/mL and partly bactericidal effect at 32 µg/mL. **Conclusion:** The formation of intramolecular hydrogen bonds was considered as an important factor in assisting the molecule to attain the correct configuration for biological activity. The different substituents in the phenyl ring of biscoumarins had different antibacterial activities against *S aureus*.

**Keywords:** biscoumarin; single crystal; *Staphylococcus aureus*; minimum inhibitory concentration

## S5.9

### Trp64Arg (rs4994) polymorphism of β3-adrenergic receptor gene is associated with hyperuricemia in a Chinese male population

Qiong HUANG, Liu-fu ZHANG, Yan CHENG, Wei WEI\*. *Institute of Clinical Pharmacology, Anhui Medical University, key Laboratory of Anti-inflammatory and Immune Medicine (Anhui Medical University), Ministry of Education, Hefei 230032, China*

\*To whom correspondence should be addressed.

E-mail: wwei@ahmu.edu.cn

**Aim:** β3-Adrenergic receptor (β3-AR) gene is associated with insulin resistance and may affect serum uric acid levels. Our aim was to determine the possible association between β3-AR gene Trp64Arg polymorphism (rs4994) and hyperuricemia in a Chinese male population. **Methods:** A total of 410 hyperuricemic and 420 normouricemic male subjects were genotyped in this study. The genotypic and allelic frequencies were compared between the two groups. Body mass index (BMI), waist to hip ratio (WHR), systolic blood pressure (SBP), diastolic blood pressure (DBP), serum uric acid, urea nitrogen, creatinine, triglyceride, total cholesterol, low-density lipoprotein-cholesterol (LDL-c), high-density lipoprotein-cholesterol (HDL-c) and fasting plasma glucose (FPG) were determined. **Results:** The frequencies of CC genotype and C allele for Trp64Arg polymorphism were higher in hyperuricemic group than in normouricemic group ( $P < 0.01$  and  $P < 0.05$ , respectively). In both hyperuricemic and normouricemic groups, subjects with mutated C allele of Trp64Arg polymorphism showed significantly higher average uric acid levels than TT genotype carriers ( $P < 0.01$  and  $P < 0.01$ , respectively). Univariate and multivariate logistic regression showed that carrier of mutated C allele of Trp64Arg polymorphism was significantly associated with hyperuricemia occurrence ( $P = 0.003$ , OR = 1.587, 95% CI 1.175–2.145 and  $P = 0.003$ , OR = 1.676, 95% CI 1.051–3.617). **Conclusion:** Trp64Arg polymorphism was associated with hyperuricemia in a Chinese male population and should be an independent risk factor for hyperuricemia.

**Keywords:** β3-adrenergic receptor; genetic polymorphism; hyperuricemia; serum uric acid; association analysis

**Acknowledgements:** This study was supported by the National Natural Science Foundation of China Grants 81202596 and 81202541, Specialized Research Fund for the Doctoral Program of Higher Education (20113420120006), Grants for Scientific Research of BSKY (XJ201021) and Young Top-notch Talent Support Programs from Anhui Medical University (2012), Research Foundation of Anhui Medical University (2011XKJ011), Anhui Province Natural Science Foundation in University (KJ2012Z158), Foundation for Key Teacher by Anhui Medical University (2013).

## S5.10

### Effect of NR2B antagonist against Aβ<sub>25-35</sub> induced acute synaptic plasticity disruption in rat hippocampal slices

Yan HUANG, Zeng-yao HU, Gang LIU, Wen-xia ZHOU\*, Yong-xiang ZHANG\*. *Beijing Institute of Pharmacology and Toxicology, Beijing 100850, China*

\*To whom correspondence should be addressed.

E-mail: zhouwx@bmi.ac.cn (Wen-xia ZHOU); zhangyx@bmi.ac.cn (Yong-xiang ZHANG).

**Aim:** Amyloid beta (Aβ)-induced synaptic dysfunction is thought to be an important event in early stage of Alzheimer's disease (AD). In this study, the effect of NR2B antagonist against Aβ<sub>25-35</sub> induced acute synaptic plasticity disruption was observed. **Methods:** Hippocampus slices from adult rats were used for LTP experiments. **Results:** The results shown that treating hippocampal slices with Aβ<sub>25-35</sub> or NMDA suppressed the induction of long-term potentiation (LTP) significantly. Slices

treated with Aβ<sub>25-35</sub> by incubation could be reserved by ifenprodil incubation. LTP deficits induced by NMDA could also be reserved by ifenprodil. It was interesting that ifenprodil perfusion could not prevent the disruption of LTP in slices treated with Aβ<sub>25-35</sub> by perfusion. Co-applying ifenprodil and NMDA has protective effect on LTP against Aβ<sub>25-35</sub> perfusion. It was well established that Aβ could increase glutamate release and Aβ incubation or application *in vivo* might induce a high level of glutamate in incubation fluids or Aβ treated tissue. So perfusion of Aβ and NMDA mimics the situation of Aβ incubation. As the NMDARs are mainly consisting of NR2B and NR2A in the hippocampal tissues of adult rats, co-applying ifenprodil with NMDA means mainly activating NR2A. **Conclusion:** Our results indicated that blocking NR2B and activating NR2A might be a potential treatment strategy for AD in early stages.

**Keywords:** amyloid beta; NR2A; NR2B; long-term potentiation; hippocampus

**Acknowledgements:** This work was supported by grants from National Natural Science Foundation of China (81202505 and 81100239); Chinese National Technology Major Project of New Drug Development (2012ZX09J12201-002 and BWS11J052).

## S5.11

### Angiotensin II type I receptor-mediated oxidative stress in rostral ventrolateral medulla underlies the elevated blood pressure after stroke

Faith CH LI, Alice YW CHANG. *Center for Translational Research in Biomedical Sciences, Kaohsiung Chang Gung Memorial Hospital, Kaohsiung 83301, Taiwan, China*

Stroke is a major health problem throughout the world. The elevated blood pressure (BP) after stroke is a common complication associated with poor outcome and is less understood. Since the crucial role of angiotensin II (Ang II) and rostral ventrolateral medulla (RVLM) in central regulation of BP is well-known, the present study delineated the role of Ang II type I receptor (AT1R) at RVLM in the elevation of BP after transient stroke. The stroked Sprague-Dawley rats received 2-h left middle cerebral artery occlusion (MCAO) operation and BP were measured by tail-cuff plethysmography method in conscious conditions before and 24-h after stroke. The levels of Ang II, AT1R mRNA or protein, superoxide production, and activities of NADPH oxidase (Nox) or mitochondrial respiratory chain in RVLM were determined by ELISA, RT-PCR or chemiluminescence method. The elevation of BP was observed at 24 h after MCAO, accompanied by an increase of Ang II and the mRNA or protein levels of AT1R in RVLM. Moreover, the elevated BP, increased superoxide levels, augmented Nox activity and decreased activities of mitochondrial respiratory chain in RVLM at 24 h after stroke were significantly reduced by bilateral microinjection of AT1R antagonist candesartan (5 nmol), Nox inhibitor apocynin (10 nmol) or mitochondrial superoxide anion scavenger mitoTEMPO (500 pmol) into RVLM immediately after MCAO. We concluded that the AT1R-mediated oxidative stress in the RVLM participates in the elevation of BP after stroke.

**Keywords:** MCAO; stroke; AT1R; RVLM; oxidative stress

**Acknowledgements:** This work is supported by research grants number CMRPG8B1241 to Alice YW CHANG from the Chang Gung Medical Foundation, China.

## S5.12

### Id proteins are critical downstream effectors of BMPR-II in pulmonary arterial smooth muscle cells

Xiao-hui LI<sup>1</sup>, Ying LI<sup>2</sup>, Yuan-jian LI<sup>1</sup>. <sup>1</sup>*Department of Pharmacology, School of Pharmaceutical Science, Central South University, Changsha 410078, China;* <sup>2</sup>*The Third Xiangya Hospital, Central South University, Changsha 410023, China*

**Aim:** Bone morphogenetic protein type II receptor (BMPR-II) mutations are responsible for over 70% of cases of heritable pulmonary arterial hypertension (PAH) and up to 15%–40% of sporadic idiopathic disease. Loss of BMP signalling promotes pulmonary vascular remodelling via modulation of pulmonary artery smooth muscle cell (PASMC) migration, differentiation and proliferation. Id proteins (Id1-4), are endogenous inhibitors of basic helix-loop-helix transcription factor binding to DNA, and are major downstream transcriptional targets of BMP signalling. However, the impact of BMPR-II mutation on the expression of the range of Id proteins and the contribution of individual Id proteins to abnormal PASMC function remains unclear. **Methods:** Human PASMCs were incubated with BMP4, 6, 9 or with a range of cytokines. Expression of Id proteins (Id1-4) was determined using real-time PCR and Western blotting. The responses in control cells were compared with PASMCs harboring BMPR-II mutations and cells in which BMPR-II was knocked down by siRNA transfection. Id3 expression in pulmonary vessel was also investigated in the lungs of wild type mice and BMPR-II mutant

mice. Furthermore, lentiviral over-expression of Id3 was employed to investigate the role of Id3 protein on cell cycle regulation and proliferation in PASCs. **Results:** BMP4 and BMP6, but not BMP9, induced mRNA expression of all four Id genes in PASCs, though Id1 and Id3 demonstrated higher levels of induction compared to Id2 and Id4. Other cytokines and growth factors tested had minimal effect on Id gene expression. The time course and dose responsiveness of Id1 and Id3 induction by BMPs was confirmed by immunoblotting. The BMP stimulated induction of Id1 and Id3 was markedly reduced in BMPR-II mutant PASCs and in control PASCs following siRNA silencing of BMPR-II. Pulmonary arteries in BMPR-II mutant mice demonstrated a lower expression of Id3 than in wild type mice. Lentiviral overexpression of Id3 reduced cell cycle progression and inhibited proliferation of PASCs. **Conclusion:** Id proteins, and particularly Id1 and Id3, are critical downstream effectors of BMP signaling in PASCs. Loss of BMPR-II function reduces the induction of Id genes in PASCs, and Id3 regulates the proliferation of PASCs via cell cycle inhibition.

**Keywords:** pulmonary hypertension; BMPR-II; Id proteins; PASCs; cell cycle

**Acknowledgements:** Thanks for funding support from National Natural Science Foundation of China (No. 81200035) and China Postdoc Foundation (No. 2012M521567).

### S5.13

#### The analgesic and anticancer effects of minimally toxic fraction from Liu-Shen-Wan and the underlying mechanisms

Xiao-jun LI<sup>1</sup>, Mei-mei JIA<sup>1</sup>, Yu-sang LI<sup>1</sup>, Yan-ling YANG<sup>2</sup>, Xian-qing MAO<sup>2</sup>, He-bin TANG<sup>1</sup>.  
<sup>1</sup>Department of Pharmacology, College of Pharmacy, South-Central University for Nationalities, Wuhan 430074, China; <sup>2</sup>Laboratory of Molecular and Cellular Oncology, Department of Oncology, Public Research Center for Health (CRP-Santé), 84, Val Fleuri, L-1526, Luxembourg

**Aim:** Liu-Shen-Wan (LSW), an ancient preparation used to treat localized infection with pain, was reported to possess anticancer activity. We obtained a LSW supernatant (LSWS) fraction from ultrasound-assisted ethanol extraction (yield 15.9%) which proved to be safer than LSW in terms of hepatotoxicity. The mechanism responsible for LSW's analgesic and anticancer activity was investigated. **Methods:** Involvement of substance P/neurokinin-1 receptor (NK-1R) in these effects was studied in mouse dorsal root ganglion (DRG) cells and HL-60 and HepG2 cancer cells. **Results:** The LSWS (1 and 10 µg/mL) exhibited a potent inhibitory effect on the bradykinin-evoked rapid release of substance P from DRG cells. At concentrations of 0.1 µg/mL and higher, the LSWS resulted in a significant increase in the percentage of apoptotic HL-60 and HepG2 cancer cells. The LSWS also exerted a concentration-related inhibitory effect on HepG2 cells. The LSWS significantly downregulated the NK-1R expression in both HepG2 and bradykinin-treated DRG cells. In addition, it suppressed the caspase-dependent apoptotic pathway, but induced mitochondria-mediated apoptosis in HepG2 cells by downregulating the Bcl-2/Bax expression ratio. **Conclusion:** The substance P/NK-1R system was partly responsible for the analgesic and anticancer activity of LSWS. Our findings will be useful for developing more effective and less toxic LSW preparations.

**Keywords:** Liu-Shen-Wan; anticancer; analgesic; substance P; neurokinin-1 receptor

**Acknowledgements:** This work was supported by National Natural Science Foundation of China 81101538 and Hubei Natural Science Foundation 2012FFC13501.

### S5.14

#### N8-regulated entry of CD8<sup>+</sup> T cells into the CNS is required for the initiation of experimental autoimmune encephalomyelitis

Qiong LUO<sup>1</sup>, Fang Y GONG<sup>1</sup>, Yang SUN<sup>1\*</sup>, Wen LIU<sup>1</sup>, Yue H KE<sup>2</sup>, Zi C HUA<sup>1</sup>, Qiang XU<sup>1\*</sup>.  
<sup>1</sup>State Key Laboratory of Pharmaceutical Biotechnology, School of Life Sciences, Nanjing University, Nanjing 210093, China; <sup>2</sup>Laboratory of Cell Signaling and Modeling Genetics, Institute of Molecular Pathology, Department of Basic Medical Sciences, School of Medicine, Zhejiang University, Hangzhou 310058, China

\*To whom correspondence should be addressed.

**Aim:** In contrast to T cell priming in the periphery, therapeutic strategies targeting the initiation step of T cell trafficking into the central nervous system (CNS) have not been extensively investigated. In this study, the effect of N8 on experimental autoimmune encephalomyelitis (EAE) and the unique mechanism were elucidated. **Methods:** C57BL/6 mice were immunized with MOG35-55. Mice were monitored for clinical severity of disease and histopathologic features in the CNS. Levels of cytokines in serum were measured by ELISA. Effects of N8 on expressions of chemokines and cytokines in the CNS were determined by quantitative

PCR. **Results:** N8-treated mice developed conventional T helper 1 (T<sub>H1</sub>) and T helper 17 (T<sub>H17</sub>) responses but were highly resistant to the induction of EAE. Treatment with N8 resulted in decreased CNS accumulation of lymphocytes and increased functional expression of the chemokine receptor CXCR7 on CD8<sup>+</sup> T cells. Moreover, adoptive transfer of T cells from 2D2-transgenic mice reconstituted EAE susceptibility in N8-treated mice, indicating that N8 only targets the initial migration of pioneer T cells. **Conclusion:** N8 almost completely abolished the development of EAE by blocking the initial infiltration of pioneer CD8<sup>+</sup> T cells into the uninfamed CNS. Taken together, these results support N8 as a potential lead compound for the treatment of relapsing-remitting multiple sclerosis.

**Keywords:** N8; pioneer CD8<sup>+</sup> T cells; CXCR7; EAE; multiple sclerosis

### S5.15

#### Pharmacological study of a novel urea transporter inhibitor PU-48 as a diuretic

Hui-wen REN, Fei LI, Bao-xue YANG. State Key Laboratory of Natural and Biomimetic Drugs, Department of Pharmacology, School of Basic Medical Sciences, Peking University, Beijing 100191, China

**Aim:** The purpose of this study is to determine the feasibility of urea transporters (UT) as novel diuretic drug targets and to discover urea transporter inhibitors with diuretic activity. **Methods:** *In vitro* erythrocyte lysis model was used to get the potential urea transporter inhibitors. Inhibition activity of hits on UT-A was assayed in an MDCK cell line that stably expresses rat UT-A1. Cytotoxicity was assayed with MDCK cell viability after incubation with compounds for 72 h by cell counting kit-8 (CCK-8) assay. Diuretic activity of the compounds was determined in rats. Urine output, blood and urinary osmolality and urea concentration were measured to evaluate the diuretic role of compounds. **Results:** Structure-activity analysis indicated that the most potent compound, thienoquinolin PU-48, reversibly inhibited UT-B-facilitated urea transport in human, rabbit, rat and mouse erythrocytes with IC<sub>50</sub> of 0.18, 0.68, 0.35, and 0.77 µmol/L, respectively. PU-48 did not affect urea transport in UT-B null mouse erythrocytes. PU-48 showed no significant cellular toxicity at concentrations up to 80 µmol/L. Subcutaneous delivery of PU-48 (at 3.12, 12.5, and 50 mg/kg) to rats produced an increase of urine output and a decrease of the urine urea concentration and osmolalities without electrolyte disturbance and liver or renal damages, which indicates that PU-48 has significant diuretic effect by urea-selective diuresis. Moreover, there was no significant difference in UT-A1-A3, UT-B, and AQP1-4 protein abundance in the medulla between PU-48 treated rats and control. **Conclusion:** These results indicate that PU-48 or its analogues might be developed as a new diuretic to increase renal fluid clearance in diseases associated with water retention without causing electrolyte imbalance, as well as a tool drug to establish 'chemical knockout' analysis of the physiological functions of UTs.

**Keywords:** diuretics; urea transporter; urine concentrating mechanism; drug discovery

**Acknowledgements:** This work was supported by National Natural Science Foundation of China grants 30870921, 31200869, 81261160507, and 81170632, Drug Discovery Program grant 2009ZX09301-010-30, Research Fund for the Doctoral Program of Higher Education 20100001110047, Program of Introducing Talents of Discipline to Universities, International Science & Technology Cooperation Program of China 2012DFA11070, and Beijing Natural Science Foundation grant 7102105.

### S5.16

#### Low expression of β-arrestin2 correlated with hepatocellular carcinoma progression

Wu-yi SUN, Shan-shan HU, Jing-jing WU, Sen ZHANG, Wei WEI\*. Institute of Clinical Pharmacology, Anhui Medical University, Key Laboratory of Anti-inflammatory and Immune Medicine (Anhui Medical University), Ministry of Education, Hefei 230032, China

\*To whom correspondence should be addressed.

**Aim:** β-Arrestins are ubiquitous cytosolic proteins which initially be regarded as potential characters in G protein-coupled receptors (GPCR) desensitization, sequestration, and internalization. Recently, it has been documented that β-arrestins play a vital role in many tumors growth and metastasis. The present study aimed to explore the function and prognostic role of β-arrestins in hepatocellular carcinoma (HCC). **Methods:** The expression of β-arrestin1 and β-arrestin2 in paraffin-embedded human liver tissues and in tumors from HCC patients who underwent surgery resection were detected by immunohistochemistry (IHC), qPCR and Western blot, respectively. The prognostic significance of β-arrestin2 was validated using the Kaplan-Meier survival estimates and the log-rank tests. The expression of β-arrestin1 and β-arrestin2 in stepwise metastatic human HCC cell

lines (HCCLM3, MHCC97H, MHCC97L, SMMC-7721, and HepG2) and normal liver cell line L-02 were measured by RT-PCR and Western blot analysis. pcDNA3 expression plasmids encoding HA-tagged  $\beta$ -arrestin2 was used to investigate the role of  $\beta$ -arrestin2 in the metastasis and invasion of HCC. **Results:** IHC analysis showed that expression of  $\beta$ -arrestin2 was higher in paraffin-embedded normal liver tissues than in HCC tumor tissues, but the expression of  $\beta$ -arrestin1 was not significantly changed. The similar results were presented in tumors from HCC patients who underwent surgery resection. Univariate and multivariate analysis revealed that  $\beta$ -arrestin2 was a significant predictor for overall survival. *In vitro*, the RT-PCR and Western blot analysis showed that  $\beta$ -arrestin2 was low expressed in the highly metastatic HCC cell lines and high expressed in normal liver cell line L02, but there was no difference of  $\beta$ -arrestin1 expression among human HCC cell lines and L02. Furthermore, HCCLM3 cells were transfected with plasmid pcDNA3/ $\beta$ -arrestin2-HA. Cell scratching and transwell experiments showed that overexpression of  $\beta$ -arrestin2 reduced the ability of cell migration and invasion. Moreover, overexpression of  $\beta$ -arrestin2 significantly attenuated the pAkt activation and enhanced the level of E-cadherin. However, ERK activation had no significant change. **Conclusion:**  $\beta$ -Arrestin2 upregulation inhibits HCC cell invasion. And  $\beta$ -arrestin2 might be a novel target in HCC diagnosis and therapy.

**Keywords:** hepatocellular carcinoma;  $\beta$ -arrestins; invasion; metastasis; prognosis

**Acknowledgements:** This work was supported by grants from Specialized Research Fund for the Doctoral Program of Higher Education of China (No 20113420120002), Natural Science Foundation of the Higher Education Institutions of Anhui Province (No KJ2012A153), the National Natural Science Foundation of China (30973543 and 81173075). The authors acknowledge the help of the staff members of the Institute of Clinical Pharmacology, Anhui Medical University in conducting the study.

#### S5.17

##### Vascular endothelial growth factor receptor 2 maintains resting cardiac vagal baroreflex and heart functions as revealed by $Kdr^{+/-}$ mice

Ching-Yi TSAI, Samuel HH CHAN. *Center for Translational Research in Biomedical Sciences, Kaohsiung Chang Gung Memorial Hospital, Kaohsiung 83301, Taiwan, China*

**Aim:** The nucleus tractus solitarius (NTS), the principal terminal site of the baroreceptor afferents, contributes to cardiac vagal baroreflex by its connection with the nucleus ambiguus (NA), the origin of the vagus nerve. We investigated the role of vascular endothelial growth factor receptor 2 (VEGFR2), also known as  $Kdr$ /Flk-1, in cardiac vagal baroreflex and heart function under resting conditions. **Methods:** Blood pressure and heart rate (HR) of heterozygous ( $Kdr^{+/-}$ ) and wild-type ( $Kdr^{+/+}$ ) mice were recorded under conscious state by radiotelemetry. Magnetic resonance imaging/diffusion tensor imaging (MRI/DTI) was performed using a 9.4 T Animal MR scanner. Echocardiography was obtained using a 40-MHz linear-array transducer. **Results:** Flk-1 mRNA and protein in the NTS and heart, power spectrum of HR, an indicator of overall functionality of brain stem cardiovascular regulation, were significantly reduced in  $Kdr^{+/-}$  mice. DTI of the brain stem revealed that the connectivity between the NTS and NA was reduced in  $Kdr^{+/-}$  mice, concurrent with a decrease in cardiac vagal baroreflex and an increase in HR. Echocardiographic analysis further showed that  $Kdr^{+/-}$  mice were inferior to  $Kdr^{+/+}$  mice in terms of ejection fraction, fractional shortening, circumferential strain, and radial strain.

**Conclusion:** We conclude that VEGFR2 expression in the NTS is required for the maintenance of cardiac vagal baroreflex manifested in the form of connectivity between the NTS and NA, and for sustained heart functions under resting conditions.

**Keywords:** VEGFR2; cardiac vagal baroreflex; MRI/DTI

#### S5.18

##### Nicotinamide phosphoribosyltransferase-mediated $NAD^+$ biosynthesis contributes to post-ischemic neovascularization: role of endothelial progenitors

Pei WANG, Chao-yu MIAO\*. *Department of Pharmacology, Second Military Medical University, Shanghai 200433, China*

\*To whom correspondence should be addressed.

Email: cymiao@smmu.edu.cn

**Aim:** The aim of the present study was to determine whether nicotinamide phosphoribosyltransferase (NAMPT)-mediated nicotinamide adenine dinucleotide ( $NAD^+$ ) biosynthesis is important for post-ischemic neovascularization. **Methods:** EPCs were isolated from bone-marrow (BM) for *in vitro* study. Two strains of transgenic mice overexpressing NAMPT (Tg mice) and H247A-mutant dominant negative NAMPT ( $\Delta$ Tg mice) were generated for *in vivo* and mechanism study. Diabetic *db/db* mice and patients with EPC dysfunction were administered with a substrate of NAMPT (vitamin B<sub>3</sub>) to test the potential translational therapeutic

value. **Results:** In mice, ischemia increased the circulating and BM  $NAD^+$  levels and EPCs number, which were abolished by  $NAD^+$  synthesis blockade with NAMPT inhibition. NAMPT inhibition also inhibited EPCs function *in vitro*. EPCs obtained from Tg mice, but not  $\Delta$ Tg mice, demonstrated enhanced proliferation, migration and tube formation. Furthermore, post-ischemic neovascularization after hind-limb or cerebral ischemia was improved in Tg mice but not in the  $\Delta$ Tg mice. NAMPT-overexpression increased the protein levels of endothelial nitric oxide synthase (eNOS) in EPCs in a SIRT1 deacetylase-dependent manner. Lastly, administration of vitamin B<sub>3</sub> increased the number of circulating EPCs in diabetic *db/db* mice as well as in patients, and accelerated wound healing in *db/db* mice. **Conclusion:** These findings suggest that NAMPT-mediated  $NAD^+$  metabolic pathway plays a critical role in biological function of EPCs and contributes to post-ischemic neovascularization.

**Keywords:** endothelial progenitor cells;  $NAD^+$ ; NAMPT; neovascularization; SIRT1

**Acknowledgements:** This work was supported by grants from the National Basic Research Program of China (2009CB521902), the National Natural Science Foundation of China (81100866, 81130061 and 30525045), the Program of Shanghai Subject Chief Scientist (10XD1405300), the Shanghai Municipal Education Commission (12ZZ078), the Shanghai 'Shu Guang' Project (10GG19) and the SMMU Young Investigator Foundation.

#### S5.19

##### Tumor necrosis factor- $\alpha$ regulates the signaling of prostaglandin E<sub>2</sub> receptor 4 through the interaction between TRAF2 and GRK2 in human fibroblast-like synoviocytes

Qing-tong WANG<sup>#</sup>, Bei HUANG<sup>#</sup>, Kang-kang LIU, Qiong HUANG, Wu-yi SUN, Cheng-yi WU, Shang-xue YAN, Ling-ling ZHANG, Jing-yu CHEN, Hua-xun WU, Ai-wu ZHOU, Yun-fang ZHANG, Li-hua LIU, Wei WEI\*. *Institute of Clinical Pharmacology, Anhui Medical University, Key Laboratory of Anti-inflammatory and Immune Medicine (Anhui Medical University), Ministry of Education, Hefei 230032, China*

<sup>#</sup>These authors contributed equally to this work.

\*To whom correspondence should be addressed.

E-mail: wwei@ahmu.edu.cn

**Aim:** In our previous work, we established that upon tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) stimulation, the prostaglandin E<sub>2</sub> (PGE<sub>2</sub>)-PGE<sub>2</sub> receptor (EP)-guanine nucleotide-binding protein (G protein)-adenylyl cyclase (AC)-cyclic adenosine monophosphate (cAMP) signal transduction pathway of synoviocytes in collagen-induced arthritis (CIA) rats was unbalanced. The present study was designed to explore how TNF- $\alpha$  regulated the signaling of EP4 receptor through its downstream molecules in human fibroblast-like synoviocytes (hFLS). **Methods:** TNF- $\alpha$ -induced endogenous PGE<sub>2</sub> production was inhibited by Indometacin (INN), a COX-2 selective inhibitor. The concentration of PGE<sub>2</sub> in culture supernatants was determined with a human-specific enzyme linked immunosorbent assay (ELISA) kit. Proliferation of hFLS was detected by thiazolyl blue tetrazolium bromide (MTT) assay. Cellular cAMP level was measured using <sup>125</sup>I-cAMP radioimmunoassay (RIA) kit. The levels of TNF receptor (TNFR) 1 and TNFR2 were quantified by flow cytometry. The expressions of total TNF receptor-associated factor 2 (TRAF2), stimulatory G protein (G $\alpha$ s), EP4, as well as the levels of EP4 and G protein coupled receptor kinase 2 (GRK2) in plasma membrane and cytoplasm were analyzed by Western blotting. The ability of TRAF2 to interact biochemically with GRK2 or  $\beta$ -arrestin2 and the level of activated G $\alpha$ s were determined by co-immunoprecipitation. The gene expression of TRAF2 was silenced by small interfering RNA (siRNA) transfection. **Results:** We were surprised to find that TRAF2 could interact with GRK2, which contributed to the regulation of EP4 signaling by TNF- $\alpha$ . Further studies revealed that silencing TRAF2 gene could significantly decrease the membrane translocation of GRK2 resulted in an up-regulated expression of membrane EP4 and intracellular cAMP. **Conclusion:** These results suggest a novel form of cross-talk between TNF- $\alpha$  and G protein coupled receptors (GPCRs) EP4. TNF- $\alpha$  promotes desensitization of EP4 by regulating the compound of TRAF2-GRK2; furthermore, TRAF2 could control the translocation of GRK2.

**Keywords:** tumor necrosis factor- $\alpha$ ; TNF receptor-associated factor 2; G protein coupled receptor kinase2; EP4; human fibroblast-like synoviocytes

**Acknowledgements:** This work was financially supported by the National Natural Science Foundation of China (30973543, 81173075, 81202541, and 81202596), the Anhui Provincial Natural Science Foundation (1208085QH146), the Anhui Province Natural Science Foundation in University (KJ2011Z181 and KJ2011Z180), the Specialized Research Fund for the Doctoral Program of Higher Education of China (20113420120002), Grants for Scientific Research of BSKY (XJ201213), and the Foundation for Key Teacher by Anhui Medical University (2013).

**S5.20****Proliferation effects of dorsolateral prostate induced by exposure to low-dose bisphenol A in adult rats**

Jian-hui WU, Han YAN, Xin SU, Qi PAN, Si-chong XU, Bo GUI, Zu-yue SUN. *National Evaluation Centre for the Toxicology of Fertility Regulating Drugs, Shanghai Institute of Planned Parenthood Research, Shanghai 200032, China*

**Aim:** Since low-dose bisphenol A (BPA) could aggravates symptom of benign prostatic hyperplasia (BPH) in rat BPH model, we hypothesized that it will has the same effect on prostate of adult rats, thus conducted present study and further investigated its underlying mechanisms. **Methods:** 50 adult male rats were divided into 5 groups at random, 10 per group, and were ig 10 mL/kg vehicle, 200 µg/kg Tamoxifen (TAM), 10.0, 30.0, and 90.0 µg/kg BPA for 4 weeks. Animal were terminated on the next day after the last treatment. Blood were collected to detect levels of testosterone (T), estradiol (E<sub>2</sub>), prolactin (PRL) and prostate specific antigen (PSA). The dorsolateral prostate (DLP) lobes were dissected and weighed, then fixed in formalin solution for histological examination, and further detected gene expression by microarray analysis and RT-PCR analysis. **Results:** 1) There were no significant differences in animal body weight between control and BPA groups, while it was lower in TAM group. 2) Between control and BPA groups, there were no differences in levels of E<sub>2</sub> and PRL, but T levels in BPA groups were decreased, and PSA levels in 10.0 µg/kg ( $P<0.05$ ) and 90.0 µg/kg BPA ( $P<0.01$ ) groups were increased. In TAM group, levels of E<sub>2</sub>, PRL ( $P<0.05$ ) and PSA ( $P<0.01$ ) were increased than that of control, but T level was lower than that of control. 3) Compared with control, total prostate weight, volume and prostate index in BPA groups were increased, and weight and index of DLP were increased than that of control ( $P<0.01$ ). While in TAM group, all indexes were decreased than that of control. 4) In BPA groups, the height of DLP epithelium were higher than that of control ( $P<0.01$ ), but it decreased with BPA dose increasing, while in TAM group, it was lower than that of control ( $P<0.01$ ). 5) Microarray analysis and RT-PCR analysis results showed that BPA induced up-regulation of *Pcna* and *Egf*, and down-regulated expression of *Cdh2* in DLP, especially down-regulated the gene *Cdh2* expression. **Conclusion:** Environment exposure to low dose of BPA may induce DLP of adult rats to proliferate through activating DNA replication and biosynthesis of steroids pathways.

**Keywords:** bisphenol A; proliferation; benign prostatic hyperplasia; genome oligo microarray analysis; RT-PCR; rat

**Acknowledgements:** This research was supported by grant 21007041 from National Natural Science Foundation of China and grant 2011ZX09301-005 from the Ministry of Science & Technology of China.

**S5.21****P-glycoprotein mediated efflux of aconitine *in vitro*, *in situ* and *in vitro***

Cui-ping YANG, Zheng LI, Tian-hong ZHANG, Fei LIU, Jing-lai LI, Xiao-ying WANG, Jin-xiu RUAN, Zhen-qing ZHANG\*. *Key Laboratory of Drug Metabolism and Pharmacokinetics, Institute of Pharmacology and Toxicology, Academy of Military Medical Sciences, Beijing100850, China*

\*To whom correspondence should be addressed.

**Aim:** Aconitine (AC) is a toxic alkaloid from some Chinese medicinal herbals with origin of genus *Aconitum*, which have been widely used for analgesic, cardiotoxic and anti-rheumatism treatment. To assess the mechanism of its low and variable oral bioavailability, the role of P-glycoprotein (P-gp) in AC absorption was evaluated in this study. **Methods:** The bidirectional transport of AC across Caco-2 and MDR1-MDCKII cells was investigated with or without P-gp inhibitors (verapamil and cyclosporin A). This was followed by an *in situ* single-pass intestinal perfusion study in rat to evaluate the effect of verapamil on the intestinal permeability of AC and determination of pharmacokinetic profile of AC following oral administration with or without pre-treatment of verapamil in rats. **Results:** The efflux of AC across Caco-2 and MDR1-MDCKII cells was greater than its influx and both P-gp inhibitors significantly decreased the efflux of AC. The *in situ* study indicated verapamil co-perfused with AC exhibited significant increase in intestinal permeability by 12.9-fold, from  $(0.22\pm 0.12)\times 10^{-5}$  cm/s to  $(2.85\pm 0.85)\times 10^{-5}$  cm/s. After oral dosing, the mean area under the plasma-concentration curve (AUC) of AC significantly increased from  $150.73\pm 29.62$  h ng/mL without verapamil to  $1002.75\pm 105.9$  h ng/mL with verapamil ( $P<0.001$ ). **Conclusion:** This work updates the involvement of P-gp in low oral bioavailability of AC and provides significant information for clinical application of aconite and their formulations containing AC.

**Keywords:** aconitine; P-glycoprotein; permeability; oralabsorption

**Acknowledgements:** This study was financially supported by the National Major Scientific and Technical Special Projects for Innovative Drug of China (No

2012ZX09301003-001-007).

**S5.22****Translational regulation of RPA2 via internal ribosomal entry site and by eIF3a**

Ji-ye YIN<sup>1,2</sup>, Zi-zheng DONG<sup>1</sup>, Ran-yi LIU<sup>1</sup>, Zhao-qian LIU<sup>2</sup>, Jian-ting ZHANG<sup>1</sup>. <sup>1</sup>Department of Pharmacology/Toxicology, IU Simon Cancer Center, Indiana University School of Medicine, Indianapolis, IN 46202, USA; <sup>2</sup>Institute of Clinical Pharmacology, Hunan Key Laboratory of Pharmacogenetics, Central South University, Changsha 410078, China.

**Aim:** RPA2 is a subunit of a trimeric RPA complex important for DNA repair and replication. Although it is known that RPA activity is regulated by post-translational modification, whether RPA expression is regulated and the mechanism therein is currently unknown. eIF3a, the largest subunit of eIF3, is an important player in translational control and has been suggested to regulate translation of a subset of mRNAs important for tumorigenesis, metastasis, cell cycle progression, drug response, and DNA repair. In the present study, we aimed to explore if RPA2 expression is regulated at translational level via internal ribosome entry site (IRES)-mediated initiation in response to DNA damage. **Methods:** Dual luciferase reporter assay, *in vitro* transcription and translation were used to identify and locate IRES element of RPA2. Realtime PCR and Western blotting were used to detect RPA2 mRNA and protein expression level, respectively. UV crosslinking and pull down assay was performed to explore the binding of eIF3a and RPA2 IRES. **Results:** In the present study, we show that RPA2 expression is regulated at translational level via internal ribosome entry site (IRES)-mediated initiation in response to DNA damage. We also found that eIF3a suppresses RPA2 synthesis and inhibits its cellular IRES activity by directly binding to the IRES element of RPA2 located at -50 to -150 bases upstream of the translation start site. **Conclusion:** RPA2 expression is translationally regulated via IRES and by eIF3a and that this regulation is partly accountable for cellular response to DNA damage and survival.

**Keywords:** eIF3a; replication protein A (RPA); cellular internal ribosome entry sites (IRES); DNA repair; DNA damage; translational control

**S5.23****Traditional Chinese medicine with reinforcing kidney activity ameliorate experimental autoimmune encephalomyelitis in rats by modulating neuroinflammatory and neurotrophic responses**

Lin-lin YIN<sup>1</sup>, Yong-yan CHEN<sup>2</sup>, Li-li LIN<sup>1</sup>, Qi ZHANG<sup>2</sup>, Lin LI<sup>1</sup>. <sup>1</sup>Department of Pharmacology, Xuan Wu Hospital of Capital Medical University, Beijing Geriatric Medical Research Center, Key Laboratory for Neurodegenerative Diseases of Ministry of Education, Beijing 100053, China; <sup>2</sup>Department of Clinical Pharmacology, School of Pharmaceutical Sciences, Zhengzhou University, Zhengzhou 450001, China

**Aim:** The present study was designed to determine whether traditional Chinese medicine with reinforcing kidney activity such as cornel iridoid glycoside (CIG) and epimedium flavonoids (EF) had effect on the development of experimental autoimmune encephalomyelitis (EAE) in rats and to elucidate its underlying mechanisms. **Methods:** EAE model was induced by immunization of adult female Lewis rats with purified guinea-pig myelin basic protein (MBP) 68-86. CIG and EF were administrated intragastrically once a day after immunization until d 21 post immunization (PI). Histopathological staining, enzyme-linked immunosorbent assay (ELISA), biochemical methods and Western blotting approaches were used to evaluate the disease incidence and severity, neuroinflammatory and neurotrophic response in the central nervous system (CNS). **Results:** CIG or EF intragastrical administration significantly reduced clinical score of neurological deficit in EAE rats; alleviated demyelination and inflammatory infiltration; and inhibited gliosis activation, nuclear transcription factor (NF-κB) expression in the spinal cord of EAE rats. Treatment with CIG or EF also enhanced neurotrophic factors such as nerve growth factor (NGF) or brain-derived nerve factor (BDNF) expression, increased the number of oligodendrocytes and protected the ultrastructure of myelin sheaths and axons in the spinal cord of EAE rats. **Conclusion:** Our results showed that CIG and EF could inhibit the development of MBP-induced EAE in rats and our findings suggest that traditional Chinese medicine with reinforcing kidney activity may be useful for the treatment of multiple sclerosis. This effect involved reducing neuroinflammation and enhancing myelination and neurotrophins.

**Keywords:** experimental autoimmune encephalomyelitis; cornel iridoid glycoside; epimedium flavonoids; neuroinflammation; neurotrophin

**S5.24****Berberine improves glucose consumption in cardiomyocytes via activation of 5'-adenosine monophosphate-activated protein kinase**

Ming ZHANG, Zhaojie MENG, Li CHEN\*. *Department of Pharmacology, Norman*

Bethune Medical College, Jilin University, Changchun 130021, China

\*To whom correspondence should be addressed.

**Aim:** Insulin resistance plays an important role in the pathogenesis of diabetic cardiomyopathy. Berberine (BER) is a plant alkaloid which promotes hypoglycemia via increasing insulin sensitivity in peripheral tissues. Little is known of BBR's role in regulating glucose metabolism in heart. **Methods:** We examined the effect and mechanism of BER on glucose consumption and glucose uptake in insulin sensitive or insulin resistant rat H9c2 cardiomyocyte cells. H9c2 myoblast cells were differentiated into cardiomyocytes and incubated with insulin for 24 h to induce insulin resistance. **Results:** BER-treatment of H9c2 cells increased glucose consumption and glucose uptake compared to controls. In addition, BER-treatment attenuated the reduction in glucose consumption and glucose uptake in insulin resistant H9c2 cells. Compound C, an inhibitor of AMP-activated protein kinase (AMPK), abolished the enhancement of glucose consumption and glucose uptake mediated by BBR in both insulin sensitive and insulin resistant H9c2 cells compared to controls. **Conclusion:** BER significantly increased AMPK activity, but had little effect on the activity of protein kinase B (AKT) in insulin resistant H9c2 cells, suggesting that berberine improves insulin resistance in H9c2 cardiomyocytes at least in part via stimulation of AMPK activity.

**Keywords:** berberine; glucose consumption; cardiomyocyte; AMPK

### S5.25

**Lycium barbarum polysaccharide LBPF4-OL, a new Toll like receptor 4-MD2 complex activator and inducer, can increase the p38 MAPK phosphorylation and decrease JNK and ERK1/2 phosphorylation**

Xiao-rui ZHANG, Wen-xia ZHOU\*, Yong-xiang ZHANG\*. *Institute of Pharmacology and Toxicology, Beijing 100850, China*

\*To whom correspondence should be addressed.

E-mail: zhouwx@bmi.ac.cn (Wen-xia ZHOU); zhangyx@bmi.ac.cn (Yong-xiang ZHANG)

**Aim:** LBPF4-OL is a polysaccharide from traditional Chinese medicine *Lycium barbarum*. In this research, we investigated the molecular mechanism of LBPF4-OL on macrophages. **Methods:** RT-PCR and ELISA were used in the analysis of cytokine level. Lymphocyte proliferation was analyzed with <sup>3</sup>H-TdR incorporation method. TLR4 gene mutation mice C3H/HeJ and antibodies block were used to verify the binding sites. Flow cytometry (FCM) and Western-blot (WB) were used to analysis TLR4 level and MAPK signal pathway, respectively. **Results:** We found that LBPF4-OL induced the TNF- $\alpha$  and IL-1 $\beta$  secretion in concentration dependent manner. Anti-TLR4 significantly inhibited LBPF4-OL induced TNF- $\alpha$  and IL-1 $\beta$  secretion. Anti-TLR2 (0.31, 0.62, 1.25, and 2.5  $\mu$ g/mL) concentration dependently inhibited IL-1 $\beta$  secretion, but only in the concentration of 0.62  $\mu$ g/mL inhibited TNF- $\alpha$  secretion. Anti-CR3 did not affect the TNF- $\alpha$  and IL-1 $\beta$  secretion. In C3H/HeJ mice, the results showed that LBPF4-OL significantly induced TNF- $\alpha$  and IL-1 $\beta$  secrete on C3H/HeJ mice macrophage, but had no influence on those of C3H/HeN mice. FCM results showed that LBPF4-OL significantly induced the expression of TLR4/MD2 both on peritoneal macrophage and on Raw264.7 cells. WB results showed that LBPF4-OL could strengthen p38 MAPK phosphorylation, but inhibited ERK1/2 and JNK phosphorylation. **Conclusion:** *Lycium barbarum* polysaccharide LBPF4-OL is a new Toll like receptor 4-MD2 complex activator and inducer, which can increase p38-MAPK and decrease JNK and ERK1/2 phosphorylation level.

**Keywords:** *Lycium barbarum* polysaccharide; TLR4/MD2; MAPK

**Acknowledgements:** This work was supported by funds from the National Natural Science Foundation of China (No 81102451).

### S5.26

**Upregulation of TRPM7 channels by angiotensin II triggers phenotypic switching of vascular smooth muscle cells of ascending aorta**

Zheng ZHANG<sup>1</sup>, Mi WANG<sup>2</sup>, Xiao-han FAN<sup>3</sup>, Jing-hui CHEN<sup>4</sup>, Yong-yuan GUAN<sup>3</sup>, Yong-bo TANG<sup>3</sup>. <sup>1</sup>Department of Pharmacology, School of Pharmaceutical Sciences, Central South University, Changsha 410087, China; <sup>2</sup>Department of Cardiology, The Second Xiangya Hospital, Central South University, Changsha 410011, China; <sup>3</sup>Department of Pharmacology, Zhongshan School of Medicine, Sun Yat-Sen University, Guangzhou 510080, China; <sup>4</sup>Department of Anesthesiology, Guangzhou Women and Children's Medical Center, Guangzhou 510623, China

**Aim:** The present study was designed to explore whether the Ca<sup>2+</sup>-permeable transient receptor potential melastatin 7 (TRPM7) channel is involved in Ang II-induced phenotype switching of ascending aortic VSMCs, and to dissect the

molecular mechanisms by which TRPM7 modulates VSMC phenotype.

**Methods:** Whole-cell and single-channel patch clamping was carried out to measure TRPM7 currents in mouse aortic smooth muscle cells. TRPM7-mediated Ca<sup>2+</sup> entry was assessed by Fura-2/AM loading. Cell cycle distribution was analyzed by flow cytometry. **Results:** As revealed by current recording, Ang II infusion increased TRPM7 whole-cell currents in ascending aortic VSMCs. The increase in TRPM7 currents was found to result from enhanced expression of TRPM7 protein, rather than elevated single channel activity (open probability and slope conductance) and/or reduced Mg<sup>2+</sup>-mediated channel block. Mechanistically, Ang II elevated TRPM7 expression via Ang II type 1 receptor-mediated ERK1/2 signaling. As indicated by the expression levels of VSMC differentiation marker genes, phenotypic switching of ascending aorta occurred during Ang II infusion. Meanwhile, ERK1/2-Elk-1 signaling pathway known to suppress VSMC differentiation was activated in Ang II-infused ascending aorta. Knockdown of TRPM7 with small interfering RNA (siRNA) established a causative role of TRPM7 in Ang II-induced phenotypic change and promotion of cell proliferation. Moreover, TRPM7 was shown to be required for Pyk2-ERK1/2-Elk-1 pathway activation by Ang II, which potentiated TRPM7 channel function and thus activated the Ca<sup>2+</sup>-sensitive kinase Pyk2. Finally, TRPM7 knockdown attenuated Ang II-induced displacement of myocardin from SM22 promoter, but the effects could be reversed by expression of constitutively active c-Src. **Conclusion:** Our data establish that upregulation of TRPM7 channels by Ang II contributes to the development of the proliferative phenotype of ascending aortic VSMCs, and TRPM7 channel suppresses VSMC gene expression via Ca<sup>2+</sup> influx-mediated activation of Pyk2-ERK1/2-Elk-1 pathway.

**Keywords:** angiotensin II; TRPM7; phenotypic modulation; smooth muscle cell; Ca<sup>2+</sup> signal

**Acknowledgements:** This work was supported by the National Natural Science Foundation of China (No 30900530), Science and Technology Planning Project of Guangdong Province China (No 2011B080701012), Doctoral Fund of Ministry of Education of China (for young teachers, No 20090171120055), and the Fundamental Research Funds for the Central Universities in China (No 11ykpy02).

### S5.27

**Effects of Jin Chai antiviral capsule in vivo approach of influenza virus infection through inhibition of virus-induced NF- $\kappa$ B activation**

Ju-ying ZHONG<sup>1</sup>, Xiao-lan CUI<sup>2</sup>, Yu-jing SHI<sup>2</sup>, Ying-jie GAO<sup>2</sup>, Hong-xin CAO<sup>3</sup>. <sup>1</sup>Experimental Research Center, China Academy of Chinese Medical Sciences, Beijing 100700, China; <sup>2</sup>Pharmacology Laboratory, Institute of Chinese Materia Medica, China Academy of Chinese Medical Sciences, Beijing 100700, China; <sup>3</sup>China Academy of Chinese Medical sciences, Beijing 100700, China

\*To whom correspondence should be addressed.

E-mail: caohx@mail.caoms.ac

**Aim:** This study is to investigate the treatment of Jin Chai antiviral capsule for influenza virus PR8/34 (H1N1) infection. **Methods:** The model of pneumonia was established by dropping influenza virus into the nose of normal mice, using specific IKK inhibitors at non-toxic concentrations. This study analyzed the expression of NF- $\kappa$ B virus load cytokines in mouse model at the 1st, 3th, 5th, and 7th d after affected. **Results:** 1) By immunohistochemical assay that NF- $\kappa$ B pathway are activated upon PR8/34 (H1N1). By real-time PCR assay that both IKK inhibitors at non-toxic concentrations and Jin Chai antiviral capsule dose groups lead to decrease in signaling virus load and reduced cytokine (ELISA). Jin Chai antiviral capsule can decrease the expression of NF- $\kappa$ B. IKK $\alpha$ , large dose groups are significantly decrease the expression of NF- $\kappa$ B. IKK $\alpha$  compared with model group at each time point ( $P < 0.05$ ,  $P < 0.01$ ). Middle dose groups are significantly decrease the expression of NF- $\kappa$ B. IKK $\alpha$  compared with model group at the 3th d and the 5th d ( $P < 0.05$ ), small dose groups are significantly decrease the expression of NF- $\kappa$ B. IKK $\alpha$  compared with model group at the 3th d ( $P < 0.05$ ). Specific IKK inhibitors used in non-toxic concentration, also were given the Jin Chai antiviral capsule (large dose groups), assay that there was no obvious difference between the IKK inhibitors group and IKK inhibitors+Jin Chai antiviral capsule group (virus load cytokines). **Conclusion:** Effects of Jin Chai antiviral capsule approach of influenza virus infection through inhibition of virus-induced NF- $\kappa$ B activation.

**Keywords:** Jin Chai antiviral capsule; influenza virus H1N1; NF- $\kappa$ B pathway

**Acknowledgements:** This study was supported by National Significant New Drugs Creation-research and Development of Jin Chai Antivirus Capsule (No 2009zx09301-005).