



# Serum calprotectin as promising diagnostic aid in predicting relapse in proteinase 3-antineutrophil cytoplasmatic antibodies-associated vasculitis

Lieve Van Hoovels<sup>1</sup>, Xavier Bossuyt<sup>2</sup>

<sup>1</sup>Department of Laboratory Medicine, OLV Hospital, Aalst, Belgium; <sup>2</sup>Department of Laboratory Medicine, University Hospitals Gasthuisberg, Leuven, Belgium

*Correspondence to:* Dr. Xavier Bossuyt. Laboratory Medicine, University Hospitals Gasthuisberg, Herestraat 49, Leuven 3000, Belgium. Email: Xavier.Bossuyt@uzleuven.be.

*Comment on:* Pepper RJ, Draibe JB, Caplin B, *et al.* Association of Serum Calprotectin (S100A8/A9) Level With Disease Relapse in Proteinase 3-Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. *Arthritis Rheumatol* 2017;69:185-93.

Received: 19 March 2017; Accepted: 30 March 2017; Published: 19 April 2017.

doi: 10.21037/jlpm.2017.03.04

**View this article at:** <http://dx.doi.org/10.21037/jlpm.2017.03.04>

The recently published article of Pepper and colleagues (1) focuses on the deficit in diagnostic aids for predicting recurrent disease in antineutrophil cytoplasmatic antibodies (ANCA)-associated vasculitis (AAV). Although the diagnostic value of ANCA in AAV is well established and undisputed, the clinical benefit of ANCA for disease monitoring seems to be limited to targeted determination in patients clinically suspected for relapse (2,3), in patients with renal disease (4), and in patients that received plasmapheresis for severe AAV (5).

New biomarkers for prediction of recurrent AAV disease remain of major interest. Relapses in AAV are common and can lead to irreversible organ damage, higher risk of infections and chronic comorbidities, secondary to vasculitis or its treatment (6-8). Patients who were diagnosed before development of advanced renal insufficiency have a better prognosis. Combining identification of the disease phenotype at risk for relapse with appropriate use of readily available validated biomarkers that can predict relapses should guide an optimally targeted maintenance therapy. Several studies have found that a diagnosis of granulomatosis with polyangiitis (GPA) [versus microscopic polyangiitis (MPA)], proteinase 3 (PR3)-ANCA [versus myeloperoxidase (MPO)-ANCA] and having had relapses in the past confer the highest risk for subsequent relapses (3).

New insights into the pathogenesis of AAV expand not only the therapeutic options but also the diagnostic tools to predict AAV flares. Despite the central role of ANCA antibodies in AAV pathogenesis, also other biomarkers

of the inflammatory process are drawing attention, like inflammatory cytokines (IL-1) (9), complement factors (C5a) (10) and the percentage of CD5+ B cells (11,12).

Calprotectin has gained widespread interest in studies of acute and chronic inflammation and associated diseases. Calprotectin is a heterodimeric complex of 2 S100 calcium-binding proteins, myeloid-related protein (MRP)-8 (S100A8) and MRP-14 (S100A9), expressed in granulocytes and monocytes. Calprotectin is an important proinflammatory factor of innate immunity acting as endogenous damage-associated molecule regulating myeloid cell function via toll-like receptor-4 activation (13). Its release at sites of inflammation makes calprotectin a potent acute-phase reactant, with normal serum levels in the range of 1–6 mg/L which can increase more than 100-fold with active inflammation (14).

Fecal calprotectin is a sensitive and specific marker of intestinal inflammation, and its analysis is universally accepted and applied for diagnosis and follow up of inflammatory bowel diseases (15,16).

Plasma calprotectin may also be a clinically useful biomarker in several other inflammatory rheumatic diseases, including Still's Disease, ankylosing spondylitis, psoriatic arthritis, systemic lupus erythematosus, primary Sjögren's syndrome and juvenile idiopathic arthritis (17). A recent systematic review by Abildtrup and colleagues, including 17 studies (1988–2013) with 1,065 patients enrolled, addresses calprotectin as an indicator of disease

activity in rheumatoid arthritis (RA). A high calprotectin concentration is found in synovial fluid obtained from the joints of RA patients. Because of its low molecular weight (36.5 kDa), calprotectin diffuses into the circulation, where it can be measured, keeping in mind the short half-life of 5 hours (17,18). Because serum calprotectin directly reflects inflammation in the synovium, serum calprotectin is a more sensitive biomarker of disease activity in RA than conventional inflammatory indices such as erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP). Patients with active RA have elevated blood calprotectin levels, which decrease significantly with effective therapy (18). For this reason, serum calprotectin has been proposed as a promising biomarker to predict response to biological therapy in active RA patients at baseline and could be used to monitor response to treatment (19). Furthermore, serum calprotectin levels were independently associated with radiographic progression in RA and high baseline calprotectin levels predict future erosive damage (18).

Relative limited data can be found on the diagnostic role of calprotectin in vasculitis. The study of Croce and colleagues, which was performed in an experimental murine model of vascular injury, demonstrates that serum calprotectin broadly regulates vascular inflammation and controls the inflammatory response at several levels, including regulation of macrophage and neutrophil accumulation, macrophage cytokine production and regulation of vascular smooth muscle cell proliferation (20). Earlier studies examining calprotectin in vascular inflammation found high serum levels in acute Kawasaki disease and the authors concluded that circulating calprotectin levels might be a predictor of the severity of vasculitis (21). Elevated serum levels of calprotectin were also observed in patients with Behçet's disease (BD), a form of systemic vasculitis. However, these serum levels did not correlate with CRP levels, ESR, white blood cell count or BDCAF score (BD current activity form) (22). In the study of Kawasaki and colleagues, serum calprotectin levels were associated with the severity of renal injury and endothelial cell dysfunction in Henoch-Schönlein purpura (HSP) patients (23).

When systemic vasculitis also involves the gastrointestinal system, as might be the case for Takayasu arteritis, HSP, AAV, polyarteritis nodosa and BD, high fecal calprotectin levels can be indicative for active gastrointestinal disease (24).

Earlier studies of Pepper and colleagues already reported elevated serum calprotectin levels in patients with active generalized AAV compared to patients in

remission and healthy controls. Persistently elevated calprotectin levels were also found in patients who went on to relapse (25), which is comparable to the non-responders in RA treatment (19).

In their most recent study, Pepper *et al.* (1) addressed the question whether calprotectin can be used as a biomarker to predict AAV relapse. For AAV patients who attained complete remission, defined by a Birmingham Vasculitis Activity Score for GPA (Wegener's Granulomatosis) (BVAS/WG) of 0 and the successful discontinuation of prednisone, serum calprotectin levels were measured at baseline and at months 1, 2 and 6 following treatment initiation. Patients were treated either with oral cyclophosphamide and glucocorticoids or with rituximab and glucocorticoids. Patients in the cyclophosphamide group were switched to maintenance therapy with azathioprine if complete remission was achieved between month 4 to 6. Not only the level of calprotectin, but also the relative changes in serum levels over time correlated with the duration of complete remission and time to relapse. A total of 144 patients, 93 with PR3-ANCA and 51 with MPO-ANCA, were included in the analysis. As only 6 MPO-ANCA patients relapsed, the analysis was focused on the PR3-ANCA cohort. A significantly higher risk of relapse was associated with an increase in calprotectin between baseline and month 2 and 6 for all patients, suggesting that an individual patient's level of calprotectin from the time of diagnosis stratifies the patient's relapse risk. Subgroup analysis demonstrated that patients treated with rituximab and with increased levels of serum calprotectin at baseline had the greatest risk for future relapse. This is in concordance with the previous observation that elevated serum calprotectin levels predict relapse in rituximab treated RA patients (19). Nevertheless, some patients treated with cyclophosphamide/azathioprine, relapsed despite decreases in serum calprotectin, which requires further investigation.

The results of Pepper *et al.* (1) on the role of calprotectin in predicting AAV relapse are promising and merit further investigation.

## Acknowledgments

*Funding:* None.

## Footnote

*Provenance and Peer Review:* This article was commissioned

and reviewed by the Section Editor Zaixing Yang (Department of Laboratory Medicine, Taizhou First People's Hospital, Taizhou, China).

*Conflicts of Interest:* Both authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/jlpm.2017.03.04>). The authors have no conflicts of interest to declare.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

1. Pepper RJ, Draibe JB, Caplin B, et al. Association of Serum Calprotectin (S100A8/A9) Level With Disease Relapse in Proteinase 3-Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. *Arthritis Rheumatol* 2017;69:185-93.
2. Verstockt B, Bossuyt X, Vanderschueren S, et al. There is no benefit in routinely monitoring ANCA titres in patients with granulomatosis with polyangiitis. *Clin Exp Rheumatol* 2015;33:S-72-6.
3. Specks U. Accurate relapse prediction in ANCA-associated vasculitis-the search for the Holy Grail. *J Am Soc Nephrol* 2015;26:505-7.
4. Kemna MJ, Damoiseaux J, Austen J, et al. ANCA as a predictor of relapse: useful in patients with renal involvement but not in patients with nonrenal disease. *J Am Soc Nephrol* 2015;26:537-42.
5. Avery TY, Bons J, van Paassen P, et al. Diagnostic ANCA algorithms in daily clinical practice: evidence, experience, and effectiveness. *Lupus* 2016;25:917-24.
6. Robson J, Doll H, Suppiah R, et al. Damage in the anca-associated vasculitides: long-term data from the European vasculitis study group (EUVAS) therapeutic trials. *Ann Rheum Dis* 2015;74:177-84.
7. Miloslavsky EM, Specks U, Merkel PA, et al. Outcomes of nonsevere relapses in antineutrophil cytoplasmic antibody-associated vasculitis treated with glucocorticoids. *Arthritis Rheumatol* 2015;67:1629-36.
8. Kronbichler A, Jayne DR, Mayer G. Frequency, risk factors and prophylaxis of infection in ANCA-associated vasculitis. *Eur J Clin Invest* 2015;45:346-68.
9. Millet A, Martin KR, Bonnefoy F, et al. Proteinase 3 on apoptotic cells disrupts immune silencing in autoimmune vasculitis. *J Clin Invest* 2015;125:4107-21.
10. Kallenberg CG, Heeringa P. Complement system activation in ANCA vasculitis: A translational success story? *Mol Immunol* 2015;68:53-6.
11. Unizony S, Lim N, Phippard DJ, et al. Peripheral CD5+ B cells in antineutrophil cytoplasmic antibody-associated vasculitis. *Arthritis Rheumatol* 2015;67:535-44.
12. Bunch DO, Mendoza CE, Aybar LT, et al. Gleaning relapse risk from B cell phenotype: decreased CD5+ B cells portend a shorter time to relapse after B cell depletion in patients with ANCA-associated vasculitis. *Ann Rheum Dis* 2015;74:1784-6.
13. Vogl T, Tenbrock K, Ludwig S, et al. Mrp8 and Mrp14 are endogenous activators of Toll-like receptor 4, promoting lethal, endotoxin-induced shock. *Nat Med* 2007;13:1042-9.
14. Ehrchen JM, Sunderkötter C, Foell D, et al. The endogenous Toll-like receptor 4 agonist S100A8/S100A9 (calprotectin) as innate amplifier of infection, autoimmunity, and cancer. *J Leukoc Biol* 2009;86:557-66.
15. van Rhee PF, Van de Vijver E, Fidler V. Faecal calprotectin for screening of patients with suspected inflammatory bowel disease: diagnostic meta-analysis. *BMJ* 2010;341:c3369.
16. Basso D, Zambon CF, Plebani M. Inflammatory bowel diseases: from pathogenesis to laboratory testing. *Clin Chem Lab Med* 2014;52:471-81.
17. Kopeć-Mędrek M, Widuchowska M, Kucharz EJ. Calprotectin in rheumatic diseases: a review. *Reumatologia* 2016;54:306-9.
18. Abildtrup M, Kingsley GH, Scott DL. Calprotectin as a biomarker for rheumatoid arthritis: a systematic review. *J Rheumatol* 2015;42:760-70.
19. Choi IY, Gerlag DM, Herenius MJ, et al. MRP8/14 serum levels as a strong predictor of response to biological treatments in patients with rheumatoid arthritis. *Ann Rheum Dis* 2015;74:499-505.
20. Croce K, Gao H, Wang Y, et al. Myeloid-related

protein-8/14 is critical for the biological response to vascular injury. *Circulation* 2009;120:427-36.

21. Hirono K, Foell D, Xing Y, et al. Expression of myeloid-related protein-8 and -14 in patients with acute Kawasaki disease. *J Am Coll Cardiol* 2006;48:1257-64.
22. Oktayoglu P, Mete N, Caglayan M, et al. Elevated serum levels of calprotectin (MRP8/MRP14) in patients with Behçet's disease and its association with disease activity and quality of life. *Scand J Clin Lab Invest* 2015;75:106-12.
23. Kawasaki Y, Ohara S, Abe Y, et al. The role of serum myeloid-related protein 8/14 complex in Henoch-Schönlein purpura nephritis. *Pediatr Nephrol* 2012;27:65-71.
24. Hatemi I, Hatemi G, Çelik AF. Systemic vasculitis and the gut. *Curr Opin Rheumatol* 2017;29:33-8.
25. Pepper RJ, Hamour S, Chavele KM, et al. Leukocyte and serum S100A8/S100A9 expression reflects disease activity in ANCA-associated vasculitis and glomerulonephritis. *Kidney Int* 2013;83:1150-8.

doi: 10.21037/jlpm.2017.03.04

**Cite this article as:** Van Hoovels L, Bossuyt X. Serum calprotectin as promising diagnostic aid in predicting relapse in proteinase 3- antineutrophil cytoplasmatic antibodies-associated vasculitis. *J Lab Precis Med* 2017;2:10.