



The metastatic niche in the liver: tilling the soil for pancreatic cancer progression

Alexander Hann¹, Bruno Sainz Jr^{2,3,4}, Patrick C. Hermann¹

¹Department of Internal Medicine I, Ulm University, Ulm, Germany; ²Department of Biochemistry, Autonomous University of Madrid (UAM), Madrid, Spain; ³Institute of Biomedical Research “Alberto Sols” (IIBM) CSIC-UAM, Madrid, Spain; ⁴Chronic Diseases and Cancer Area, Ramón and Cajal Institute of Health Research (IRYCIS), Madrid, Spain

Correspondence to: Dr. Patrick C. Hermann, MD, PhD. Department of Internal Medicine I, Ulm University, Albert-Einstein-Allee 23, 89081 Ulm, Germany. Email: patrick.hermann@uni-ulm.de.

Comment on: Grünwald B, Harant V, Schaten S, *et al.* Pancreatic Premalignant Lesions Secrete Tissue Inhibitor of Metalloproteinases-1, Which Activates Hepatic Stellate Cells Via CD63 Signaling to Create a Premetastatic Niche in the Liver. *Gastroenterology* 2016;151:1011–1024.e7.

Abstract: Despite new developments in adjuvant and palliative treatment, pancreatic ductal adenocarcinoma (PDAC) still has a very poor prognosis. One of the main factors that prevent surgical resection of the tumor in curative intent is metastasis at the time of diagnosis. This perspective article summarizes new insights on the influence of premalignant lesions and PDAC itself on the formation of the metastatic niche in the liver.

Keywords: Pancreatic cancer; cancer stem cells; metastasis

Submitted Jan 04, 2017. Accepted for publication Jan 17, 2017.

doi: 10.21037/tcr.2017.02.30

View this article at: <http://dx.doi.org/10.21037/tcr.2017.02.30>

Pancreatic ductal adenocarcinoma (PDAC), the fourth leading cause of cancer-related death, has a poor overall survival rate with less than 7% of patients surviving for longer than 5 years after diagnosis (1), largely due to the lack of effective screening methods and therapeutic options. In fact, compared to other cancer entities that have benefited from significant improvements in screening and therapy, PDAC is predicted to become the second most lethal cancer by 2030 (2).

Despite this rather dismal outlook for pancreatic cancer, the field of cancer therapy as a whole has been positively changing over the last 5 years (3). Almost 14 years after the introduction of Gemcitabine as the standard of care for the treatment of PDAC, new chemotherapeutic options now promise an increase in the overall survival of PDAC patients. New first line palliative therapies for stage III and IV cancer such as FOLFIRINOX (4) or gemcitabine combined with nab-paclitaxel (5) are already used in daily practice, and the number of second line palliative therapies, like OFF (6) or nanoliposomal irinotecan combined with 5FU (7), are increasing. Likewise, there is a shift in

intensifying adjuvant therapy by using Gemcitabine in combination with capecitabine (8), and the results of the recent APACT trial suggest a potential benefit of adding Nab-paclitaxel to Gemcitabine in the adjuvant setting (9).

While these data are quite promising, most patients still present with a metastatic disease at the time of diagnosis with little to no hope for cure. Early detection of the disease would undoubtedly increase the number of tumors diagnosed at an early stage, thus increasing the rate of resections in curative intent that currently can only be offered to a small minority of PDAC patients (10); however, thus far no screening tools or methods have been developed to achieve such a goal. Additionally, 63% of patients that receive a resection for curative intent will eventually develop metastatic disease, local recurrence, or both months or years following surgical intervention, and only 22% of patients survive without tumor recurrence after surgery with adjuvant therapy (11). Thus understanding the mechanisms that are involved in the metastatic process is pivotal in order to improve the survival of both patients undergoing surgical resection and patients with locally advanced disease.




Figure 1 Overview of selected key factors for forming the hepatic metastatic niche influenced by malignant precursor lesions (PanIN). PanIN, pancreatic intraepithelial neoplasia.

The liver is the main venous drainage point of the mesenteric organs and thus the most common site of pancreatic cancer metastasis. While its anatomical situation is important, many other factors also promote metastatic spread to the liver. For example, the properties of the metastatic cell(s) that promotes metastases and the factors that support the implantation of this cell(s) in the liver are likely equally as important. Regarding the latter, among the key and most prominent cellular players that promote the formation of the “metastatic niche” in the liver are immune cells, such as natural killer cells (NK), T-cells, Kupffer cells (tissue-resident macrophages), tissue infiltrating monocytes/macrophages and dendritic cells (12). Additionally non-immune cells such as stellate cells have also been shown to play an important role in the regulation of the tumor microenvironment, and in the maintenance of cancer stem cells (13). Soluble factors also promote the crosstalk between tumor cells and the surrounding tissue, and important secreted factors including transforming growth factor beta (TGF- β) and the chemokine stromal cell-derived factor-1 (SDF-1) have been shown to be important for cancer cell intravasation and migration to the liver. TGF- β can act on tumor cells as well as on stromal cells. In cancer cells, TGF- β has been shown to induce epithelial-to-mesenchymal transition (EMT), thus promoting invasion (14). The upregulated expression of SDF-1 in metastatic target organs such as the liver plays an importing role for the “homing” of metastatic tumor cells, mediated by the SDF-1 receptor CXCR4, which is expressed

on metastatic subpopulations of cancer (stem) cells (15).

Based on detailed analyses of soluble serum factors in cancer-bearing patients and experimental mice, a new signaling crosstalk has been proposed in recent years. Based on the fact that certain serum factors were shown to be altered in mice bearing malignant precursor lesions in the pancreas [called pancreatic intraepithelial neoplasia (PanIN)], the hypothesis of “crosstalk between the premalignant lesion and the future metastatic organ” was born (Figure 1). Two soluble serum factors have been identified to prepare the metastatic niche in the liver already during the early stages of PDAC development: cancer-cell-derived exosomes and tissue inhibitor of metalloproteinases-1 (TIMP1). Glypican-1 (GPC1) containing exosomes were identified in PanIN-bearing mice and in the serum of PDAC patients (16). Importantly, the level of these exosomes correlated with tumor burden before and after tumor resection, and the level of cancer-cell-derived exosomes correlated with the survival of these patients. In addition, pancreas-derived exosomes containing macrophage migration inhibitory factor (MIF) were shown to be taken up by Kupffer cells to stimulate hepatic stellate cells to produce fibronectin by secretion of TGF- β . These factors prepared the metastatic niche by arresting bone marrow-derived macrophages in the liver. Blocking of MIF-containing exosomes prevented niche formation and pancreatic cancer metastasis in mice (17).

The second factor that has been shown to prepare the hepatic metastatic niche during PDAC development is TIMP1. While this molecule was initially described as a protease inhibitor (18), studies have ascribed specific biological activities to TIMP1 that are independent of metalloproteinase activity. For example, TIMP1 was reported to promote cell growth in different cell lines (19). Interestingly, a study by Jung *et al.* showed that the main binding partner of TIMP1 on the cell surface is CD63 (20). In a study by Seubert *et al.*, increasing TIMP1 levels in mice led to an increase in tumor cell homing to the liver (21), and livers “primed” with high serum levels of TIMP1 were shown to produce SDF-1, fibronectin, and TGF- β . Furthermore, an enrichment of hepatic neutrophils was also observed. These neutrophils are recruited along the SDF-1/CXCR4 axis that is used by CXCR4-expressing tumor cells to invade the liver and other metastatic sites rich in SDF-1. As we have demonstrated before, this mechanism of metastasis can be abolished by blocking CXCR4, either using the small molecule AMD3100 or a specific inhibitory antibody (15). In a recent study Grünwald *et al.* shed more

light on how pancreatic tumors prepare the metastatic niche in the liver already in the early stages of tumor formation: TIMP1 activates hepatic stellate cells via CD63 binding, and upon this activation HSCs produce endogenous TIMP1 and SDF-1 (22). Again, neutrophils are attracted to the liver, which in addition to other cell types, help prepare the metastatic niche. Accordingly, serum TIMP1 levels were shown to be elevated in patients with PDAC, highlighting an association between TIMP1 and PDAC patient survival (23).

The question still remains, “Will these improvements in our understanding of the mechanisms promoting the formation of the (hepatic) metastatic niche lead to a therapeutic improvement of our present clinical scenarios?” More than 50% of PDAC patients are determined to be at the metastatic stage at the time of diagnosis, precluding them from curative surgery, and approximately 78% of the patients that initially present with resectable disease will develop metastasis, local recurrence of the tumor, or will succumb to the tumor during the following months (11), suggesting that metastatic spread already occurred before surgery. Only case series are available evaluating neoadjuvant chemotherapeutic intervention for locally advanced unresectable tumors with FOLFIRINOX or Gemcitabine combined with Nab-paclitaxel, and the results thus far are disappointing as only very few locally advanced tumors are down-graded sufficiently to convert them into surgically resectable (24).

There is no doubt that patients who present with a locally advanced unresectable tumor would indeed greatly benefit from therapies that protect against metastatic spread until neoadjuvant therapies can sufficiently reduce tumors to a size permitting surgical resection. So while we continue to develop methods and systems to improve early tumor detection, identify groups at high risk, and improve our ability to effectively reduce the primary tumor mass to allow for curative resection, we should simultaneously step up research efforts focused on metastasis prevention, which would complement these aforementioned key objectives and add a new armament to the arsenal of weapons available to fight pancreatic cancer.

Acknowledgments

Funding: PC Hermann is supported by a Max Eder Fellowship of the German Cancer Aid [111746], by the Hector Foundation Cancer Research Grant (M.65.1), and Baustein Grants of the University Ulm [PC Hermann (LSBN0097) and A Hann (LSBR.0035)]. B Sainz Jr is

supported by a Ramón y Cajal Merit Award from the Ministerio de Economía y Competitividad, Spain, a Clinic and Laboratory Integration Program (CLIP) grant from the Cancer Research Institute, NY, NY, USA (B Sainz Jr) and a Conquer Cancer Now Award from the Concern Foundation, LA, CA, USA (B Sainz Jr).

Footnote

Provenance and Peer Review: This article was commissioned and reviewed by the Section Editor Xiaotian Sun (The 307th Hospital of Chinese People’s Liberation Army, Beijing, China).

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/tcr.2017.02.30>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Kleeff J, Korc M, Apte M, et al. Pancreatic cancer. *Nat Rev Dis Primers* 2016;2:16022.
2. Rahib L, Smith BD, Aizenberg R, et al. Projecting cancer incidence and deaths to 2030: the unexpected burden of thyroid, liver, and pancreas cancers in the United States. *Cancer Res* 2014;74:2913-21.
3. Hann A, Bohle W, Egger J, et al. Improvement in advanced pancreatic cancer survival with novel chemotherapeutic strategies - experience of a community based hospital. *Z Gastroenterol* 2016;54:1138-42.
4. Conroy T, Desseigne F, Ychou M, et al. FOLFIRINOX versus gemcitabine for metastatic pancreatic cancer. *N*

- Engl J Med 2011;364:1817-25.
5. Von Hoff DD, Ervin T, Arena FP, et al. Increased survival in pancreatic cancer with nab-paclitaxel plus gemcitabine. *N Engl J Med* 2013;369:1691-703.
 6. Oettle H, Riess H, Stieler JM, et al. Second-line oxaliplatin, folinic acid, and fluorouracil versus folinic acid and fluorouracil alone for gemcitabine-refractory pancreatic cancer: outcomes from the CONKO-003 trial. *J Clin Oncol* 2014;32:2423-9.
 7. Wang-Gillam A, Li CP, Bodoky G, et al. Nanoliposomal irinotecan with fluorouracil and folinic acid in metastatic pancreatic cancer after previous gemcitabine-based therapy (NAPOLI-1): a global, randomised, open-label, phase 3 trial. *Lancet* 2016;387:545-57.
 8. Neoptolemos JP, Palmer D, Ghaneh P, et al. ESPAC-4: A multicenter, international, open-label randomized controlled phase III trial of adjuvant combination chemotherapy of gemcitabine (GEM) and capecitabine (CAP) versus monotherapy gemcitabine in patients with resected pancreatic ductal adenocarcinoma. *J Clin Oncol* 2016;34:abstr LBA4006.
 9. Tempero MA, Cardin DB, Goldstein D, et al. AFACT: Phase III randomized trial of adjuvant treatment with nab-paclitaxel (nab-P) plus gemcitabine (Gem) versus Gem alone in patients (pts) with resected pancreatic cancer (PC). *J Clin Oncol* 2016;34:abstr TPS473.
 10. Chari ST, Kelly K, Hollingsworth MA, et al. Early detection of sporadic pancreatic cancer: summative review. *Pancreas* 2015;44:693-712.
 11. Neoptolemos JP, Stocken DD, Bassi C, et al. Adjuvant chemotherapy with fluorouracil plus folinic acid vs gemcitabine following pancreatic cancer resection: a randomized controlled trial. *JAMA* 2010;304:1073-81.
 12. Canning C, O'Brien M, Hegarty J, et al. Liver immunity and tumour surveillance. *Immunol Lett* 2006;107:83-8.
 13. Lonardo E, Frias-Aldeguer J, Hermann PC, et al. Pancreatic stellate cells form a niche for cancer stem cells and promote their self-renewal and invasiveness. *Cell Cycle* 2012;11:1282-90.
 14. Chiang SP, Cabrera RM, Segall JE. Tumor cell intravasation. *Am J Physiol Cell Physiol* 2016;311:C1-C14.
 15. Hermann PC, Huber SL, Herrler T, et al. Distinct populations of cancer stem cells determine tumor growth and metastatic activity in human pancreatic cancer. *Cell Stem Cell* 2007;1:313-23.
 16. Melo SA, Luecke LB, Kahlert C, et al. Glypican-1 identifies cancer exosomes and detects early pancreatic cancer. *Nature* 2015;523:177-82.
 17. Costa-Silva B, Aiello NM, Ocean AJ, et al. Pancreatic cancer exosomes initiate pre-metastatic niche formation in the liver. *Nat Cell Biol* 2015;17:816-26.
 18. Brew K, Nagase H. The tissue inhibitors of metalloproteinases (TIMPs): an ancient family with structural and functional diversity. *Biochim Biophys Acta* 2010;1803:55-71.
 19. Hayakawa T, Yamashita K, Tanzawa K, et al. Growth-promoting activity of tissue inhibitor of metalloproteinases-1 (TIMP-1) for a wide range of cells. A possible new growth factor in serum. *FEBS Lett* 1992;298:29-32.
 20. Jung KK, Liu XW, Chirco R, et al. Identification of CD63 as a tissue inhibitor of metalloproteinase-1 interacting cell surface protein. *EMBO J* 2006;25:3934-42.
 21. Seubert B, Grünwald B, Kobuch J, et al. Tissue inhibitor of metalloproteinases (TIMP)-1 creates a premetastatic niche in the liver through SDF-1/CXCR4-dependent neutrophil recruitment in mice. *Hepatology* 2015;61:238-48.
 22. Grünwald B, Harant V, Schaten S, et al. Pancreatic Premalignant Lesions Secrete Tissue Inhibitor of Metalloproteinases-1, Which Activates Hepatic Stellate Cells Via CD63 Signaling to Create a Premetastatic Niche in the Liver. *Gastroenterology* 2016;151:1011-1024.e7
 23. Poruk KE, Firpo MA, Scaife CL, et al. Serum osteopontin and tissue inhibitor of metalloproteinase 1 as diagnostic and prognostic biomarkers for pancreatic adenocarcinoma. *Pancreas* 2013;42:193-7.
 24. Winner M, Goff SL, Chabot JA. Neoadjuvant therapy for non-metastatic pancreatic ductal adenocarcinoma. *Semin Oncol* 2015;42:86-97.

Cite this article as: Hann A, Sainz B Jr, Hermann PC. The metastatic niche in the liver: tilling the soil for pancreatic cancer progression. *Transl Cancer Res* 2017;6(Suppl 1):S217-S220. doi: 10.21037/tcr.2017.02.30