



# Diabetes burden in Asia: the role of metabolic surgery

Chih-Hung Lin<sup>1,2</sup>, Lee-Ming Chuang<sup>1,2</sup>

<sup>1</sup>Graduate Institute of Clinical Medicine, College of Medicine, National Taiwan University, Taipei 100, Taiwan; <sup>2</sup>Department of Internal Medicine, National Taiwan University Hospital, Taipei 100, Taiwan

*Contributions:* (I) Conception and design: All authors; (II) Administrative support: None; (III) Provision of study materials or patients: None; (IV) Collection and assembly of data: None; (V) Data analysis and interpretation: None; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

*Correspondence to:* Dr. Lee-Ming Chuang, MD, PhD. Department of Internal Medicine, National Taiwan University Hospital, 7, Chung Shan S. Rd, Taipei 100, Taiwan. Email: leeming@ntu.edu.tw.

**Abstract:** Type 2 diabetes mellitus (T2DM) is one of the largest epidemics the world has ever faced. As a result of rapid economic development and urbanization, Asia has the fastest-growing population affected by T2DM worldwide due to the shift to a sedentary lifestyle and over-nutrition status. The understandings of genetic predisposition have elucidated the potential connection between obesity and T2DM in the Asian populations. On the other hand, the Asian populations tend to have android adiposity, which may further contribute to the pathogenesis of T2DM. In this article, we'll discuss the potential role of metabolic surgery in the management of T2DM in the Asian populations.

**Keywords:** Type 2 diabetes mellitus (T2DM); Asia; genomics; visceral adiposity

Received: 05 December 2016; Accepted: 20 December 2016; Published: 11 February 2017.

doi: 10.21037/ales.2017.01.04

View this article at: <http://dx.doi.org/10.21037/ales.2017.01.04>

## Diabetes as a global epidemic

Diabetes mellitus (DM), characterized by clinical symptoms and signs of chronic hyperglycemia, is one of the largest and fastest-growing epidemics the world has ever faced. The non-communicable diseases (NCD) Risk Factor Collaboration stated that the number of adults with DM worldwide increased from 108 million in 1980 to 422 million in 2014 due to rise in prevalence, population growth and aging (1). According to the global estimate of International Diabetes Federation (IDF), there were 415 million people affected by DM in 2015, and the number will rise up to 642 million, which equals to a prevalence of 10% (2). DM can be categorized into several different types according to their distinctive pathophysiology. Among these various types, type 2 DM (T2DM) is the one mainly responsible for the current global epidemic, both in developed and developing countries (3). Being aware of the rapidly escalating menace of DM as a NCD, the United Nations (UN) General Assembly passed Resolution 61/225 in

2006, which stated that DM should be recognized as an international public health challenge and for each nation to target prevention and control of the emerging threat (4). In 2011, the UN General Assembly made a political declaration on the prevention and control of NCDs (5). In 2012, the World Health Organization (WHO) declared to reduce 25% of premature mortality from NCDs by 2025 (6). While the fight against T2DM has become a global issue, in this article we'll focus on the burden of T2DM and the role of metabolic surgery in Asia.

## Diabetes burden in Asia

Being the most populated area around the world, Asia has the fastest-growing population affected by T2DM, with many Asian countries suffer from folds of increase in disease prevalence (1,2). In 2015, the estimated population with DM in Asia was around 232 million (2). In China, the prevalence of T2DM has increased from 0.9% in 1980 to 10.6% in 2015 (2,7). In Malaysia, the prevalence of T2DM has increased from 2.1% in 1982 to 16.1% in 2015 (2,8,9).

In Indonesia, the prevalence of T2DM has increased from 1.6% in 1981 to 5.7% in 2015 (2,9,10). In India, an astounding high prevalence of T2DM (18.2%) was noticed in the urban area (11). It is generally believed that changes in lifestyle and dietary patterns due to rapid economic and social development are the major factors associated with the rapid growth of T2DM in Asia, especially in developing countries (12). Urbanization and westernized diet have led to sedentary lifestyle and over-nutrition- both are well-established risk factors for T2DM. The healthcare cost for T2DM increases as complications gradually occur or when hospitalization, surgery, or treatment of newer medications is required. However, despite of the rapid economic growth, wealth inequality is rather common in developing countries in Asia. In the lower economic groups, up to 34% of their income might have to be spent to cover costs for the care of diabetes (13,14). Together with scarcity of adequate health-care facilities and low educational status, these phenomena have undoubtedly lay obstacles in optimizing care for T2DM.

### Genetic predispositions

Genome-wide association studies (GWAS) have identified many susceptible genes for T2DM in Asian population, including *TCF7L2* (15,16), *PPARG* (17), *SLC30A8* (17), *HHEX* (17), *IGF2BP2* (17), *CDKN2A/CDKN2B* (17), *FTO* (17), *CDKAL1* (17,18), *KCNQ1* (19,20), and *KCNJ11* (21). Among these genes, *FTO* was found to be associated with the risk of T2DM and obesity in Asian populations. Genomic study performed in 3,041 patients with T2DM and 3,678 control subjects of Asian ancestry from Hong Kong and Korea identified the association between *FTO* and increased risk of T2DM. Furthermore, the A allele of rs8050136 at *FTO* was associated with increased body-mass index (BMI) (22). Chang *et al.* identified the rs9939609 A allele of *FTO* was strongly associated with obesity and BMI in the Chinese population, with each additional copy of the rs9936609 A allele was associated with a BMI increase of approximately 0.37 kg/m<sup>2</sup> (23). Further studies targeted on specific population groups also consistently confirmed the association of *FTO* with obesity-related traits in Chinese, Japanese, Koreans, Vietnamese, Filipino, Malay, and Asian Indian populations (24). In a meta-analysis that combined data of 96,551 individuals of Asian ancestry, each additional minor allele increases the risk of obesity by 1.25-fold, which is similar to the effect seen in European populations.

Despite the increase in BMI for each additional minor allele is less than in those with European ancestry, it might reflect the differences in adiposity phenotype in Asian populations (25).

### Visceral adiposity in Asian populations

Asians generally have a lower BMI than do people of many other races, with a different body composition. Abdominal obesity and adiposity is a characteristic feature in many Asian populations, especially in Southeast Asia (12). Compared to Caucasians with similar waist circumferences, Asians tend to have more visceral adipose tissue (VAT) (26). Excess VAT is known as an independent risk factor for T2DM, and is more strongly correlated to the disease states than subcutaneous adipose tissue (SAT) (27-29). Increased lipolysis originated from excess VAT leads to the elevation of non-esterified fatty acids and triglycerides in blood and skeletal muscle, which hinders muscular utilization of glucose. Excess VAT also increases insulin resistance by changing the secretion of cytokines, especially of leptin and adiponectin (30), and leads to pro-inflammatory conditions (31).

### Effects of metabolic surgery on adiposity

Most studies which focused on the change of fat mass deposit found significant reductions in both VAT and SAT within the first few months after metabolic surgery. However, studies with longer post-operation follow-up durations revealed further declines in VAT instead of SAT (32). In patients who received Roux-en-Y gastric bypass, Faria *et al.* identified that the reduction of VAT served as an indicator of the remission of metabolic syndrome 1 year after the procedure, even with identical BMI and SAT area (33). Studies also demonstrated reduction of plasma leptin and elevation of plasma adiponectin levels after metabolic surgery (32), which might suggest potential improvement to the obesity-related low-grade inflammation and the associated metabolic disorders. These lines of evidence might suggest a specific role of metabolic surgery in the management of T2DM in obese Asian patients. Indeed, the IDF statement declares that metabolic surgery is an appropriate management for obese T2DM patients who fail to achieve treatment targets with medical therapies, and the BMI cut-off point for intervention should be lower in Asians populations compared to other races (*Table 1*) (34).

**Table 1** Eligibility and prioritization for metabolic surgery in Asian populations not achieving optimal T2DM control with medical therapy according to the IDF statement

BMI range (kg/m <sup>2</sup> )	Eligible for surgery	Prioritized for surgery
<27.5	No	No
27.5–32.5	Yes–conditional*	No
32.5–37.5	Yes	Yes–conditional*
>37.5	Yes	Yes

\*, HbA<sub>1c</sub> >7.5% despite fully optimized conventional therapy, especially if weight is increasing, or other obesity-related co-morbidities, e.g., hypertension and sleep apnea, are not achieving targets on conventional therapies. Adapted from Dixon *et al.* Bariatric surgery: an IDF statement for obese type 2 diabetes (34). T2DM, type 2 diabetes mellitus; IDF, International Diabetes Federation; BMI, body-mass index.

## Conclusions

Rapid economic development and urbanization have resulted in a tremendous increase in the prevalence of T2DM in Asia over the past few decades. The genetic predisposition and distinct body composition create a unique association among obesity, adiposity and the pathogenesis of T2DM in the Asian populations. With the effects on body weight reduction as well as change of the configuration of adipose tissue, the role of metabolic surgery in the management of T2DM in the Asian populations warrants further investigation.

## Acknowledgments

*Funding:* None.

## Footnote

*Provenance and Peer Review:* This article was commissioned by the Guest Editor (Wei-Jei Lee) for the series “Laparoscopic Metabolic Surgery for the Treatment of Type 2 Diabetes in Asia” published in *Annals of Laparoscopic and Endoscopic Surgery*. The article has undergone external peer review.

*Conflicts of Interest:* Both authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/ales.2017.01.04>). The series “Laparoscopic Metabolic Surgery for the Treatment of Type 2 Diabetes in Asia” was commissioned by the editorial office without any

funding or sponsorship. The authors have no other conflicts of interest to declare.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

1. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in diabetes since 1980: a pooled analysis of 751 population-based studies with 4.4 million participants. *Lancet* 2016;387:1513–30.
2. International Diabetes Federation. IDF Diabetes Atlas, 7th edn. Brussels, Belgium: International Diabetes Federation, 2015.
3. Zimmet PZ, Magliano DJ, Herman WH, et al. Diabetes: a 21st century challenge. *Lancet Diabetes Endocrinol* 2014;2:56–64.
4. International Diabetes Federation. United Nations resolution 61/225: world diabetes day [2006]. Available online: <https://www.idf.org/sites/default/files/UN%20Resolution%20on%20World%20Diabetes%20Day%20of%20Dec%202006.pdf>
5. United Nations General Assembly. Political declaration of the high-level meeting of the General Assembly on the Prevention and Control of Non-Communicable Diseases [2011]. Available online: [http://www.un.org/ga/search/view\\_doc.asp?symbol=A/66/L.1](http://www.un.org/ga/search/view_doc.asp?symbol=A/66/L.1)
6. World Health Organisation. 65th World Health Assembly closes with new global health measures [2012]. Available online: [http://www.who.int/mediacentre/news/releases/2012/wha65\\_closes\\_20120526/en/index.html](http://www.who.int/mediacentre/news/releases/2012/wha65_closes_20120526/en/index.html)
7. Yin J, Kong AP, Chan JC. Prevention and Care Programs Addressing the Growing Prevalence of Diabetes in China. *Curr Diab Rep* 2016;16:130.
8. Zaini A. Where is Malaysia in the midst of the Asian

- epidemic of diabetes mellitus? *Diabetes Res Clin Pract* 2000;50 Suppl 2:S23-8.
9. Ramachandran A, Snehalatha C, Shetty AS, et al. Trends in prevalence of diabetes in Asian countries. *World J Diabetes* 2012;3:110-7.
  10. Waspadji S, Ranakusuma AB, Suyono S, et al. Diabetes mellitus in an urban population in Jakarta, Indonesia. *Tohoku J Exp Med* 1983;141 Suppl:219-28.
  11. Ramachandran A, Mary S, Yamuna A, et al. High prevalence of diabetes and cardiovascular risk factors associated with urbanization in India. *Diabetes Care* 2008;31:893-8.
  12. Ramachandran A, Ma RC, Snehalatha C. Diabetes in Asia. *Lancet* 2010;375:408-18.
  13. Yach D, Stuckler D, Brownell KD. Epidemiologic and economic consequences of the global epidemics of obesity and diabetes. *Nat Med* 2006;12:62-6.
  14. Ramachandran A, Ramachandran S, Snehalatha C, et al. Increasing expenditure on health care incurred by diabetic subjects in a developing country: a study from India. *Diabetes Care* 2007;30:252-6.
  15. Grant SF, Thorleifsson G, Reynisdottir I, et al. Variant of transcription factor 7-like 2 (TCF7L2) gene confers risk of type 2 diabetes. *Nat Genet* 2006;38:320-3.
  16. Cauchi S, El Achhab Y, Choquet H, et al. TCF7L2 is reproducibly associated with type 2 diabetes in various ethnic groups: a global meta-analysis. *J Mol Med (Berl)* 2007;85:777-82.
  17. Frayling TM. Genome-wide association studies provide new insights into type 2 diabetes aetiology. *Nat Rev Genet* 2007;8:657-62.
  18. Steinthorsdottir V, Thorleifsson G, Reynisdottir I, et al. A variant in CDKAL1 influences insulin response and risk of type 2 diabetes. *Nat Genet* 2007;39:770-5.
  19. Unoki H, Takahashi A, Kawaguchi T, et al. SNPs in KCNQ1 are associated with susceptibility to type 2 diabetes in East Asian and European populations. *Nat Genet* 2008;40:1098-102.
  20. Yasuda K, Miyake K, Horikawa Y, et al. Variants in KCNQ1 are associated with susceptibility to type 2 diabetes mellitus. *Nat Genet* 2008;40:1092-7.
  21. Florez JC. Clinical review: the genetics of type 2 diabetes: a realistic appraisal in 2008. *J Clin Endocrinol Metab* 2008;93:4633-42.
  22. Ng MC, Park KS, Oh B, et al. Implication of genetic variants near TCF7L2, SLC30A8, HHEX, CDKAL1, CDKN2A/B, IGF2BP2, and FTO in type 2 diabetes and obesity in 6,719 Asians. *Diabetes* 2008;57:2226-33.
  23. Chang YC, Liu PH, Lee WJ, et al. Common variation in the fat mass and obesity-associated (FTO) gene confers risk of obesity and modulates BMI in the Chinese population. *Diabetes* 2008;57:2245-52.
  24. Loos RJ, Yeo GS. The bigger picture of FTO: the first GWAS-identified obesity gene. *Nat Rev Endocrinol* 2014;10:51-61.
  25. Li H, Kilpeläinen TO, Liu C, et al. Association of genetic variation in FTO with risk of obesity and type 2 diabetes with data from 96,551 East and South Asians. *Diabetologia* 2012;55:981-95.
  26. Lear SA, Humphries KH, Kohli S, et al. Visceral adipose tissue accumulation differs according to ethnic background: results of the Multicultural Community Health Assessment Trial (M-CHAT). *Am J Clin Nutr* 2007;86:353-9.
  27. Boyko EJ, Fujimoto WY, Leonetti DL, et al. Visceral adiposity and risk of type 2 diabetes: a prospective study among Japanese Americans. *Diabetes Care* 2000;23:465-71.
  28. Gallagher D, Kelley DE, Yim JE, et al. Adipose tissue distribution is different in type 2 diabetes. *Am J Clin Nutr* 2009;89:807-14.
  29. Liu J, Fox CS, Hickson DA, et al. Impact of abdominal visceral and subcutaneous adipose tissue on cardiometabolic risk factors: the Jackson Heart Study. *J Clin Endocrinol Metab* 2010;95:5419-26.
  30. Havel PJ. Control of energy homeostasis and insulin action by adipocyte hormones: leptin, acylation stimulating protein, and adiponectin. *Curr Opin Lipidol* 2002;13:51-9.
  31. Leinonen E, Hurt-Camejo E, Wiklund O, et al. Insulin resistance and adiposity correlate with acute-phase reaction and soluble cell adhesion molecules in type 2 diabetes. *Atherosclerosis* 2003;166:387-94.
  32. Frikke-Schmidt H, O'Rourke RW, Lumeng CN, et al. Does bariatric surgery improve adipose tissue function? *Obes Rev* 2016;17:795-809.
  33. Faria G, Pestana D, Aral M, et al. Metabolic score: insights on the development and prediction of remission of metabolic syndrome after gastric bypass. *Ann Surg* 2014;260:279-86.
  34. Dixon JB, Zimmet P, Alberti KG, et al. Bariatric surgery: an IDF statement for obese Type 2 diabetes. *Diabet Med* 2011;28:628-42.

doi: 10.21037/ales.2017.01.04

**Cite this article as:** Lin CH, Chuang LM. Diabetes burden in Asia: the role of metabolic surgery. *Ann Laparosc Endosc Surg* 2017;2:21.