



# Feasibility, safety and limits of robotic approach for lymphadenectomy in biliary tumours

Julie Deyrat<sup>^</sup>, Stylianos Tzedakis

AP-HP, Centre Université de Paris, Groupe Hospitalier Cochin Port Royal, DMU Cancérologie et spécialités médico-chirurgicales, Service de chirurgie hépatobiliaire, digestive et endocrinienne, Paris, France

*Correspondence to:* Stylianos Tzedakis, MD. Department of Hepatobiliary, Digestive and Endocrine Surgery, Cochin - Port-Royal, AP-HP, Centre Université de Paris, 75005 Paris, France. Email: stylianos.tzedakis@aphp.fr.

*Comment on:* Ratti F, Cipriani F, Ingallinella S, *et al.* Robotic Approach for Lymphadenectomy in Biliary Tumours: The Missing Ring Between the Benefits of Laparoscopic and Reproducibility of Open Approach? *Ann Surg* 2022. [Epub ahead of print]. doi: 10.1097/SLA.0000000000005748.

**Keywords:** Robotic; minimally invasive; lymphadenectomy; biliary tumour

Submitted Mar 23, 2023. Accepted for publication Apr 26, 2023. Published online May 22, 2023.

doi: 10.21037/hbsn-23-149

**View this article at:** <https://dx.doi.org/10.21037/hbsn-23-149>

We read with great interest the study of Ratti *et al.* (1) aiming to compare robotic, laparoscopic and open hepatectomies in terms of feasibility, safety and oncological adequacy in surgery for biliary tumours (intrahepatic cholangiocarcinoma, perihilar cholangiocarcinoma and gallbladder cancer). The authors conducted a retrospective monocentric study focusing on the quality of lymphadenectomy, operative time and postoperative morbidity and mortality up to day 90. Using a weighted propensity score analysis by inverse probability of treatment weighting (IPTW), 25 patients operated by robotic approach were compared with 97 patients operated by laparoscopy and 113 by laparotomy. The authors showed a significant advantage of the robotic approach over laparoscopy and laparotomy respectively concerning the rate of adequate lymphadenectomy with a retrieval of at least 6 lymph nodes (96% *vs.* 86.6%;  $P=0.043$  and 76.9%;  $P=0.032$ ). Operative time for lymphadenectomy (35 *vs.* 50 minutes;  $P=0.044$ ) and conversion rate (4% *vs.* 8.2%;  $P=0.043$ ) were significantly reduced with the robotic approach when compared with laparoscopy. Minimal invasive surgery (robotic and laparoscopic approaches) was associated with a reduction in severe morbidity, length of hospital stays and length of recovery compared with open procedures. There was no significant difference in overall

and disease-free survival at one year follow-up between the three procedures. Moreover, duration of lymph node dissection, conversion rate and blood loss between the 25 first hepatectomies did not show any learning curve effect in the robotic group whereas a learning curve of 10 cases was identified in the laparoscopic group. The authors concluded that robotic approach granted better results than laparoscopic and open resection for cholangiocarcinoma surgery, but these results were also probably associated with their already advanced laparoscopic experience.

In this retrospective study, selection bias was limited by a propensity score analysis allowing the three groups (robotic, laparoscopic and open resection) to be comparable on prognostic and confusion factors. The study asserts a benefit of minimal invasive surgery for reduction of blood loss and improved post operative outcomes which is in accordance with the existing literature mostly based on retrospective studies and no randomised trial (2). Although the authors suggest a superior quality and duration of lymph node dissection with minimal invasive surgery compared with open surgery and with robotic approach over laparoscopy, this superiority is not observed for the total number of lymph node retrieved and the total operative time which is overall longer (290 *vs.* 260 minutes;

<sup>^</sup> ORCID: 0009-0006-4679-5234.

P=0.176) with the robotic approach. Laparoscopic hepatic pedicle lymphadenectomy is still under debate and was probably not systematically performed before the last American Joint Committee on Cancer (AJCC) recommendations (3,4). Indeed, a French national multicentric study recently reported worse results in laparoscopic lymph node resection quality when compared with open resection (5). However, these results were based on liver resections realised between 2000 and 2012 and therefore, before the publication of the latest AJCC recommendations (6) in which the quality of lymphadenectomy had been clearly established (>6 nodes removed to obtain a correct lymph node staging) and widely spread.

In this study, the authors do not describe any learning curve in terms of time required for lymph node dissection with robotic surgery although several studies describe a shorter learning curve compared with the laparoscopic approach (7,8). One should consider that robotic liver surgery (initiated since February 2021) was developed following laparoscopic liver surgery (initiated since 2005) permitting the authors to acquire expert skills in minimal invasive hepatic resection and thus the results of the learning curve of this study are biased. Furthermore, generalising these results is doubtful given the small number of robotic resections and the major expertise of this team in minimal invasive hepatobiliary surgery. Robotic use in hepatobiliary surgery is steadily growing and although indications are also expanding, its use is only supported by a low level of proof (9) and is still limited in highly specialised centres.

Finally, the scarce proportion of perihilar cholangiocarcinoma included in this study did not allow to evaluate robotic confection of hepaticojejunostomy which is supposed to be one of the major improvements in minimally invasive liver surgery with reported anastomotic fistulae and stenosis rates close to those of open surgery (10). Likewise, vascular resections and arterial or venous reconstructions often realised for perihilar cholangiocarcinoma surgery, and described as safe with the use of robot in selected patients for pancreatic resection in literature (11,12), were not studied. Last but not least, robotic approach may improve accessibility to tumours with a complex localisation compared with laparoscopy and facilitate the performance of difficult hepatectomies (13), which was not investigated by the authors.

In conclusion, minimal invasive surgery appears to provide better short-term outcomes than open approach for surgery of biliary cancers. The benefit of robotic approach

over laparoscopy is still yet to confirm.

## Acknowledgments

*Funding:* None.

## Footnote

*Provenance and Peer Review:* This article was commissioned by the editorial office, *Hepatobiliary Surgery and Nutrition*. The article did not undergo external peer review.

*Conflicts of Interest:* Both authors have completed the ICMJE uniform disclosure form (available at <https://hbsn.amegroups.com/article/view/10.21037/hbsn-23-149/coif>). The authors have no conflicts of interest to declare.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

1. Ratti F, Cipriani F, Ingallinella S, et al. Robotic Approach for Lymphadenectomy in Biliary Tumours: The Missing Ring Between the Benefits of Laparoscopic and Reproducibility of Open Approach? *Ann Surg* 2022. [Epub ahead of print]. doi: 10.1097/SLA.0000000000005748.
2. Spiegelberg J, Iken T, Diener MK, et al. Robotic-Assisted Surgery for Primary Hepatobiliary Tumors-Possibilities and Limitations. *Cancers (Basel)* 2022;14:265.
3. Regmi P, Hu HJ, Paudyal P, et al. Is laparoscopic liver resection safe for intrahepatic cholangiocarcinoma? A meta-analysis. *Eur J Surg Oncol* 2021;47:979-89.
4. Amin MB, Greene FL, Edge SB, et al. The Eighth Edition AJCC Cancer Staging Manual: Continuing to build a bridge from a population-based to a more

- "personalized" approach to cancer staging. *CA Cancer J Clin* 2017;67:93-9.
5. Hobeika C, Cauchy F, Fuks D, et al. Laparoscopic versus open resection of intrahepatic cholangiocarcinoma: nationwide analysis. *Br J Surg* 2021;108:419-26.
  6. Spolverato G, Bagante F, Weiss M, et al. Comparative performances of the 7th and the 8th editions of the American Joint Committee on Cancer staging systems for intrahepatic cholangiocarcinoma. *J Surg Oncol* 2017;115:696-703.
  7. Efanov M, Alikhanov R, Tsvirkun V, et al. Comparative analysis of learning curve in complex robot-assisted and laparoscopic liver resection. *HPB (Oxford)* 2017;19:818-24.
  8. Chen PD, Wu CY, Hu RH, et al. Robotic major hepatectomy: Is there a learning curve? *Surgery* 2017;161:642-9.
  9. Liu R, Wakabayashi G, Kim HJ, et al. International consensus statement on robotic hepatectomy surgery in 2018. *World J Gastroenterol* 2019;25:1432-44.
  10. Bustos R, Fernandes E, Mangano A, et al. Robotic hepaticojejunostomy: surgical technique and risk factor analysis for anastomotic leak and stenosis. *HPB (Oxford)* 2020;22:1442-9.
  11. Allan BJ, Novak SM, Hogg ME, et al. Robotic vascular resections during Whipple procedure. *J Vis Surg* 2018;4:13.
  12. Kauffmann EF, Napoli N, Menonna F, et al. Robotic pancreatoduodenectomy with vascular resection. *Langenbecks Arch Surg* 2016;401:1111-22.
  13. Luberic K, Sucandy I, Modasi A, et al. Applying IWATE criteria to robotic hepatectomy: is there a "robotic effect"? *HPB (Oxford)* 2021;23:899-906.

**Cite this article as:** Deyrat J, Tzedakis S. Feasibility, safety and limits of robotic approach for lymphadenectomy in biliary tumours. *HepatoBiliary Surg Nutr* 2023;12(3):404-406. doi: 10.21037/hbsn-23-149