



The power of prevention: how tenofovir and entecavir are changing the game in hepatocellular carcinoma

Endrit Shahini¹, Rossella Donghia², Antonio Facciorusso³

¹Gastroenterology Unit, National Institute of Gastroenterology-IRCCS “Saverio de Bellis”, Castellana Grotte, Italy; ²Data Science Unit, National Institute of Gastroenterology-IRCCS “Saverio de Bellis”, Castellana Grotte, Italy; ³Gastroenterology Unit, Department of Medical Sciences, University of Foggia, Foggia, Italy

Correspondence to: Prof. Antonio Facciorusso, MD, PhD. Gastroenterology Unit, Department of Medical Sciences, University of Foggia, Via L Pinto 1, 71122 Foggia, Italy. Email: antonio.facciorusso@unifg.it.

Comment on: Choi WM, Yip TC, Wong GL, *et al.* Hepatocellular carcinoma risk in patients with chronic hepatitis B receiving tenofovir- *vs.* entecavir-based regimens: Individual patient data meta-analysis. *J Hepatol* 2023;78:534-42.

Keywords: Hepatocellular carcinoma (HCC); liver cancer; cirrhosis, hepatitis B virus (HBV)

Submitted Oct 12, 2023. Accepted for publication Oct 26, 2023. Published online Nov 09, 2023.

doi: 10.21037/hbsn-23-528

View this article at: <https://dx.doi.org/10.21037/hbsn-23-528>

Introduction

Individual patient data (IPD) meta-analysis by Choi *et al.* (1) compared hepatocellular carcinoma (HCC) risk between the entecavir (ETV) and tenofovir (TDF) in treatment-naïve chronic hepatitis B (CHB) patients using a multivariable Cox proportional hazards model from 11 Asian studies, totaling 42,939 patients receiving nucleos(t)ide analogues (NAs) for more than one year. TDF was associated with a significantly lower HCC risk than ETV, particularly in patients with hepatitis B e antigen (HBeAg) positivity. The authors concluded, however, that a longer follow-up period may be necessary to clarify the impact disparities between therapies across the various subgroups.

Current American, European, and Asian-Pacific clinical practice guidelines recommend both ETV and TDF as first-line therapies for the treatment of CHB, particularly in NAs-naïve patients. Nonetheless, the relative HCC risk for TDF versus ETV treatment is controversial, and only two randomized controlled trials (RCTs) have demonstrated comparable efficacy, and excellent safety profiles (1). Importantly, the two RCTs involving 720 patients had the same follow-up duration (3 years).

This ambiguity has sparked intense interest and heated debate about which of these two preferred drugs is more effective at lowering the risk of HCC. Several meta-analyses produced contradictory results in this regard, with

some showing concordant results and others showing no difference between these two NAs (the summary table is available at: <https://cdn.amegroups.com/static/public/hbsn-23-528-1.xlsx>) (1-17).

Methodology of previous metanalyses

The heterogeneity of observational studies has hampered the majority of prior meta-analyses comparing the two NAs on HCC risk development (the summary table available at: <https://cdn.amegroups.com/static/public/hbsn-23-528-1.xlsx>). *Figure 1* depicts some of the critical factors that revolve around meta-analysis study carried out in accordance with appropriate standards.

Although the meta-analyses examined were all well conducted, the type of data presented proved to be very heterogeneous. The number of patients was not always stratified by groups (TDF and ETV) but presented as “overall”. The age variable was not presented consistently. They present different centrality indices (mean or median) in some cases and are absent in others. The follow-up data considered various time series (months or years). Cirrhosis, HBeAg positivity, hepatitis B virus (HBV) viremia, NAs naïve status, virologic response, and cumulative HCC incidence are all missing or incompletely presented for the majority of them. The effect size varies greatly since the




Figure 1 Some of the essential elements of a meta-analysis study conducted in accordance with appropriate standards. Created with “BioRender.com” (Agreement number: HK25ZPG7JM, Academic License).

relative risk (RR) was used in some cases and the hazard ratio (HR) in others. Only a few authors reported an unadjusted HR. Even heterogeneity’s I^2 was not always reported.

Furthermore, with the exception of Jeong’s paper (lack of the full text) (12), all of the meta-analyses adhere to the fundamental characteristics of a good meta-analysis. All meta-analyses are typically carried out using either “fixed effects” or “random effects” models. All authors explained different or combined uses based on different assumptions. Additionally, all authors conducted a sensitivity analysis, which is the most important part of a meta-analysis since it determines the robustness of the observed results, and subgroup analysis was used.

In terms of publication bias, Zhang *et al.* (3) and Choi *et al.* [2021] (8) used the funnel plot to test the publication bias, but did not specify the test used; Huang *et al.* (17) and Choi *et al.* [2023] (1) did not mention the bias analysis. Oh *et al.* (15), on the other hand, used an unusual test (the AS-Thomson test). Finally, all authors except Dave *et al.* (7) and Choi *et al.* (8) used the Newcastle-Ottawa scale or the Jadad scale as primary tools for assessing the quality of observational studies.

Discussion

While the majority of the observational studies were of high

quality and were conducted on a national or multicenter scale using (or not) propensity score matching (PSM) to reduce selection bias and confounding factors, some studies were affected by inadequate sample size, different study designs with diverse data sets, various ethnicities, and diverse proportions of cirrhotic patients.

Notably, the different sources of search strategy among the studies may be to blame for the inclusion of some studies in the meta-analysis process. Two recent meta-analyses did not include most of the recent high-quality studies (3,6). Among studies showing no differences in the HCC risk among the two NAs, the cohort study by Kim *et al.* [2019] (18) and the retrospective study by Lee *et al.* [2020] (19) were not included in the meta-analyses by Zhang *et al.* [2019] (3), Wang *et al.* [2020] (6), and Oh *et al.* [2022] (15); the same studies were not included in the meta-analysis by Choi *et al.* [2023] (1). Besides, data from the Spanish prospective, multicenter database CIBERHEP study of Riveiro-Barciela *et al.* [2017] (20) were only included in the meta-analyses of Wang *et al.* [2020] (6), Yuan *et al.* [2021] (13), and Dave *et al.* [2021] (7). Tseng *et al.* [2020] (10), Wang *et al.* [2020] (6), Tan *et al.* [2022] (16), and Huang *et al.* [2022] (17) did not include data from Papatheodoridis *et al.*’s multicentric European PAGE-B study [2016] (21) showing similar results between ETV and TDF monotherapies.

According to Asian research, CHB patients treated with TDF have a lower cumulative incidence rate of HCC, whereas studies conducted outside of Asia found that the incidence rates of HCC are equal between TDF and ETV treatments (the summary table available at: <https://cdn.amegroups.com/static/public/hbsn-23-528-1.xlsx>). These geographic disparities, may be caused by diverse demographic baseline characteristics, HBV genotypes, and healthcare systems. Age should also be considered as a potentially confounding variable, as ETV may have been chosen preferentially for older patients due to potential adverse bone effects associated with TDF or disease severity. HBV genotype C is also predominant in many Asian countries. Since patients with cirrhosis are more likely to develop HCC, the impact of antiviral treatment may be most clearly seen in studies on cirrhotic patients. Antiviral therapy, however, may be most beneficial early in the disease’s course. Besides, many studies did not take into account the use of aspirin, statins, and metformin, which are well-known drugs that reduce HCC risk, or the inclusion of treatment-experienced patients.

Additionally, given the moderate to significant

heterogeneity of patient characteristics within/between studies, as well as ETV's earlier introduction than TDF in East Asia (the former in 2006, the latter in 2011), where the majority of studies have been conducted, longer follow-up periods for ETV-treated patients are more frequently available. Specifically, patients treated with TDF had a significantly lower HCC rate in studies where ETV follow-up was more than 12 months longer than TDF, whereas studies with more equal follow-up between arms found no significant difference (4,10). In the meta-analysis of Choi *et al.* [2021] (8), TDF was associated with a 23% lower HCC risk compared with ETV. On the subgroup analysis, this beneficial effect persisted in cirrhotic patients. However, there was a disparity in the follow-up periods between the 2 groups even after PSM since the ETV group had longer follow-up than the TDF group, a difference of up to 33 months. Additionally, in the PAGE-B cohort of 1,935 CHB, long-term monotherapy with ETV or TDF was associated with equivalent HCC risk during similar follow-up times (7.6 and 7.5 years, respectively) (21).

According to research conducted in Asia, particularly in Korea, the controversy surrounding the use of both NAs in the treatment of HCC stems from the arbitrary nature of significance levels, which leads to contradictory conclusions from similar datasets (22). Additionally, although high statistical power can produce statistically significant results, they are not always clinically relevant. Also, it is possible that unadjusted meta-analysis estimates can produce HRs which show that TDF decreases the risk of HCC development more than ETV because patients on ETV may have a higher risk of HCC than those on TDF. Patients with prior NAs resistance, indeed, may have been at a higher risk of HCC due to incomplete viral suppression.

In addition, subgroup analyses, which are often utilized to evaluate the effects of heterogeneity on a limited number of studies, can lead to both false-positive and false-negative results. By applying PSM or covariate-adjusted estimations, meta-analyses can decrease the influence of within-study heterogeneity. However, even when PSM and covariate adjustments are used, the resulting estimations may not always be completely accurate, particularly when significant clinical variables are not included in the analysis, as is the case when administrative claims databases are used in meta-analyses.

Regardless, utilizing IPD allows for a standard analytical approach across all studies. An IPD meta-analysis has the advantage of accounting for biases across all datasets using consistent methodologies, thereby providing a more

robust estimate. Despite the obstacles involved in obtaining agreement and ethical approval from a sufficient number of studies, as well as the potential lack of versatility due to the majority of studies being conducted in East Asia, this approach would address many of the issues brought up by aggregate meta-analyses.

Ultimately, it is imperative to carefully consider the source of funding for studies, as it can have an indirect impact on the results and lead to potential bias.

Conclusions

High-quality multicenter RCTs are unlikely to be conducted to identify subgroups that benefit the most from a specific NA due to the lengthy follow-up period and large number of participants required due to the low HCC incidence in CHB patients.

Future observational studies, even if limited by their nature, could compensate for this gap if they focus on distinct subgroups such as those with cirrhosis, different ethnicities, and HBeAg seropositivity.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Hepatobiliary Surgery and Nutrition*. The article did not undergo external peer review.

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://hbsn.amegroups.com/article/view/10.21037/hbsn-23-528/coif>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the

original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- Choi WM, Yip TC, Wong GL, et al. Hepatocellular carcinoma risk in patients with chronic hepatitis B receiving tenofovir- vs. entecavir-based regimens: Individual patient data meta-analysis. *J Hepatol* 2023;78:534-42.
- Han Y, Zeng A, Liao H, et al. The efficacy and safety comparison between tenofovir and entecavir in treatment of chronic hepatitis B and HBV related cirrhosis: A systematic review and Meta-analysis. *Int Immunopharmacol* 2017;42:168-75.
- Zhang Z, Zhou Y, Yang J, et al. The effectiveness of TDF versus ETV on incidence of HCC in CHB patients: a meta analysis. *BMC Cancer* 2019;19:511.
- Li M, Lv T, Wu S, et al. Tenofovir versus entecavir in lowering the risk of hepatocellular carcinoma development in patients with chronic hepatitis B: a critical systematic review and meta-analysis. *Hepatol Int* 2020;14:105-14.
- Gu L, Yao Q, Shen Z, et al. Comparison of tenofovir versus entecavir on reducing incidence of hepatocellular carcinoma in chronic hepatitis B patients: A systematic review and meta-analysis. *J Gastroenterol Hepatol* 2020;35:1467-76.
- Wang X, Liu X, Dang Z, et al. Nucleos(t)ide Analogues for Reducing Hepatocellular Carcinoma in Chronic Hepatitis B Patients: A Systematic Review and Meta-Analysis. *Gut Liver* 2020;14:232-47.
- Dave S, Park S, Murad MH, et al. Comparative Effectiveness of Entecavir Versus Tenofovir for Preventing Hepatocellular Carcinoma in Patients with Chronic Hepatitis B: A Systematic Review and Meta-Analysis. *Hepatology* 2021;73:68-78.
- Choi WM, Choi J, Lim YS. Effects of Tenofovir vs Entecavir on Risk of Hepatocellular Carcinoma in Patients With Chronic HBV Infection: A Systematic Review and Meta-analysis. *Clin Gastroenterol Hepatol* 2021;19:246-258.e9.
- Liu H, Shi Y, Hayden JC, et al. Tenofovir Treatment Has Lower Risk of Hepatocellular Carcinoma than Entecavir Treatment in Patients with Chronic Hepatitis B: A Systematic Review and Meta-Analysis. *Liver Cancer* 2020;9:468-76.
- Tseng CH, Hsu YC, Chen TH, et al. Hepatocellular carcinoma incidence with tenofovir versus entecavir in chronic hepatitis B: a systematic review and meta-analysis. *Lancet Gastroenterol Hepatol* 2020;5:1039-52.
- Cheung KS, Mak LY, Liu SH, et al. Entecavir vs Tenofovir in Hepatocellular Carcinoma Prevention in Chronic Hepatitis B Infection: A Systematic Review and Meta-Analysis. *Clin Transl Gastroenterol* 2020;11:e00236.
- Jeong S, Cho Y, Park SM, et al. Differential Effectiveness of Tenofovir and Entecavir for Prophylaxis of Hepatocellular Carcinoma in Chronic Hepatitis B Patients Depending on Coexisting Cirrhosis and Prior Exposure to Antiviral Therapy: A Systematic Review and Meta-analysis. *J Clin Gastroenterol* 2021;55:e77-86.
- Yuan J, Peng Y, Hao FB, et al. No difference in hepatocellular carcinoma risk in chronic hepatitis B patients treated with tenofovir vs entecavir: evidence from an updated meta-analysis. *Aging (Albany NY)* 2021;13:7147-65.
- Yuan BH, Li RH, Huo RR, et al. Lower risk of hepatocellular carcinoma with tenofovir than entecavir treatment in subsets of chronic hepatitis B patients: an updated meta-analysis. *J Gastroenterol Hepatol* 2022;37:782-94.
- Oh H, Lee HY, Kim J, et al. Systematic Review with Meta-Analysis: Comparison of the Risk of Hepatocellular Carcinoma in Antiviral-Naive Chronic Hepatitis B Patients Treated with Entecavir versus Tenofovir: The Devil in the Detail. *Cancers (Basel)* 2022;14:2617.
- Tan DJH, Ng CH, Tay PWL, et al. Risk of Hepatocellular Carcinoma With Tenofovir vs Entecavir Treatment for Chronic Hepatitis B Virus: A Reconstructed Individual Patient Data Meta-analysis. *JAMA Netw Open* 2022;5:e2219407.
- Huang ZH, Lu GY, Qiu LX, et al. Risk of hepatocellular carcinoma in antiviral treatment-naïve chronic hepatitis B patients treated with entecavir or tenofovir disoproxil fumarate: a network meta-analysis. *BMC Cancer* 2022;22:287.
- Kim SU, Seo YS, Lee HA, et al. A multicenter study of entecavir vs. tenofovir on prognosis of treatment-naïve chronic hepatitis B in South Korea. *J Hepatol* 2019;71:456-64.
- Lee SW, Kwon JH, Lee HL, et al. Comparison of tenofovir and entecavir on the risk of hepatocellular carcinoma and mortality in treatment-naïve patients with chronic hepatitis B in Korea: a large-scale, propensity score analysis. *Gut* 2020;69:1301-8.
- Riveiro-Barciela M, Tabernero D, Calleja JL, et al.

- Effectiveness and Safety of Entecavir or Tenofovir in a Spanish Cohort of Chronic Hepatitis B Patients: Validation of the Page-B Score to Predict Hepatocellular Carcinoma. *Dig Dis Sci* 2017;62:784-93.
21. Papatheodoridis G, Dalekos G, Sypsa V, et al. PAGE-B predicts the risk of developing hepatocellular carcinoma in Caucasians with chronic hepatitis B on 5-year antiviral therapy. *J Hepatol* 2016;64:800-6.
22. Choi WM, Yip TC, Lim YS, et al. Methodological challenges of performing meta-analyses to compare the risk of hepatocellular carcinoma between chronic hepatitis B treatments. *J Hepatol* 2022;76:186-94.

Cite this article as: Shahini E, Donghia R, Facciorusso A. The power of prevention: how tenofovir and entecavir are changing the game in hepatocellular carcinoma. *HepatoBiliary Surg Nutr* 2023;12(6):936-940. doi: 10.21037/hbsn-23-528