



Shifting paradigm of quality of life after pancreatoduodenectomy for pancreas cancer

Kyuseok Im, Victoria V. O'Connor

Kaiser Permanente Los Angeles Medical Center, Los Angeles, CA, USA

Correspondence to: Victoria V. O'Connor, MD. Kaiser Permanente Los Angeles Medical Center, 4760 Sunset Blvd., Los Angeles, CA 90027, USA.

Email: Victoria.v.o'connor@kp.org.

Comment on: Zhang C, Zirona A, Vierkant RA, *et al.* Quality of Life and Gastrointestinal Symptoms in Long-term Survivors of Pancreatic Cancer Following Pancreatoduodenectomy. *Ann Surg* 2024;279:842-9.

Keywords: Quality of life (QOL); psychosocial; pancreas cancer; pancreatoduodenectomy

Submitted Feb 02, 2024. Accepted for publication Mar 04, 2025. Published online Mar 25, 2025.

doi: 10.21037/hbsn-2025-79

View this article at: <https://dx.doi.org/10.21037/hbsn-2025-79>

As long-term survival has improved with advancements in pancreatic cancer treatment in the modern era, quality of life (QOL) among pancreatic cancer patients following pancreaticoduodenectomy (PD), i.e., Whipple procedure, warrants understanding and addressing within the medical community. The article by Zhang *et al.*, titled “Quality of Life and Gastrointestinal Symptoms in Long-Term Survivors of Pancreatic Cancer Following Pancreaticoduodenectomy”, provides valuable insights into the balance between the overall survival benefits of this complex procedure and patient-reported long-term post-operative effects on patient well-being.

Historically, PD, first described by Alessandro Codivilla in the late 19th century, was a technical marvel and was later refined by Dr. Whipple in the 1930s (1,2). Despite further surgical advancements in the 1950s and 1960s, high morbidity and mortality rates persisted, raising questions about whether the risks of surgery were justified given the generally poor prognosis of pancreatic cancer (3). Even as Cameron *et al.* established new benchmarks for post-Whipple morbidity and mortality in the 2000s, this technical success could not overcome the persistent challenge of low overall survival rates (4). The perception of both pancreatic cancer and the Whipple procedure remained negative for many years, with both the disease and the operation becoming synonymous with a bleak prognosis. However, modern advancements in systemic treatments, imaging modalities, surgical techniques, and post-operative

management have led to improvements in mortality and morbidity associated with the Whipple procedure, as well as increased long-term survival among pancreas cancer patients (5). As these outcomes improved, so has our understanding of the physiological and psychological impacts of PD.

The current study by Zhang *et al.* sheds light on key aspects of pancreatic cancer survivorship, particularly QOL and gastrointestinal (GI) symptoms. The study found that long-term survivors of pancreatic cancer who had undergone PD reported a relatively higher QOL compared to the general US population. Although patients experienced chronic post-operative GI symptoms, including pancreatic insufficiency, reflux, and delayed gastric emptying, these symptoms did not significantly affect their overall QOL. Most patients indicated they would undergo surgery again. However, those who reported poorer QOL attributed it to post-operative complications such as pancreatic fistulas, delayed gastric emptying, and incisional hernias. Additionally, patients who expressed reluctance to undergo PD again reported a higher frequency of GI symptoms. These insights should be integrated into the management of recovery and the discussion of long-term lifestyle adjustments. Using the evidence from Zhang and similar studies, surgeons can offer evidence-based reassurance to patients who may be anxious about undergoing this major procedure, while also setting realistic expectations for recovery and the potential long-term

physiological effects. Overall, long-term cancer survival appears to be the primary driver of QOL, despite ongoing physical challenges.

In our own study, we found that while patients may experience chronic GI issues following surgery, the overall impact on QOL is more significantly influenced by mental well-being and financial factors than by the physiological consequences of PD (6). Furthermore, there were notable differences between patients who underwent PD for malignant versus benign conditions. Patients treated for malignancy reported a significantly worse QOL, citing psychological and socioeconomic challenges related to pancreatic cancer, including feelings of diminished utility, hopelessness, loss of control, uncertainty about the future, dissatisfaction with physical appearance, memory issues, fear of relapse, and financial strain. In contrast, physical health did not have a significant association with QOL, regardless of whether the pathology was benign or malignant. These findings highlight a concern for the undertreatment of depression in patients with pancreatic cancer. Large healthcare systems must better address this gap by providing comprehensive support and counseling, even for patients who have undergone curative surgery, as non-surgical factors tend to have a more profound impact on QOL than the physiological effects of the surgery itself.

While the management of GI issues related to PD, such as malabsorption, diarrhea, and weight loss, is essential, multiple studies emphasize the significant psychological and socioeconomic challenges faced by cancer patients. Although one might expect QOL to improve with remission, Firkins *et al.* report that mental health issues continue to negatively impact QOL among long-term cancer survivors (7). This link between cancer and mental health is further emphasized by Michoglou *et al.*, who explore the mechanisms connecting depression with pancreatic cancer and highlight the complexities of managing mental health in the context of cancer itself, rather than solely focusing on surgery-related morbidities (8). While pylorus-preserving surgery might be expected to improve QOL by alleviating GI symptoms like dumping syndrome, diarrhea, and weight loss, Ohtsuka *et al.* found that pylorus-preservation has no significant effect on QOL (9). They concluded that psychosocial QOL remains low regardless of the surgical approach, underscoring the need for long-term mental health support, independent of the physiological effects of surgery.

As prognosis and survivorship for pancreatic cancer continue to improve, recognizing that the psychosocial

impacts often outweigh the physical consequences of surgical treatment becomes increasingly important. The need for comprehensive support grows to address both the physical and non-physical challenges patients face. In a large survey of cancer survivors, unmet needs related to physical health accounted for less than 40% of all reported needs (10). The remaining unmet needs spanned financial, educational, autonomy-related, healthcare system resources, emotional, mental, interpersonal, societal, existential, and personal identity concerns. Dr. Zhang and colleagues contribute to this evolving understanding of holistic cancer care, demonstrating that post-surgical physical sequelae are generally less disruptive to long-term survivors' QOL than the ongoing socioeconomic and psychological burdens associated with a history of pancreatic cancer. This highlights the need for oncologic surgical care to extend beyond managing post-operative symptoms to include long-term support for the intangible morbidities that persist after curative-intent resection.

In summary, as the field of pancreatic cancer treatment, including its cornerstone treatment modality PD, continues to evolve, we are experiencing a paradigm shift in what defines “best outcomes” for patients. This should include not only objective survival but also the multifaceted factors that contribute to QOL. While scientific, medical, and surgical advancements will continue to enhance post-operative physiologic outcomes and long-term survival, the psychosocial toll of the disease must not be overlooked. A deeper understanding of these non-physical aspects of patient well-being, and the application of this knowledge to develop more nuanced, comprehensive care strategies, should remain a central focus in advancing the field. Pancreatic cancer survivors require more than just surgical intervention—they need and deserve ongoing socioeconomic and psychological support to achieve a meaningful QOL.

Acknowledgments

None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *HepatoBiliary Surgery and Nutrition*. The article did not undergo external peer review.

Funding: None.

Conflicts of Interest: Both authors have completed the ICMJE uniform disclosure form (available at <https://hbsn.amegroups.com/article/view/10.21037/hbsn-2025-79/coif>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Schnelldorfer T, Sarr MG. Alessandro Codivilla and the first pancreatoduodenectomy. *Arch Surg* 2009;144:1179-84.
2. Whipple AO, Parsons WB, Mullins CR. Treatment of carcinoma of the ampulla of Vater. *Ann Surg* 1935;102:763-79.
3. Gudjonsson B. Cancer of the pancreas. 50 years of surgery. *Cancer* 1987;60:2284-303.
4. Cameron JL, Riall TS, Coleman J, et al. One thousand consecutive pancreaticoduodenectomies. *Ann Surg*. 2006;244:10-5.
5. Del Chiaro M, Sugawara T, Karam SD, et al. Advances in the management of pancreatic cancer. *BMJ* 2023;383:e073995.
6. Silva T, Wagler A, Nelson D, et al. Quality of life after pancreatoduodenectomy: Is the outcome predetermined by the diagnosis? *J Surg Oncol* 2023;128:1080-6.
7. Firkins J, Hansen L, Driessnack M, et al. Quality of life in "chronic" cancer survivors: a meta-analysis. *J Cancer Surviv* 2020;14:504-17.
8. Michoglou K, Ravinthiranathan A, San Ti S, et al. Pancreatic cancer and depression. *World J Clin Cases* 2023;11:2631-6.
9. Ohtsuka T, Yamaguchi K, Ohuchida J, et al. Comparison of quality of life after pylorus-preserving pancreatoduodenectomy and Whipple resection. *Hepatogastroenterology* 2003;50:846-50.
10. Burg MA, Adorno G, Lopez ED, et al. Current unmet needs of cancer survivors: analysis of open-ended responses to the American Cancer Society Study of Cancer Survivors II. *Cancer* 2015;121:623-30.

Cite this article as: Im K, O'Connor VV. Shifting paradigm of quality of life after pancreatoduodenectomy for pancreas cancer. *HepatoBiliary Surg Nutr* 2025;14(2):316-318. doi: 10.21037/hbsn-2025-79