

# Identification of potential associated factors for stress urinary incontinence in women: a retrospective study

Dongmei Wei<sup>1,2</sup>^, Jian Meng<sup>1,2</sup>^, Yueting Zhang<sup>1,2</sup>^, Yueyue Chen<sup>1,2</sup>^, Jijie Li<sup>2,3</sup>^, Xiaoyu Niu<sup>1,2</sup>^

<sup>1</sup>Department of Gynecology and Obstetrics, West China Second Hospital, Sichuan University, Chengdu, China; <sup>2</sup>Key Laboratory of Birth Defects and Related Diseases of Women and Children (Sichuan University), Ministry of Education, Chengdu, China; <sup>3</sup>Medical Records Management Department, West China Second Hospital, Sichuan University, Chengdu, China

*Contributions:* (I) Conception and design: D Wei, X Niu; (II) Administrative support: D Wei, Y Chen; (III) Provision of study materials or patients: J Meng, Y Zhang (IV) Collection and assembly of data: Y Chen, Y Zhang; (V) Data analysis and interpretation: D Wei, J Li; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

*Correspondence to:* Xiaoyu Niu. Department of Gynecology and Obstetrics, West China Second Hospital, Sichuan University, No. 20, Section 3, South Renmin Road, Wuhou District, Chengdu, China. Email: hxfeynxy@163.com.

**Background:** This study sought to analyze the potential associated factors for female stress urinary incontinence (SUI).

**Methods:** A total of 5,013 women were screened for pelvic floor function at the West China Second Hospital of Sichuan University from January 2015 to January 2019. Of these, 410 patients were diagnosed with SUI. A single-factor Chi-square test and multi-factor logistic regression analysis were conducted to examine the relationship between pre-pregnancy urinary incontinence, vaginal delivery, menopause, and hormone therapy, chronic cough, and smoking, and postpartum SUI.

**Results:** The postpartum SUI rate in patients with urinary incontinence during pregnancy was 19.33%, while that of patients without urinary incontinence was only 5.44%. The rates of urinary incontinence in patients experiencing vaginal delivery or cesarean delivery were 13.62% and 4.36%, respectively. The SUI incidences in patients with or without a family genetic history of SUI were 28.46% and 7.48%, respectively. The incidence rates of SUI in smoking and non-smoking patients were 18.92% and 8.39%. The rate of SUI in patients with chronic cough (16.46%) behaved significantly differently from those with non-chronic cough (8.21%). The occurrence of SUI was highly correlated with the following factors, including pre-pregnancy urinary incontinence (OR =5.256; 95% CI: 2.061–13.409; P<0.001), urological incontinence during the pregnancy period (OR =2.965; 95% CI: 2.111–4.163; P<0.001), vaginal delivery (OR =4.028; 95% CI: 2.909–5.577; P<0.001), and genetic history (OR =4.341; 95% CI: 2.8–6.73; P<0.001).

**Conclusions:** The occurrence of SUI is highly related to a history of urinary incontinence, the delivery mode, chronic cough, smoking, and genetic history. Further, urinary incontinence before and during pregnancy, natural delivery, and genetic history are important independent high-associated factors for SUI. Our findings show the importance of screening for the above associated factors in association with SUI.

Keywords: Stress urinary incontinence (SUI); incontinence; vaginal delivery; genetic history

Submitted Jun 24, 2022. Accepted for publication Aug 17, 2022. doi: 10.21037/atm-22-3539 View this article at: https://dx.doi.org/10.21037/atm-22-3539

^ ORCID: Dongmei Wei, 0000-0001-6922-0296; Jian Meng, 0000-0002-5875-4177; Yueting Zhang, 0000-0002-1238-3526; Jijie Li, 0000-0002-4762-9709; Xiaoyu Niu, 0000-0002-9562-1795; Yueyue Chen, 0000-0002-2745-4622.

# Introduction

Stress urinary incontinence (SUI) is a common gynecologic urinary disorder defined as involuntary urine outflow, which is characterized by the normal absence of enuresis and an automatic flow of urine following a sudden increase in abdominal pressure (1). The incidence of urinary dysfunction in adult females is approximately 33%, and 50% of urinary incontinence is SUI (2). At present, there is no ideal treatment for SUI. Surgery is a common treatment, however, it is extremely painful for SUI patients (3). There is therefore a need to explore risk factors for SUI in order to provide early intervention for patients.

The initiation and progression of SUI has been linked to age, being overweight, diabetes, and obstetric trauma (4,5). Further, vaginal delivery, menopausal and hormone therapy are known to affect the functionality of the inferior ureter (6,7). The multifactorial risk of SUI can be categorized into predisposing, triggering and dysfunctional factors that have been shown to have a very complex impact on the development of SUI (8). There are several limitations and shortcomings in the previous studies. In addition, the sampling methods of previous studies may not be representative of all women. Thirdly, the diagnosis of SUI in some studies was based on the patient's chief complaint without clinical examination data such as radiological or urodynamic findings. The current study sought to increase awareness of this common medical problem to improve the treatment effect.

Our present study showed that the occurrence of SUI was highly associated with a history of urinary incontinence, the delivery mode, chronic cough, smoking, and genetic history. Further, urinary incontinence before and during pregnancy, natural delivery, and genetic history were important independent high-associated factors for SUI. Our findings would provide the first screening for SUI-related relevant factors. We present the following article in accordance with the STROBE reporting checklist (available at https://atm. amegroups.com/article/view/10.21037/atm-22-3539/rc).

## Methods

This hospital-based retrospective study analyzed the data of 5013 patients who presented with or without urinary incontinence from January 2015 to January 2019 at the West China Second Hospital of Sichuan University. The identification of SUI was dependent on the Guidelines for the diagnosis and treatment of female SUI. According to Wei et al. Potential associated factors for SUI women

the classic symptoms of SUI, urine overflow occurs during varying levels of increased abdominal pressure (such as laughing, coughing, sneezing, or walking); the abnormal urine overflow disappeared if the pressure stopped. The divergent exposure factors included a history of urinary incontinence, age, body mass index (BMI), job (i.e., mental worker or manual worker), number of deliveries, the delivery method (i.e., Caesarean section or vaginal delivery), diabetes, hypertension, genetic history, smoking, chronic cough, and constipation. The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013). This study was approved by the Medical Ethics Association of West China Second Hospital of Sichuan University [2018.No.(069)]. All patients signed informed consent.

## Statistical analysis

The patient characteristics were expressed as the absolute number (percentage) or mean  $\pm$  standard deviation (SD). For those whose measurement data was normally distributed, the independent sample *t* test was used for those whose variances between groups were homogeneous. Count data and categorical variables were tested by chi-square test. The factors with P $\leq$ 0.10 in univariate analysis were further included in multivariate logistic regression analysis. The relationship between pre-pregnancy urinary incontinence, vaginal delivery, menopause, and hormone therapy, chronic cough, smoking, and postpartum urinary incontinence was calculated. All statistical tests were 2-sided probability tests. P value <0.05 was considered statistically significant. Statistical analyses were conducted using SPSS26.0 software.

# **Results**

# The characteristics of patients in this study and potential associated factors for the occurrence of SUI

First, we conducted a single-factor analysis to identify the relevant factors related to SUI (*Table 1*). Among the patients diagnosed with SUI, the postpartum urinary incontinence rate of patients with urinary incontinence before pregnancy was 45.16%, while that of patients without urinary incontinence before pregnancy was only 6.73% (P<0.001). The postpartum urinary incontinence rate of patients with urinary incontinence during pregnancy was 19.33%, while that of patients without urinary incontinence during pregnancy during urinary incontinence during pregnancy was 19.33%, while that of patients without urinary incontinence during pregnancy was 19.33%.

# Annals of Translational Medicine, Vol 10, No 18 September 2022

# Page 3 of 7

Table 1 Patient features in this study, and potential associated factors for the occurrence of SUI based on a single-factor analysis

| Patient features                      | Non-SUI | SUI | $\chi^2$ value | P value |
|---------------------------------------|---------|-----|----------------|---------|
| Urinary incontinence before pregnancy |         |     | 63.6245        | <0.0001 |
| Negative                              | 3,547   | 256 |                |         |
| Positive                              | 17      | 14  |                |         |
| Urinary incontinence during pregnancy |         |     | 116.8176       | <0.0001 |
| Negative                              | 3,201   | 184 |                |         |
| Positive                              | 363     | 87  |                |         |
| Age                                   |         |     | 0.0527         | 0.8184  |
| <55 years                             | 2,763   | 263 |                |         |
| ≥55 years                             | 1,818   | 169 |                |         |
| BMI                                   |         |     | 0.5597         | 0.4544  |
| <25 kg/m²                             | 3,664   | 339 |                |         |
| ≥25 kg/m²                             | 917     | 46  |                |         |
| dof                                   |         |     | 1.2491         | 0.2637  |
| Mental worker                         | 3,229   | 328 |                |         |
| Manual worker                         | 376     | 46  |                |         |
| Number of deliveries                  |         |     | 3.0686         | 0.0798  |
| 1                                     | 2,635   | 234 |                |         |
| 2                                     | 1,848   | 196 |                |         |
| Delivery method                       |         |     | 123.507        | <0.0001 |
| Caesarean section                     | 2,304   | 105 |                |         |
| Vaginal delivery                      | 1,922   | 303 |                |         |
| Diabetes                              |         |     | 0.0775         | 0.7807  |
| Negative                              | 4,187   | 388 |                |         |
| Positive                              | 253     | 22  |                |         |
| Hypertension                          |         |     | 0.2489         | 0.6178  |
| Negative                              | 4,114   | 376 |                |         |
| Positive                              | 69      | 5   |                |         |
| Genetic history                       |         |     | 130.921        | <0.0001 |
| Negative                              | 4,166   | 337 |                |         |
| Positive                              | 176     | 70  |                |         |
| Smoking                               |         |     | 3.9786         | 0.0331  |
| Negative                              | 4,400   | 403 |                |         |
| Positive                              | 30      | 7   |                |         |

Table 1 (continued)

# Page 4 of 7

# Wei et al. Potential associated factors for SUI women

#### Table 1 (continued)

| · · · ·          |         |     |                |         |
|------------------|---------|-----|----------------|---------|
| Patient features | Non-SUI | SUI | $\chi^2$ value | P value |
| Cough            |         |     | 6.8981         | 0.0086  |
| Negative         | 4,358   | 390 |                |         |
| Positive         | 66      | 13  |                |         |
| Constipation     |         |     | 2.4143         | 0.1202  |
| Negative         | 3,409   | 301 |                |         |
| Positive         | 1,029   | 109 |                |         |
|                  |         |     |                |         |

SUI, stress urinary incontinence.

Table 2 The multivariate logistic regression analysis indicated the independent associated factors for SUI

| Item  | OR    | 95% CI      | P value |
|---|-------|-------------|---------|
| Urinary incontinence before pregnancy (positive vs. negative) | 5.256 | 2.06-13.409 | 0.0005  |
| Urinary incontinence during pregnancy (positive vs. negative) | 2.965 | 2.111-4.163 | <0.0001 |
| Age (≥55 <i>vs.</i> <55 years)                                | 0.792 | 0.558–1.123 | 0.1903  |
| BMI (≥25 <i>vs.</i> <25 kg/m²)                                | 1.273 | 0.89–1.819  | 0.1862  |
| Mental worker vs. manual worker                               | 0.954 | 0.558-1.632 | 0.8636  |
| Number of deliveries (2 vs. 1)                                | 1.454 | 1.057-2.002 | 0.0216  |
| Natural delivery vs. Caesarean section                        | 4.028 | 2.909–5.577 | <0.0001 |
| Diabetes (positive vs. negative)                              | 1.192 | 0.659–2.157 | 0.5612  |
| Hypertension (positive vs. negative)                          | 0.684 | 0.149–3.133 | 0.625   |
| Genetic history   | 4.341 | 2.8-6.73    | <0.0001 |
| Smoking (positive vs. negative)                               | 1.933 | 0.392-9.522 | 0.418   |
| Cough (positive vs. negative)                                 | 1.432 | 0.514-3.992 | 0.4923  |
| Constipation (positive vs. negative)                          | 1.151 | 0.83–1.595  | 0.3992  |

SUI, stress urinary incontinence.

pregnancy was only 5.44% (P<0.001). The rates of urinary incontinence in patients experiencing vaginal delivery or cesarean delivery were 13.62% and 4.36%, respectively (P<0.001). The prevalence of urological incontinence in patients with or without a family genetic history of SUI were 28.46% and 7.48% (P<0.001). The prevalence of urological incontinence in smoking and non-smoking patients were 18.92% and 8.39%. The rate of SUI in patients with chronic cough (16.46%) behaved significantly differently from those with non-chronic cough (8.21%).

# Multivariate logistic regression of the independent associated factors for SUI

Based on the single-factor analysis, to exclude the influence of confounding factors on the occurrence of SUI, we conducted a multi-factor logistic analysis. As *Table 2* shown, the multivariate logistic regression revealed that the occurrence of SUI was highly correlated with the following factors, including urinary incontinence before pregnancy [odds ratio (OR) =5.256, 95% confidence interval (CI):

# Annals of Translational Medicine, Vol 10, No 18 September 2022

2.061–13.409, P<0.001], urinary incontinence during pregnancy (OR =2.965, 95% CI: 2.111–4.163, P<0.001), vaginal delivery (OR =4.028, 95% CI: 2.909–5.577, P<0.001), and genetic history (OR =4.341, 95% CI: 2.8–6.73, P<0.001).

# Discussion

Normally, the complete support of adjacent tissues of the posterior pubic urethra is necessary to maintain urine control under pressure (8,9). In cases of vaginal wall laxity and pelvic tissue damage, the contractile muscles of the pubococcygeus caudalis cannot act as maintenance and the urethra autonomously fails to close, leading to the development of urinary incontinence (10,11). Somewhat concerningly, the number of patients with SUI has been steadily growing over the past decade (12,13). This study showed that the development of SUI was highly associated with a history of urinary incontinence, mode of delivery, chronic cough, smoking and genetic history. In addition, incontinence before and during pregnancy, spontaneous delivery and genetic history were independent factors associated with SUI.

Age is thought to be associated with decreased sensitivity, detrusor activity, and bladder capacity, all of which can induce SUI (14,15). Previous research has shown an association between age and SUI (16,17), however, we were unable to confirm any relationship between age and SUI in the present study. Further, previously existing urinary incontinence in patients has been recognized as a potential risk factor for SUI (18,19). SUI is more prevalent in individuals who have experienced incontinence. In line with previous studies, we found a high proportion of patients had urinary incontinence after pregnancy. It is thought that decreased estrogen receptor levels are involved in the development of premenopausal SUI, as SUI appears to occur in conjunction with estrogen deficiency (20). However, the present study did not find any difference between menopausal and non-menopausal patients. In the current study, we were not able to confirm a significant correlation between SUI and menopausal and nonmenopausal patients.

There is emerging evidence that hypertension or hypertension treatment drugs trigger or exacerbate SUI (21,22). For example, adrenergic-receptor antagonists have been shown to inhibit the closure of the bladder neck and cause SUI (21). Diuretics can inhibit the reabsorption of urine and increase the urine produced by the kidneys, thereby aggravating the original urine (22). However, there was no difference in the incidence of SUI between hypertensive and non-hypertensive patients.

Obesity has been described as a cause of SUI, as increased intraabdominal pressure leads to increased intravesical pressure (23). However, our results showed that patients with different BMIs had similar SUI rates. Similarly, early research has suggested that SUI is more common in diabetic women than non-diabetic women, which indicates that diabetic microvascular complications reduce the normal control function of the bladder (24,25). Additionally, no such difference in the predominance of SUI was observed in the present study.

Some studies have suggested that multiple pregnancies, multiple births, and perineal lacerations are associated factors for SUI (26,27). Currently, most scholars have believed that changes in the anatomical pelvic-floor structure supporting tissue and the pelvic-floor impairment function are the main causes of SUI (28). Multiple pregnancies and childbirths have the cumulative effects on the damage to the pelvic floor sphincter and ligaments (29). Postpartum pelvic floor tissue cannot be restored to its original muscle strength and elasticity, and there is a cumulative effect of pelvic floor functional deficits or injuries. Additionally, perineal lacerations can cause during childbirth lead to the formation of scars after delivery. Perineal tension is significantly reduced, which will result in pelvic floor tissue relaxation and promotes the occurrence of SUI (30,31). In the present investigation, the rate of urinary incontinence in vaginal delivery patients was higher than that of cesarean delivery patients, which indicated that vaginal delivery increased the risk of SUI.

Given the potential for the development of SUI, a genetic history of SUI is an independent risk factor for SUI. This is suggested that a genetic history of SUI will increase patients' susceptibility to SUI. To ensure the appropriate interpretation of the sample data, relevant information needs to be gathered on the composition of SUI from similar cohorts. We found that chronic cough was more closely linked to a risk of developing SUI than no cough. However, the associations between chronic cough and SUI need to be more thoroughly explored in the future.

The present study had a number of strengths, including a sufficient sample size. To date, this study included the largest series of samples examined in association with SUI associated factors. The present study also had rigorous participant characterization and adopted comprehensive analytical approaches. Thus, the results of this large study

#### Page 6 of 7

have good suggestive and reference significance.

This study had some limitations due to its retrospective design. We observed a number of potential associations, however, due to the retrospective results, we were unable to imply direct causation. In addition, the entire testing process relied heavily on the patient's chief complaint and the patient's medical history might be affected by recall bias. The information of some patients was also missing, which might affect the replicability of our results. Finally, because our data were collected from a center, there might be the sample bias.

Despite the above limitations, our results revealed a correlation between a history of urinary incontinence, the delivery mode, chronic cough, smoking, and genetic history, and the subsequent development of SUI. In addition, urinary incontinence before and during pregnancy, vaginal delivery and genetic history were found to be important independent high correlates of SUI, which may help in counselling patients on the risk of SUI. Our findings have emphasized the importance of screening for the abovementioned associated factors in association with SUI. Additional in-depth studies evaluating the biological and histological characteristics of women with SUI need to be conducted.

# Acknowledgments

Funding: None.

# Footnote

*Reporting Checklist:* The authors have completed the STROBE reporting checklist. Available at https://atm. amegroups.com/article/view/10.21037/atm-22-3539/rc

*Data Sharing Statement:* Available at https://atm.amegroups. com/article/view/10.21037/atm-22-3539/dss

*Conflicts of Interest:* All authors have completed the ICMJE uniform disclosure form (available at https://atm. amegroups.com/article/view/10.21037/atm-22-3539/coif). The authors have no conflicts of interest to declare.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The study was conducted in accordance with the Declaration of Helsinki (as

revised in 2013). This study was approved by the Medical Ethics Association of West China Second Hospital of Sichuan University [2018.No.(069)]. All patients signed informed consent.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: https://creativecommons.org/licenses/by-nc-nd/4.0/.

# References

- 1. Pires T, Pires P, Moreira H, et al. Assessment of pelvic floor muscles in sportswomen: Quality of life and related factors. Phys Ther Sport 2020;43:151-6.
- Whooley J, Cunnane EM, Do Amaral R, et al. Stress Urinary Incontinence and Pelvic Organ Prolapse: Biologic Graft Materials Revisited. Tissue Eng Part B Rev 2020;26:475-83.
- Kelsey R. Urinary incontinence: Comparing AFS and TVT for stress urinary incontinence. Nat Rev Urol 2015;12:5.
- Palmieri S, Frigerio M, Spelzini F, et al. Risk factors for stress urinary incontinence recurrence after single-incision sling. Neurourol Urodyn 2018;37:1711-6.
- Lin YH, Chang SD, Hsieh WC, et al. Persistent stress urinary incontinence during pregnancy and one year after delivery; its prevalence, risk factors and impact on quality of life in Taiwanese women: An observational study. Taiwan J Obstet Gynecol 2018;57:340-5.
- Li T, Zhang YJ, Zhang HL, et al. Prevalence and Risk Factors of Stress Urinary Incontinence Among Perimenopausal Women and Its Influence on Daily Life in Women with Sexual Desire Problem. Curr Med Sci 2019;39:615-21.
- Swain D, Parida SP, Jena SK, et al. Prevalence and risk factors of obstetric fistula: implementation of a need-based preventive action plan in a South-eastern rural community of India. BMC Womens Health 2020;20:40.
- Kowalik CG, Dmochowski RR, De EJB. Surgery for female SUI: The ICI algorithm. Neurourol Urodyn 2019;38 Suppl 4:S21-7.
- 9. Xu C, Chen M, Fu J, et al. Urinary incontinence status and

## Annals of Translational Medicine, Vol 10, No 18 September 2022

risk factors in women aged 50-70 years: a cross-sectional study in Hunan, China. Int Urogynecol J 2021;32:95-102.

- Berger AA, Tan-Kim J, Menefee SA. Long-term Risk of Reoperation After Synthetic Mesh Midurethral Sling Surgery for Stress Urinary Incontinence. Obstet Gynecol 2019;134:1047-55.
- 11. Paz-Levy D, Weintraub AY, Reuven Y, et al. Prevalence and risk factors for urinary tract infection following stress urinary incontinence surgery with two midurethral sling procedures. Int J Gynaecol Obstet 2018;143:333-8.
- 12. Denisenko AA, Clark CB, D'Amico M, et al. Evaluation and management of female urinary incontinence. Can J Urol 2021;28:27-32.
- Padmanabhan P, Dmochowski R. Urinary incontinence in women: a comprehensive review of the pathophysiology, diagnosis and treatment. Minerva Ginecol 2014;66:469-78.
- 14. Kohler A, Kirschner-Hermanns R, Staat M, et al. Female stress incontinence: aspects of pathogenesis and functional anatomy. Aktuelle Urol 2018;49:47-51.
- Selecka M, Hagovska M, Bukova A, et al. Influence of sports groups on the risk of stress urinary incontinence in sportswomen. Eur J Obstet Gynecol Reprod Biol 2021;264:374-9.
- Mathieson R, Kippen R, Manning T, et al. Stress urinary incontinence in the mesh complication era: current Australian trends. BJU Int 2021;128:95-102.
- Jia G, Jiang C, Wang K, et al. Epidemiological investigation of urinary incontinence in peri- and postpartum women from Nanjing, China. Low Urin Tract Symptoms 2021;13:481-9.
- Schreiber Pedersen L, Lose G, Høybye MT, et al. Prevalence of urinary incontinence among women and analysis of potential risk factors in Germany and Denmark. Acta Obstet Gynecol Scand 2017;96:939-48.
- Svenningsen R, Oversand SH, Schiøtz HA, et al. Comparing risk of repeat surgery for stress urinary incontinence after mid-urethral slings and polyacrylamide hydrogel. Acta Obstet Gynecol Scand 2021;100:2186-92.
- Kim MM, Kreydin EI. The Association of Serum Testosterone Levels and Urinary Incontinence in Women. J Urol 2018;199:522-7.

**Cite this article as:** Wei D, Meng J, Zhang Y, Chen Y, Li J, Niu X. Identification of potential associated factors for stress urinary incontinence in women: a retrospective study. Ann Transl Med 2022;10(18):965. doi: 10.21037/atm-22-3539

- Kołodyńska G, Zalewski M, Rożek-Piechura K. Urinary incontinence in postmenopausal women - causes, symptoms, treatment. Prz Menopauzalny 2019;18:46-50.
- 22. Gao J, Liu X, Zuo Y, et al. Risk factors of postpartum stress urinary incontinence in primiparas: What should we care. Medicine (Baltimore) 2021;100:e25796.
- 23. Fuselier A, Hanberry J, Margaret Lovin J, et al. Obesity and Stress Urinary Incontinence: Impact on Pathophysiology and Treatment. Curr Urol Rep 2018;19:10.
- 24. Brož J, Hronová M, Brunerová L. Metabolic syndrome and diabetes mellitus in women with and without stress urinary incontinence. Int Urogynecol J 2019;30:847.
- Ströher RLM, Sartori MGF, Takano CC, et al. Metabolic syndrome in women with and without stress urinary incontinence. Int Urogynecol J 2020;31:173-9.
- Vašek P, Gärtner M, Szabová O, et al. Urinary incontinence during pregnancy. Ceska Gynekol 2019;84:73-6.
- 27. Kwon J, Suzuki T, Takaoka EI, et al. Analysis of continence reflexes by dynamic urethral pressure recordings in a rat stress urinary incontinence model induced by multiple simulated birth traumas. Am J Physiol Renal Physiol 2019;317:F781-8.
- Sangsawang B, Sangsawang N. Stress urinary incontinence in pregnant women: a review of prevalence, pathophysiology, and treatment. Int Urogynecol J 2013;24:901-12.
- Wingate JT, Erickson BA, Murphy G, et al. Multicenter Analysis of Patient Reported Outcomes Following Artificial Urinary Sphincter Placement for Male Stress Urinary Incontinence. J Urol 2018;199:785-90.
- Bergman I, Westergren Söderberg M, Lundqvist A, et al. Associations Between Childbirth and Urinary Incontinence After Midurethral Sling Surgery. Obstet Gynecol 2018;131:297-303.
- Sánchez-García O, Hernández-Aragón LG, López-García K, et al. Signs of damage in pelvic floor muscles at the end of pregnancy in rabbits. Int Urogynecol J 2019;30:977-84.

(English Language Editor: L. Huleatt)