



Cellular responses and functions of $\alpha 7$ nicotinic acetylcholine receptor activation in the brain: a narrative review

Zhe-Qi Xu^{1,2^}, Wen-Jun Zhang³, Ding-Feng Su¹, Guo-Qing Zhang², Chao-Yu Miao¹

¹Department of Pharmacology, Second Military Medical University/Naval Medical University, Shanghai, China; ²Department of Pharmacy, Eastern Hepatobiliary Surgery Hospital, Second Military Medical University/Naval Medical University, Shanghai, China; ³Department of Neurology, Dongying People's Hospital, Dongying, China

Contributions: (I) Conception and design: CY Miao, GQ Zhang; (II) Administrative support: CY Miao, GQ Zhang; (III) Provision of study materials or patients: None; (IV) Collection and assembly of data: None; (V) Data analysis and interpretation: None; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Chao-Yu Miao. Department of Pharmacology, Second Military Medical University/Naval Medical University, Shanghai 200433, China. Email: cymiao@smmu.edu.cn; Guo-Qing Zhang. Department of Pharmacy, Eastern Hepatobiliary Surgery Hospital, Second Military Medical University/Naval Medical University, Shanghai 200433, China. Email: gqzhang@smmu.edu.cn.

Abstract: The $\alpha 7$ nicotinic acetylcholine receptor ($\alpha 7$ nAChR) has been studied for many years since its discovery. Although many functions and characteristics of brain $\alpha 7$ nAChR are widely understood, much remains to be elucidated. The $\alpha 7$ nAChR is widely expressed in the central nervous system, not only in neurons but also in astrocytes, microglia, and endothelial cells. $\alpha 7$ nAChR can be activated by endogenous agonist like acetylcholine or exogenous agonists like nicotine and PNU282987. Its agonists can be divided into selective agonists and non-selective agonists. The activation of $\alpha 7$ nAChR results in a series of physiological processes which have both short-term and long-term effects on cells, for example, calcium influx, neurotransmitter release, synaptic plasticity, and excitatory transmission. It also induces other downstream events, such as inflammation, autophagy, necrosis, transcription, and apoptosis. The cellular responses to $\alpha 7$ nAChR activation vary according to cell types and conditions. For example, $\alpha 7$ nAChR activation in pyramidal neurons leads to long-term potentiation, while $\alpha 7$ nAChR activation in GABAergic interneurons leads to long-term depression. Studies have also shown some contradictory phenomena, which requires further study for clarification. Herein, the cellular responses of $\alpha 7$ nAChR activation are summarized, and the functions of $\alpha 7$ nAChR in neurons and non-neuronal cells are discussed. We also summarized contradictory conclusions to show where we stand and where to go for future studies.

Keywords: $\alpha 7$ nicotinic acetylcholine receptor ($\alpha 7$ nAChR); signaling pathway; central nervous system diseases (CNS diseases); calcium influx; synaptic plasticity

Submitted Dec 10, 2020. Accepted for publication Feb 24, 2021.

doi: 10.21037/atm-21-273

View this article at: <http://dx.doi.org/10.21037/atm-21-273>

Introduction

Nicotinic acetylcholine receptors (nAChRs) are cholinergic receptors which are activated by the endogenous neurotransmitter acetylcholine (1). Distinct from muscarinic acetylcholine receptors, nAChRs can also

respond to nicotine, belonging to the cys-loop superfamily of receptors (2,3). The nAChR is a pentameric ligand-gated ion channel containing 5 homomeric or heteromeric subunits (4). In total, 9 subunits have been found, and are named as $\alpha 2$ -7 and $\beta 2$ -4 (2,5,6). These 9 subunits can form different types of nAChRs, for example ($\alpha 4$)₃($\beta 2$)₂ nAChR,

[^] ORCID: 0000-0002-6508-1038.




Figure 1 Cellular responses to $\alpha 7$ nAChR activation. The activation of $\alpha 7$ nAChR can lead to calcium influx directly, which can trigger neurotransmitter release, action potentials, synaptic plasticity, and calcium-induced calcium release. It can also activate the JAK2-STAT3 signaling pathway and further induce anti-apoptotic and anti-inflammatory effects. In addition, it also activates the AMPK-mTOR signaling pathway and induces autophagy. $\alpha 7$ nAChR, $\alpha 7$ nicotinic acetylcholine receptor.

$(\alpha 4)_2(\beta 3)$, nAChR, and $(\alpha 3)_2(\beta 4)$, nAChR. The $\alpha 7$ nicotinic acetylcholine receptor ($\alpha 7$ nAChR) is a special subtype of nAChR which consists of 5 homometric $\alpha 7$ subunits.

The $\alpha 7$ nAChR is widely expressed in the central nervous system (CNS), not only in neurons but also in astrocytes, microglia, and endothelial cells (7-9). It is one of the most well-studied ionotropic receptors. In neurons, it is believed that $\alpha 7$ nAChR plays an important role in synaptic plasticity, contributing to dependence, learning, memory, and other cognitive functions (10,11). The activation of $\alpha 7$ nAChR can also contribute to neurotransmitter release, calcium influx, neuronal excitability, and other cellular activities (12-15). Studies have also revealed that the activation of $\alpha 7$ nAChR in microglia can induce anti-inflammatory effects (16,17). As several benefits have been found, $\alpha 7$ nAChR is considered a promising therapeutic target in CNS diseases, such as psychiatric disorders including autism and schizophrenia, neurodegenerative diseases including Alzheimer's disease (AD) and Parkinson's disease (PD), and

cerebrovascular diseases including hemorrhage stroke and ischemic stroke (18-21). Although, there were many reviews about $\alpha 7$ nAChR in the brain, many of them were focused on certain disease or were out of date. Some even reported contradictory conclusions. In this review, we will summarize recent researches on $\alpha 7$ nAChR in the brain and give an update snapshot of where we stand and where to head for in area of $\alpha 7$ nAChR. This review will summarize the cellular responses to $\alpha 7$ nAChR activation and its different roles in various types of brain cells. We present the following article in accordance with the Narrative Review reporting checklist (available at <http://dx.doi.org/10.21037/atm-21-273>).

Cellular responses to $\alpha 7$ nAChR activation

As a ligand-gated ion channel, the activation of $\alpha 7$ nAChR can lead to a series of cellular responses, including but not limited to calcium influx, neurotransmitter release, synaptic plasticity, and excitatory transmission (12-15). It also induces other downstream events, such as inflammation, autophagy, necrosis, transcription, and apoptosis (Figure 1).

$\alpha 7$ nAChR and calcium influx

When the agonist binds to the ligand-binding domain of $\alpha 7$ nAChR, the channel, which is composed of 5 $\alpha 7$ subunits, opens to let cations through. The $\alpha 7$ nAChR has a high permeability to calcium (22). To determine this important physiological parameter, early studies used heterologous systems and reported permeability ratios of $P_{\text{Ca}}/P_{\text{Na}}$ as high as 15 to 20 (3,23). However, Castro and Albuquerque used ion activities and the Goldman-Hodgkin-Katz equation for reversal potentials shifts to calculate the permeability ratios of Ca^{2+} in cultured rat hippocampal neurons. The results showed that the permeability ratio of $P_{\text{Ca}}/P_{\text{Na}}$ was 6.1 ± 0.5 , which was much lower than the results determined by heterologous systems (24). The difference in permeability may be due to the different conditions. Another method to detect the Ca^{2+} permeability of $\alpha 7$ nAChR are fluorescent Ca^{2+} indicators (25,26). This method records fluorescence signals and transmembrane currents simultaneously, and maintains good control of cell voltage and excludes any form of Ca^{2+} -induced Ca^{2+} release (CICR) (25). The result is displayed as fractional calcium current (P_f), indicating the percentage of the total current flowing through an ion channel which is carried by Ca^{2+} . A study showed that the P_f value of human neuronal $\alpha 7$ nAChR was 11.6%, indicating $\alpha 7$ nAChR has a high calcium permeability (27). The

activation of $\alpha 7$ nAChR not only raises cytoplasmic Ca^{2+} directly, but also indirectly. For example, the activation of $\alpha 7$ nAChR can depolarize neurons, and then induce calcium influx indirectly by voltage-dependent calcium channels (VDCCs) (28). Furthermore, $\alpha 7$ nAChR-mediated rising cytoplasmic calcium is partly caused by CICR from the endoplasmic reticulum (29). These are the 3 main ways that $\alpha 7$ nAChR can induce increased cytoplasmic Ca^{2+} .

$\alpha 7$ nAChR and neurotransmitter release

Neurotransmitter release usually occurs in the presynaptic membrane (30). The main processes of synaptic vesicle exocytosis and neurotransmitter release include: (I) the action potential arrives at the presynaptic terminals and depolarizes the terminal membrane; (II) the depolarized membrane induces calcium influx through VDCCs which dramatically increases the calcium concentration in the cytoplasm near the presynaptic membrane; (III) Ca^{2+} entry contributes to conformational changes in specific proteins in synaptic vesicles, leading to the fusion of synaptic vesicles to the presynaptic membrane. The whole process is an instantaneous response to the action potential, which lasts only a millisecond (31). Although neurotransmitter release is mainly triggered by VDCCs, in certain circumstances, $\alpha 7$ nAChR-mediated Ca^{2+} influx is sufficient to induce synaptic vesicles to release neurotransmitters. Gray *et al.* reported that nicotine acts on presynaptic $\alpha 7$ nAChR to enhance glutamatergic transmission in the hippocampus (13). It was also shown that activation of $\alpha 7$ nAChR in the presynaptic terminals of GABAergic neurons could increase GABA input to 5-HT dorsal raphe nucleus (DRN) neurons and lead to inhibitory postsynaptic spontaneous currents (32). This effect was triggered by $\alpha 7$ nAChR-mediated CICR in DRN GABAergic terminals.

However, neurotransmitter release is not only modulated by $\alpha 7$ nAChR-mediated calcium changes, but is also regulated by $\alpha 7$ nAChR-related signal transduction. Evidence has shown that the release of dopamine could be enhanced by the activation of $\alpha 7$ nAChR, which was modulated through protein kinase C (33,34).

$\alpha 7$ nAChR and synaptic plasticity

Synaptic plasticity, such as short-term potentiation (STP), long-term potentiation (LTP), or long-term depression (LTD), is a neuronal property characterized by the ability of synapses to strengthen or weaken in response to their

increased or decreased activity (35-37). Accumulating evidence suggests that it is one of the most important neurological foundations of learning and memory (38-40). Researchers found that in CA1 pyramidal neurons of the hippocampus, properly timed nicotine-mediated nAChR activation could contribute to the induction of LTP via presynaptic and postsynaptic pathways (41). The results showed that $\alpha 7$ nAChR contributed to most of these effects (41). However, different kinds of neurons had different outcomes. For GABAergic interneurons, the excitation caused inhibition of nearby pyramidal neurons, and the activation of $\alpha 7$ nAChR-mediated synaptic plasticity led to LTD (41). This synaptic plasticity is associated with $\alpha 7$ nAChR-mediated neurotransmitter release at the presynaptic membrane as well as $\alpha 7$ nAChR-mediated postsynaptic depolarization and calcium signals (41). Nitta *et al.* reported that infusion of β -amyloid protein into cerebral ventricles for 14 days could lead to impairments in learning and cholinergic neuronal degeneration (42). Furthermore, Chen *et al.* reported a series of impairments in CA1 hippocampal slices, including basal synaptic transmission, LTP, post-tetanic potentiation (PTP), and paired-pulse facilitation (PPF) after continuous intracerebroventricular infusion of β -amyloid protein (43). They also found that activation of $\alpha 7$ nAChR was required for LTP induction, highlighting the association between $\alpha 7$ nAChR and synaptic plasticity (43). In addition to learning and memory, synaptic plasticity also participates in neuronal connections. Crisculo *et al.* reported that they found impaired synaptic plasticity in the visual cortex of $\alpha 7$ nAChR knockout mice, indicating that $\alpha 7$ nAChR contributes to visual cortex function (44).

$\alpha 7$ nAChR and anti-inflammatory effects

The relationship between the nervous system and the immune system has been studied for several decades, and it has been consistently demonstrated that these 2 systems interact with each other. Researchers found that stimulation of the vagus nerve could attenuate inflammation by inhibiting macrophage tumor-necrosis factor (TNF) release (45). Further results suggested that this phenomenon was mediated by $\alpha 7$ nAChR responding to acetylcholine (45-47). Hence, the anti-inflammatory effect of the vagal nerve is termed the “cholinergic anti-inflammatory pathway (CAP)” (45,46).

The $\alpha 7$ nAChR is widely expressed in cells of the immune system, such as macrophages, monocytes, and lymphocytes

(48-50). Pinheiro *et al.* reported that treatment with PNU282987, an $\alpha 7$ nAChR agonist, could reduce LPS-induced acute lung injury through reducing neutrophil recruitment and IL-1 β , TNF- α , and IL-6 levels via changing the macrophage profile (51). Our lab also found that combination treatment with anisodamine (antagonist of muscarinic acetylcholine receptors) and neostigmine (cholinesterase inhibitor), which promoted the binding of endogenous acetylcholine to $\alpha 7$ nAChR, could rescue crush syndrome by reducing inflammation, and this phenomenon was absent in $\alpha 7$ nAChR knockout mice (52). Several studies have revealed that the attenuation of inflammatory cytokine release from macrophages by activating $\alpha 7$ nAChR was through the JAK2/STAT3 signaling pathway (16,52-54).

In addition to the roles of $\alpha 7$ nAChR in the peripheral CAP, many studies have investigated whether such roles exist in the CNS. In the brain, microglia are the resident immune cells, and the activation of these cells accounts for neuroinflammation (55,56). Shytle *et al.* reported for the first time that the brain CAP regulated microglial activation through $\alpha 7$ nAChR, and pretreatment with nicotine could reduce LPS-induced TNF- α release (47). Furthermore, De Simone *et al.* also demonstrated the existence of the CAP in the brain and the expression of $\alpha 7$ nAChR in microglia, and found that nicotine enhanced the expression of cyclooxygenase-2 and the synthesis of one of its major products, prostaglandin E2 (12). Our lab also demonstrated *in vivo* that activation of microglial $\alpha 7$ nAChR by PNU282987 reduced the production of IL-6, IL-1 β , IL-18, and TNF- α , thereby alleviating experimental autoimmune encephalomyelitis (57,58).

$\alpha 7$ nAChR and autophagy

Autophagy or “self-eating” is, as its name suggests, a basic cellular process that degrades and recycles misfolded or long-lived proteins and damaged organelles (59). Evidence has shown that autophagy is involved in the pathophysiological changes in many CNS diseases (60-65). Hung *et al.* found that $\alpha 7$ nAChR could bind with amyloid β (A β) and internalize into the cytoplasm to further inhibit A β -induced neurotoxicity through autophagy (66). They further demonstrated *in vivo* and *in vitro* that overexpression of LC3 led to neuroprotective effects by increasing $\alpha 7$ nAChR expression, which increased binding to A β and enhanced autophagy activity (67). Jeong *et al.* also found that $\alpha 7$ nAChR could be modulated and upregulated by melatonin, which enhanced autophagy and

exerted neuroprotective effects on prion diseases (68). Our lab has focused on $\alpha 7$ nAChR for many years and found that activation of $\alpha 7$ nAChR by PNU282987 enhanced microglia autophagy and suppressed neuroinflammation in experimental autoimmune encephalomyelitis mice (57). Enhanced autophagy induced by $\alpha 7$ nAChR activation was mediated by AMPK-mTOR-P70S6K signaling pathway activation (57). Interestingly, Hou *et al.* reported that autophagy was inhibited by $\alpha 7$ nAChR activation using PNU282987 and induced protection against myocardial ischemia/reperfusion injury *in vivo* and hypoxia/reoxygenation injury in primary cardiomyocytes *in vitro* (69). They demonstrated that these effects were mediated by JAK2 and PI3K activation in cardiomyocytes (69). These different phenomena in microglia and cardiomyocytes indicate that the relationship between autophagy and $\alpha 7$ nAChR activation varies in different organs or cells (70).

$\alpha 7$ nAChR and anti-apoptotic effects

It has been demonstrated that activation of $\alpha 7$ nAChR can affect cellular apoptosis (71-73). De Rosa *et al.* reported that nicotine upregulated $\alpha 7$ nAChR in human lymphocytes, which then decreased cortisol-induced apoptosis (71). However, the authors did not address the underlying mechanisms of this finding. Furthermore, Hejmadi *et al.* found that the presence of nicotine could protect primary cortical neurons against apoptosis induced by oxygen deprivation, which was mediated by 2 subtypes of nAChR, including $\alpha 7$ nAChR and $\beta 2$ nAChR (74). Further studies have revealed the mechanisms of nicotine-mediated anti-apoptotic effects. Toborek *et al.* reported that the anti-apoptotic effect of $\alpha 7$ nAChR activation in spinal cord neurons was dependent on increased phosphorylated ERK1/2 and total ERK1/2 activity, indicating that the ERK1/2 signaling pathway was involved in the anti-apoptotic effects of nicotine (75). It is interesting that different signaling pathways involved in $\alpha 7$ nAChR-mediated anti-apoptotic effects have been found by researchers. Parada *et al.* found that PNU282987 could rescue rotenone and oligomycin A-induced apoptosis via the JAK2/PI3K/Akt cascade in SH-SY5Y cells (72). In addition, Marrero and Bencherif reported that nicotine induced anti-apoptotic effects targeting $\alpha 7$ nAChR mainly through JAK2 activation, increased NF- κ B, and the production of Bcl-2 in PC12 cells (73). It is interesting that different signaling pathways have been found to be involved in the same phenomenon. This may be due to the different cells and different stimuli

used in studies.

Functions of $\alpha 7$ nAChR in neurons and non-neuronal cells

$\alpha 7$ nAChR in neurons

The $\alpha 7$ nAChR is widely expressed in neurons. It is also found in different locations in neurons, including the presynaptic membrane, postsynaptic membrane, and perisynaptic sites (76,77). The activation of $\alpha 7$ nAChR in neurons mainly leads to calcium-dependent events and a variety of other downstream signaling events which depend on receptor location. In general, activation of $\alpha 7$ nAChR leads to calcium influx which increases Ca^{2+} levels in neurons, triggering downstream responses (13,78,79). The basic neuronal physiological functions cannot be achieved without $\alpha 7$ nAChR. Activation of $\alpha 7$ nAChR in the presynaptic membrane leads to transmitter release, which contributes to synaptic transmission. Also, the effect of transmitter release induced by $\alpha 7$ nAChR does not need other forms of calcium influx like VDCCs or CICR (13). Postsynaptic $\alpha 7$ nAChR activation participates in intracellular signaling cascades and downstream processes. The most important function is to influence synaptic plasticity related to learning, memory, and addiction (80-82). As discussed above, properly-timed activation of $\alpha 7$ nAChR produces LTP or LTD, which remodels and strengthens connections within neuronal networks.

Furthermore, many studies have reported that $\alpha 7$ nAChR activation can have anti-apoptotic effects on neurons (83-85). A series of signaling pathways involved in the anti-apoptotic effects mediated by $\alpha 7$ nAChR have been uncovered. Huang *et al.* reported that neuroprotection by nicotine against colchicine-induced neuronal apoptosis was mediated by the PI3K/Akt signaling pathway (85). Also, Yu *et al.* demonstrated that $\alpha 7$ nAChR activation reduced β -amyloid-induced neuronal apoptosis through the PI3K signaling pathway (86). However, Chen *et al.* reported that the $\alpha 7$ nAChR agonist PHA568487 inhibited neuronal apoptosis through the TLR/Myd88/NF- κ B signaling pathway (83).

Moreover, as mentioned above, Hung *et al.* reported that $\alpha 7$ nAChR could induce A β transport into the cytoplasm and promote degradation through autophagy in SH-SY5Y cells (66). Jeong and Park also indicated that $\alpha 7$ nAChR upregulated autophagy in SH-SY5Y cells and protected against prion-mediated mitochondrial neurotoxicity (68).

In summary, $\alpha 7$ nAChR activation not only contributes to instantaneous processes such as calcium influx and neurotransmitter release, which comprise basic neuronal functions, it also participates in long-term effects like synaptic plasticity, and influences intracellular signaling pathways to produce anti-apoptotic effects and increased autophagy.

$\alpha 7$ nAChR in astrocytes

Astrocytes are star-shaped glial cells in the CNS. According to the research, astrocytes in the brain account for 20% to 40% of all glial cells. For a long time, scientists regarded astrocytes as merely support cells for neurons. However, many important functions have been recently uncovered, including but not limited to the regulation of neurotransmitters, the maintenance of the blood-brain barrier (BBB), the formation of the tripartite synapse, metabolic support, glucose sensing, the regulation of ion concentration in the extracellular space, and the repair of CNS injury through glial scar formation (87-95).

The $\alpha 7$ nAChR is also distributed in astrocytes and its activation can influence cellular functions (96). Teaktong *et al.* reported that the expression of $\alpha 7$ nAChR in astrocytes was increased in AD compared to that in dementia with Lewy bodies, which might contribute to alterations in calcium homeostasis and nitric oxide production in response to A β -mediated inflammation in AD (97). Yu *et al.* also found a significant increase of $\alpha 7$ nAChR expression in astrocytes in patients with AD or in patients carrying the Swedish APP 670/671 mutation, however, it was more pronounced in patients carrying the Swedish APP 670/671 mutation (98). They further indicated that the increased expression of $\alpha 7$ nAChR in astrocytes was positively correlated with the extent of pathological changes, especially the number of neuritic plaques in AD patients, which indicates its important role in the pathogenesis of AD (98). Although these studies demonstrated the relationship between $\alpha 7$ nAChR in astrocytes and CNS diseases, they did not describe the exact roles of $\alpha 7$ nAChR in these diseases. Later, Liu *et al.* reported the protective effect of $\alpha 7$ nAChR against dopaminergic neuron loss in a PD model via inhibition of astrocyte activation (99). In contrast to this finding, Talantova *et al.* reported that A β induced astrocytic glutamate release through $\alpha 7$ nAChR, which activated extrasynaptic NMDA receptors on neurons and damaged synapses in AD (100). Taken together, $\alpha 7$ nAChR activation in astrocytes can lead to different

effects under different conditions, which may be dependent on various ligands or brain regions.

$\alpha 7$ nAChR in microglia

Microglia are the brain's resident immune cells, which contribute to CNS inflammation. Decades ago, Wang *et al.* demonstrated that $\alpha 7$ nAChR was essential for acetylcholine inhibition of macrophage TNF- α release, which was named CAP (45). Whether its counterpart microglia also had this effect was a mystery until Shytle and colleagues uncovered it in their study (47). For the first time, Shytle *et al.* indicated that $\alpha 7$ nAChR was expressed in primary cultured microglial cells and brain slices (47). In addition, pretreatment with acetylcholine and nicotine inhibited LPS-induced TNF- α release in microglial cells, and this effect could be attenuated by α -bungarotoxin, a $\alpha 7$ nAChR antagonist (47). Many other studies have also demonstrated this effect (101-107). However, Thomsen *et al.* found that blockade of $\alpha 7$ nAChR by the antagonists methyllycaconitine (MLA) and NS6740 reduced LPS-induced microglia TNF- α release, which indicated that the $\alpha 7$ nAChR-induced anti-inflammatory effects of microglia were not due to calcium influx through ion channels (108).

Furthermore, Suzuki *et al.* also reported that the nicotine-induced increase in microglia intracellular Ca^{2+} levels was independent of extracellular Ca^{2+} , and this effect could be blocked by the phospholipase C (PLC) inhibitor U73122 and the inositol trisphosphate (IP_3) receptor blocker xestospongine C (101). In addition, nicotine-modulated LPS-mediated TNF- α release was also blocked by U73122 and xestospongine C (101). They suggested that nicotine triggered intracellular Ca^{2+} increases in microglia and their anti-inflammatory effects, which were mediated through the activation of PLC and Ca^{2+} release from the endoplasmic reticulum (101). Other signaling pathways involved in $\alpha 7$ nAChR-mediated microglial anti-inflammatory effects have also been uncovered. Parada *et al.* discovered a new signaling pathway involved in $\alpha 7$ nAChR-mediated neuronal protection in ischemic stroke (104). They suggested that activation of $\alpha 7$ nAChR in microglia contributed to the CAP through nuclear factor erythroid-2-related factor 2 (Nrf2) and heme oxygenase-1 (HO-1) activation, leading to neuroprotective effects in ischemic stroke (104). Morioka *et al.* also suggested a novel mechanism of microglial $\alpha 7$ nAChR-mediated neuroprotection beyond anti-inflammatory effects (109). They reported that treatment of microglia with nicotine significantly increased the

expression of the glutamate/aspartate transporter (GLAST), which is the main transporter in microglia, and increased glutamate uptake (109). The upregulation of GLAST was mainly through the IP_3 and Ca^{2+} /calmodulin-dependent protein kinase II (CaMKII) signaling pathway (109).

$\alpha 7$ nAChR in cerebrovascular endothelial cells

It was first reported in 2001 by Wang *et al.* that $\alpha 7$ nAChR was expressed in human bronchial epithelial and endothelial cells (110). They used both western blot and immunohistochemistry with an anti- $\alpha 7$ nAChR antibody and demonstrated that $\alpha 7$ nAChR was expressed in human aortic endothelial cells (110). They also used patch-clamp and found that the presence of fast desensitization currents activated by acetylcholine and nicotine could be blocked by MLA and α -bungarotoxin, thus further demonstrating the presence of $\alpha 7$ nAChR in endothelial cells (110). In spite of this discovery, they still did not know the exact role of $\alpha 7$ nAChR. In 2006, Li and Wang reported that the $\alpha 7$ nAChR agonist choline increased $\alpha 7$ nAChR expression, proliferation, and tube formation in human umbilical vein endothelial cells (HUVECs), and this phenomenon could be reversed by applying the antagonist α -bungarotoxin *in vitro* (111). They also investigated capillary density in a myocardial infarction (MI) rat model and reported that an $\alpha 7$ nAChR agonist increased capillary density in infarct tissues while the antagonists reversed this effect (111). This study indicated that activation of $\alpha 7$ nAChR could contribute to angiogenesis.

Ng *et al.* investigated the relationship between $\alpha 7$ nAChR and angiogenesis (112). They found that nicotine could dose-dependently enhance human microvascular endothelial cell (HMVEC) migration, which is a key process in angiogenesis (112). Furthermore, the induction of angiogenesis by other growth factors, including fibroblast growth factor (FGF) and vascular endothelial growth factor (VEGF), also relied on the activation of $\alpha 7$ nAChR, as their effect was abolished or attenuated after treatment with the $\alpha 7$ nAChR antagonist α -bungarotoxin (112). Wu *et al.* also detected the mRNA expression of $\alpha 7$ nAChR by RT-PCR in a series of human endothelial cell types: human pulmonary arterial endothelial cells (HPAEC), HMVECs, HUVECs, and human retinal endothelial cells (HREC) (113). Consistent with previous studies, they found that $\alpha 7$ nAChR was expressed in these cell types, however, there were differences in the levels of expression (113). Higher

Table 1 Studies show contradictory findings

Related functions	Author/year	Cell type/region	Primary findings	Species
$\alpha 7$ nAChR and synaptic plasticity	Ji D <i>et al.</i> (2001)	GABAergic interneuron	Activation of $\alpha 7$ nAChR by Ach can mediate synaptic plasticity leads to LTD, which can be blocked by methyllycaconitine (MLA)	Mouse
	Chen L <i>et al.</i> (2006)	Brain hippocampal slice	Activation of $\alpha 7$ nAChR by [3-(2,4-dimethoxybenzylidene)-anabaseine] (DMXB) can induce LTP which can be blocked by MLA or alpha-bungarotoxin (alpha-BTX)	Rat
$\alpha 7$ nAChR and autophagy	Hou Z <i>et al.</i> (2018)	Cardiomyocyte	Activation of $\alpha 7$ nAChR by PNU282987 inhibits Beclin-1 associated autophagy and reduces cardiomyocyte injury induced by hypoxia/reoxygenation	Rat
	Shao BZ <i>et al.</i> (2017)	Microglia	Activation of $\alpha 7$ nAChR by PNU282987 enhanced microglia autophagy and suppressed neuroinflammation	Mouse
$\alpha 7$ nAChR and anti-apoptosis effect	Parada E <i>et al.</i> (2017)	SH-SY5Y	PNU282987 can rescue rotenone and oligomycin A induced apoptosis via JAK2/PI3K/Akt cascade	Human
	Marrero and Bencherif (2009)	PC12	Activation of $\alpha 7$ nAChR induces anti-apoptosis effect through JAK2 activation and increasing NF- κ B and Bcl-2	Mouse
$\alpha 7$ nAChR in astrocytes	Liu Y <i>et al.</i> (2012)	Astrocyte and dopaminergic neuron	Activation of $\alpha 7$ nAChR has protective effect against dopaminergic neuron loss in PD model via inhibition of astrocyte activation	Mouse
	Talantova M <i>et al.</i> (2013)	Astrocyte and neuron	Amyloid- β binds to $\alpha 7$ nAChR to induce release of astrocytic glutamate, which activates NMDA receptors and damages synapse in AD	Rat, mouse, and human
$\alpha 7$ nAChR in cerebrovascular endothelial cells	Abbruscato TJ <i>et al.</i> (2002)	Brain microvessel endothelial cells	Nicotine and cotinine can increase permeability of blood-brain barrier under hypoxia condition and reduce expression of ZO-1, which can be reversed by $\alpha 7$ nAChR antagonist PHA543613	Bovine
	Kimura I <i>et al.</i> (2018)	Brain endothelial cells	PHA543613 can decrease permeability of blood-brain barrier and increase expression of claudin-5 by activating $\alpha 7$ nAChR	Rat

$\alpha 7$ nAChR, $\alpha 7$ nicotinic acetylcholine receptor; LTD, long-term depression; LTP, long-term potentiation; AD, Alzheimer's disease.

$\alpha 7$ nAChR expression was detected in HUVECs (113). They also reported that $\alpha 7$ nAChR plays a dominant role in nicotine-induced cell signaling and endothelial cell migration, proliferation, and tube formation (113).

In spite of the positive effects of $\alpha 7$ nAChR activation in these endothelial cells, studies in cerebrovascular endothelial cells showed contrasting results. Abbruscato and colleagues reported in 2002 that nicotine and its metabolite, cotinine, could increase the permeability of the BBB in a bovine brain microvessel endothelial cell model under hypoxic condition (114). Furthermore, nicotine and cotinine reduced the expression of ZO-1 (tight junction protein) which could be reversed by the $\alpha 7$ nAChR antagonist α -bungarotoxin, indicating that both the nicotine and cotinine-induced effects were mediated by the $\alpha 7$ nAChR (114). However, Kimura *et al.* reported different results in 2018. They demonstrated that the selective

$\alpha 7$ nAChR agonist PHA543613 decreased the permeability of the BBB and increased the expression of claudin-5 and occludin, which are key tight junction components (115). Additionally, they did not show a positive effect of nicotine. The different results may be due to the different agonists and different *in vitro* models used in studies.

Conclusions

Based on the discussion above, evidence from *in vivo* and *in vitro* studies has shown significant cellular responses to $\alpha 7$ nAChR activation. However, some results are contradictory and require further study (Table 1). For example, although $\alpha 7$ nAChR is a ligand-gated ion channel, it can also increase intracellular cAMP levels through adenylate cyclase 1 (AC1), which is a common signaling pathway of G protein coupled receptors (116). Does

$\alpha 7$ nAChR activate AC1 directly or indirectly? The cellular responses and functions of $\alpha 7$ nAChR in the brain show promising therapeutic effects in CNS diseases. The many benefits of $\alpha 7$ nAChR activation in the CNS may be the reason why smokers are at a lower risk of developing PD compared with non-smokers (117). As $\alpha 7$ nAChR has various effects on cells, agonists, antagonists, and other allosteric modulators are widely used in many animal models to study CNS diseases. Many agonists or allosteric modulators are involved in clinical trials to treat CNS diseases. However, whether these $\alpha 7$ nAChR targeting drugs have adverse effects or not should be studied in-depth.

Acknowledgments

Funding: None.

Footnote

Reporting Checklist: The authors have completed the Narrative Review reporting checklist. Available at <http://dx.doi.org/10.21037/atm-21-273>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/atm-21-273>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Gay EA, Yakel JL. Gating of nicotinic ACh receptors; new insights into structural transitions triggered by agonist binding that induce channel opening. *J Physiol* 2007;584:727-33.
2. Dani JA, Bertrand D. Nicotinic acetylcholine receptors and nicotinic cholinergic mechanisms of the central nervous system. *Annu Rev Pharmacol Toxicol* 2007;47:699-729.
3. Seguela P, Wadiche J, Dineley-Miller K, et al. Molecular cloning, functional properties, and distribution of rat brain alpha 7: a nicotinic cation channel highly permeable to calcium. *J Neurosci* 1993;13:596-604.
4. Brejc K, van Dijk WJ, Klaassen RV, et al. Crystal structure of an ACh-binding protein reveals the ligand-binding domain of nicotinic receptors. *Nature* 2001;411:269-76.
5. Yakel JL. Cholinergic receptors: functional role of nicotinic ACh receptors in brain circuits and disease. *Pflugers Arch* 2013;465:441-50.
6. Hurst R, Rollema H, Bertrand D. Nicotinic acetylcholine receptors: from basic science to therapeutics. *Pharmacol Ther* 2013;137:22-54.
7. Quik M, Polonskaya Y, Gillespie A, et al. Localization of nicotinic receptor subunit mRNAs in monkey brain by in situ hybridization. *J Comp Neurol* 2000;425:58-69.
8. Palma E, Conti L, Roseti C, et al. Novel approaches to study the involvement of alpha7-nAChR in human diseases. *Curr Drug Targets* 2012;13:579-86.
9. Dineley KT, Pandya AA, Yakel JL. Nicotinic ACh receptors as therapeutic targets in CNS disorders. *Trends Pharmacol Sci* 2015;36:96-108.
10. Nordman JC, Phillips WS, Kodama N, et al. Axon targeting of the alpha 7 nicotinic receptor in developing hippocampal neurons by Gprn1 regulates growth. *J Neurochem* 2014;129:649-62.
11. Zhao X, Kuryatov A, Lindstrom JM, et al. Nootropic drug modulation of neuronal nicotinic acetylcholine receptors in rat cortical neurons. *Mol Pharmacol* 2001;59:674-83.
12. De Simone R, Ajmone-Cat MA, Carnevale D, et al. Activation of alpha7 nicotinic acetylcholine receptor by nicotine selectively up-regulates cyclooxygenase-2 and prostaglandin E2 in rat microglial cultures. *J Neuroinflammation* 2005;2:4.
13. Gray R, Rajan AS, Radcliffe KA, et al. Hippocampal synaptic transmission enhanced by low concentrations of nicotine. *Nature* 1996;383:713-6.
14. Fucile S, Palma E, Mileo AM, et al. Human neuronal threonine-for-leucine-248 alpha 7 mutant nicotinic acetylcholine receptors are highly Ca²⁺ permeable. *Proc Natl Acad Sci U S A* 2000;97:3643-8.
15. Chamberlin NL, Bocchiaro CM, Greene RW, et al. Nicotinic excitation of rat hypoglossal motoneurons. *Neuroscience* 2002;115:861-70.

16. Shi S, Liang D, Bao M, et al. Gx-50 Inhibits Neuroinflammation via alpha7 nAChR Activation of the JAK2/STAT3 and PI3K/AKT Pathways. *J Alzheimers Dis* 2016;50:859-71.
17. Frasch MG, Szykaruk M, Prout AP, et al. Decreased neuroinflammation correlates to higher vagus nerve activity fluctuations in near-term ovine fetuses: a case for the afferent cholinergic anti-inflammatory pathway? *J Neuroinflammation* 2016;13:103.
18. Olincy A, Blakeley-Smith A, Johnson L, et al. Brief Report: Initial Trial of Alpha7-Nicotinic Receptor Stimulation in Two Adult Patients with Autism Spectrum Disorder. *J Autism Dev Disord* 2016;46:3812-7.
19. Haig G, Wang D, Othman AA, et al. The alpha7 Nicotinic Agonist ABT-126 in the Treatment of Cognitive Impairment Associated with Schizophrenia in Nonsmokers: Results from a Randomized Controlled Phase 2b Study. *Neuropsychopharmacology* 2016;41:2893-902.
20. Echeverria V, Yarkov A, Aliev G. Positive modulators of the alpha7 nicotinic receptor against neuroinflammation and cognitive impairment in Alzheimer's disease. *Prog Neurobiol* 2016;144:142-57.
21. Kawamata J, Suzuki S, Shimohama S. alpha7 nicotinic acetylcholine receptor mediated neuroprotection in Parkinson's disease. *Curr Drug Targets* 2012;13:623-30.
22. Barrantes GE, Westwick J, Wonnacott S. Nicotinic acetylcholine receptors in primary cultures of hippocampal neurons: pharmacology and Ca⁺⁺ permeability. *Biochem Soc Trans* 1994;22:294S.
23. Bertrand D, Galzi JL, Devillers-Thierry A, et al. Mutations at two distinct sites within the channel domain M2 alter calcium permeability of neuronal alpha 7 nicotinic receptor. *Proc Natl Acad Sci U S A* 1993;90:6971-5.
24. Castro NG, Albuquerque EX. alpha-Bungarotoxin-insensitive hippocampal nicotinic receptor channel has a high calcium permeability. *Biophys J* 1995;68:516-24.
25. Gryniewicz G, Poenie M, Tsien RY. A new generation of Ca²⁺ indicators with greatly improved fluorescence properties. *J Biol Chem* 1985;260:3440-50.
26. Neher E. The use of fura-2 for estimating Ca buffers and Ca fluxes. *Neuropharmacology* 1995;34:1423-42.
27. Fucile S, Renzi M, Lax P, et al. Fractional Ca(2+) current through human neuronal alpha7 nicotinic acetylcholine receptors. *Cell Calcium* 2003;34:205-9.
28. Rathouz MM, Berg DK. Synaptic-type acetylcholine receptors raise intracellular calcium levels in neurons by two mechanisms. *J Neurosci* 1994;14:6935-45.
29. Gueorguiev VD, Zeman RJ, Meyer EM, et al. Involvement of alpha7 nicotinic acetylcholine receptors in activation of tyrosine hydroxylase and dopamine beta-hydroxylase gene expression in PC12 cells. *J Neurochem* 2000;75:1997-2005.
30. Kavalali ET. The mechanisms and functions of spontaneous neurotransmitter release. *Nat Rev Neurosci* 2015;16:5-16.
31. Sudhof TC. Neurotransmitter release: the last millisecond in the life of a synaptic vesicle. *Neuron* 2013;80:675-90.
32. Hernandez-Vazquez F, Chavarria K, Garduno J, et al. Nicotine increases GABAergic input on rat dorsal raphe serotonergic neurons through alpha7 nicotinic acetylcholine receptor. *J Neurophysiol* 2014;112:3154-63.
33. Quarta D, Naylor CG, Barik J, et al. Drug discrimination and neurochemical studies in alpha7 null mutant mice: tests for the role of nicotinic alpha7 receptors in dopamine release. *Psychopharmacology (Berl)* 2009;203:399-410.
34. Soliakov L, Wonnacott S. Involvement of protein kinase C in the presynaptic nicotinic modulation of [(3)H]-dopamine release from rat striatal synaptosomes. *Br J Pharmacol* 2001;132:785-91.
35. Hughes JR. Post-tetanic potentiation. *Physiol Rev* 1958;38:91-113.
36. Bocchio M, Nabavi S, Capogna M. Synaptic Plasticity, Engrams, and Network Oscillations in Amygdala Circuits for Storage and Retrieval of Emotional Memories. *Neuron* 2017;94:731-43.
37. Basu J, Siegelbaum SA. The Corticohippocampal Circuit, Synaptic Plasticity, and Memory. *Cold Spring Harb Perspect Biol* 2015;7:a021733.
38. Martin SJ, Grimwood PD, Morris RG. Synaptic plasticity and memory: an evaluation of the hypothesis. *Annu Rev Neurosci* 2000;23:649-711.
39. Pignatelli M, Umanah GKE, Ribeiro SP, et al. Synaptic Plasticity onto Dopamine Neurons Shapes Fear Learning. *Neuron* 2017;93:425-40.
40. Mills F, Globa AK, Liu S, et al. Cadherins mediate cocaine-induced synaptic plasticity and behavioral conditioning. *Nat Neurosci* 2017;20:540-9.
41. Ji D, Lape R, Dani JA. Timing and location of nicotinic activity enhances or depresses hippocampal synaptic plasticity. *Neuron* 2001;31:131-41.
42. Nitta A, Itoh A, Hasegawa T, et al. beta-Amyloid protein-induced Alzheimer's disease animal model. *Neurosci Lett* 1994;170:63-6.
43. Chen L, Yamada K, Nabeshima T, et al. alpha7 Nicotinic acetylcholine receptor as a target to rescue deficit in hippocampal LTP induction in beta-amyloid infused rats.

- Neuropharmacology 2006;50:254-68.
44. Criscuolo C, Accorroni A, Domenici L, et al. Impaired synaptic plasticity in the visual cortex of mice lacking alpha7-nicotinic receptor subunit. *Neuroscience* 2015;294:166-71.
 45. Wang H, Yu M, Ochani M, et al. Nicotinic acetylcholine receptor alpha7 subunit is an essential regulator of inflammation. *Nature* 2003;421:384-8.
 46. Wang H, Liao H, Ochani M, et al. Cholinergic agonists inhibit HMGB1 release and improve survival in experimental sepsis. *Nat Med* 2004;10:1216-21.
 47. Shytle RD, Mori T, Townsend K, et al. Cholinergic modulation of microglial activation by alpha 7 nicotinic receptors. *J Neurochem* 2004;89:337-43.
 48. Zabrodskii PF, Maslyakov VV, Gromov MS. Role of alpha7-Nicotinic Acetylcholine Receptors of B Cells in the Immunotoxic Effect of Organophosphorus Compounds. *Bull Exp Biol Med* 2016;161:779-81.
 49. Zanetti SR, Ziblat A, Torres NI, et al. Expression and Functional Role of alpha7 Nicotinic Receptor in Human Cytokine-stimulated Natural Killer (NK) Cells. *J Biol Chem* 2016;291:16541-52.
 50. Liu Z, Han B, Li P, et al. Activation of alpha7nAChR by nicotine reduced the Th17 response in CD4(+)T lymphocytes. *Immunol Invest* 2014;43:667-74.
 51. Pinheiro NM, Santana FP, Almeida RR, et al. Acute lung injury is reduced by the alpha7nAChR agonist PNU-282987 through changes in the macrophage profile. *FASEB J* 2017;31:320-32.
 52. Xu ZQ, Shao BZ, Ke P, et al. Combined administration of anisodamine and neostigmine rescued acute lethal crush syndrome through alpha7nAChR-dependent JAK2-STAT3 signaling. *Sci Rep* 2016;6:37709.
 53. Lu XX, Hong ZQ, Tan Z, et al. Nicotinic Acetylcholine Receptor Alpha7 Subunit Mediates Vagus Nerve Stimulation-Induced Neuroprotection in Acute Permanent Cerebral Ischemia by a7nAChR/JAK2 Pathway. *Med Sci Monit* 2017;23:6072-81.
 54. Yang YH, Li DL, Bi XY, et al. Acetylcholine Inhibits LPS-Induced MMP-9 Production and Cell Migration via the alpha7 nAChR-JAK2/STAT3 Pathway in RAW264.7 Cells. *Cell Physiol Biochem* 2015;36:2025-38.
 55. Giulian D. Ameboid microglia as effectors of inflammation in the central nervous system. *J Neurosci Res* 1987;18:155-71,32-3.
 56. Lassmann H, Zimprich F, Rossler K, et al. Inflammation in the nervous system. Basic mechanisms and immunological concepts. *Rev Neurol (Paris)* 1991;147:763-81.
 57. Shao BZ, Ke P, Xu ZQ, et al. Autophagy Plays an Important Role in Anti-inflammatory Mechanisms Stimulated by Alpha7 Nicotinic Acetylcholine Receptor. *Front Immunol* 2017;8:553.
 58. Ke P, Shao BZ, Xu ZQ, et al. Activating alpha7 nicotinic acetylcholine receptor inhibits NLRP3 inflammasome through regulation of beta-arrestin-1. *CNS Neurosci Ther* 2017;23:875-84.
 59. Moulis M, Vindis C. Autophagy in Metabolic Age-Related Human Diseases. *Cells* 2018;7:149.
 60. Sun Y, Zhang T, Zhang Y, et al. Ischemic Postconditioning Alleviates Cerebral Ischemia-Reperfusion Injury Through Activating Autophagy During Early Reperfusion in Rats. *Neurochem Res* 2018;43:1826-40.
 61. Xu L, Zhang C, Jiang N, et al. Rapamycin combined with MCC950 to treat multiple sclerosis in experimental autoimmune encephalomyelitis. *J Cell Biochem* 2019;120:5160-8.
 62. Morandi E, Jagessar SA, Hart BA, et al. EBV Infection Empowers Human B Cells for Autoimmunity: Role of Autophagy and Relevance to Multiple Sclerosis. *J Immunol* 2017;199:435-48.
 63. Aminyavari S, Zahmatkesh M, Farahmandfar M, et al. Protective role of Apelin-13 on amyloid beta25-35-induced memory deficit; Involvement of autophagy and apoptosis process. *Prog Neuropsychopharmacol Biol Psychiatry* 2019;89:322-34.
 64. Hung COY, Livesey FJ. Altered gamma-Secretase Processing of APP Disrupts Lysosome and Autophagosome Function in Monogenic Alzheimer's Disease. *Cell Rep* 2018;25:3647-60.e2.
 65. Varmazyar R, Noori-Zadeh A, Rajaei F, et al. 17 beta-Estradiol Oxidative Stress Attenuation and Autophagy-Induced Dopaminergic Neuroprotection. *Cell J* 2019;21:1-6.
 66. Hung SY, Huang WP, Liou HC, et al. Autophagy protects neuron from Abeta-induced cytotoxicity. *Autophagy* 2009;5:502-10.
 67. Hung SY, Huang WP, Liou HC, et al. LC3 overexpression reduces Abeta neurotoxicity through increasing alpha7nAChR expression and autophagic activity in neurons and mice. *Neuropharmacology* 2015;93:243-51.
 68. Jeong JK, Park SY. Melatonin regulates the autophagic flux via activation of alpha-7 nicotinic acetylcholine receptors. *J Pineal Res* 2015;59:24-37.
 69. Hou Z, Zhou Y, Yang H, et al. Alpha7 nicotinic acetylcholine receptor activation protects against myocardial reperfusion injury through modulation

- of autophagy. *Biochem Biophys Res Commun* 2018;500:357-64.
70. Klionsky DJ, Abdelmohsen K, Abe A, et al. Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). *Autophagy* 2016;12:1-222.
 71. De Rosa MJ, Esandi Mdel C, Garelli A, et al. Relationship between alpha 7 nAChR and apoptosis in human lymphocytes. *J Neuroimmunol* 2005;160:154-61.
 72. Parada E, Egea J, Romero A, et al. Poststress treatment with PNU282987 can rescue SH-SY5Y cells undergoing apoptosis via alpha7 nicotinic receptors linked to a Jak2/Akt/HO-1 signaling pathway. *Free Radic Biol Med* 2010;49:1815-21.
 73. Marrero MB, Bencherif M. Convergence of alpha 7 nicotinic acetylcholine receptor-activated pathways for anti-apoptosis and anti-inflammation: central role for JAK2 activation of STAT3 and NF-kappaB. *Brain Res* 2009;1256:1-7.
 74. Hejmadi MV, Dajas-Bailador F, Barns SM, et al. Neuroprotection by nicotine against hypoxia-induced apoptosis in cortical cultures involves activation of multiple nicotinic acetylcholine receptor subtypes. *Mol Cell Neurosci* 2003;24:779-86.
 75. Toborek M, Son KW, Pudielko A, et al. ERK 1/2 signaling pathway is involved in nicotine-mediated neuroprotection in spinal cord neurons. *J Cell Biochem* 2007;100:279-92.
 76. Jones IW, Wonnacott S. Precise localization of alpha7 nicotinic acetylcholine receptors on glutamatergic axon terminals in the rat ventral tegmental area. *J Neurosci* 2004;24:11244-52.
 77. Marshall LM. Synaptic localization of alpha-bungarotoxin binding which blocks nicotinic transmission at frog sympathetic neurons. *Proc Natl Acad Sci U S A* 1981;78:1948-52.
 78. Shoop RD, Chang KT, Ellisman MH, et al. Synaptically driven calcium transients via nicotinic receptors on somatic spines. *J Neurosci* 2001;21:771-81.
 79. Vogt KE, Regehr WG. Cholinergic modulation of excitatory synaptic transmission in the CA3 area of the hippocampus. *J Neurosci* 2001;21:75-83.
 80. Sultan A, Yang KS, Isaev D, et al. Thujone inhibits the function of alpha7-nicotinic acetylcholine receptors and impairs nicotine-induced memory enhancement in one-trial passive avoidance paradigm. *Toxicology* 2017;384:23-32.
 81. Li J, Bo L, Zhang P, et al. Exposure to nicotine during pregnancy and altered learning and memory in the rat offspring. *Nicotine Tob Res* 2015;17:661-6.
 82. Wei XM, Yang W, Liu LX, et al. Effects of L-arginine and N(omega)-nitro-L-arginine methylester on learning and memory and alpha7 nAChR expression in the prefrontal cortex and hippocampus of rats. *Neurosci Bull* 2013;29:303-10.
 83. Chen K, Sun Y, Diao Y, et al. alpha7 nicotinic acetylcholine receptor agonist inhibits the damage of rat hippocampal neurons by TLR4/Myd88/NFkappaB signaling pathway during cardiopulmonary bypass. *Mol Med Rep* 2017;16:4770-6.
 84. Sun Y, Song D, Wang M, et al. alpha7 nicotinic acetylcholine receptor agonist attenuates the cerebral injury in a rat model of cardiopulmonary bypass by activating the Akt/GSK3beta pathway. *Mol Med Rep* 2017;16:7979-86.
 85. Huang X, Cheng Z, Su Q, et al. Neuroprotection by nicotine against colchicine-induced apoptosis is mediated by PI3-kinase--Akt pathways. *Int J Neurosci* 2012;122:324-32.
 86. Yu W, Mechawar N, Krantic S, et al. alpha7 Nicotinic receptor activation reduces beta-amyloid-induced apoptosis by inhibiting caspase-independent death through phosphatidylinositol 3-kinase signaling. *J Neurochem* 2011;119:848-58.
 87. Araque A, Parpura V, Sanzgiri RP, et al. Tripartite synapses: glia, the unacknowledged partner. *Trends Neurosci* 1999;22:208-15.
 88. Abbott NJ, Ronnback L, Hansson E. Astrocyte-endothelial interactions at the blood-brain barrier. *Nat Rev Neurosci* 2006;7:41-53.
 89. McDougal DH, Viard E, Hermann GE, et al. Astrocytes in the hindbrain detect glucoprivation and regulate gastric motility. *Auton Neurosci* 2013;175:61-9.
 90. Figley CR, Stroman PW. The role(s) of astrocytes and astrocyte activity in neurometabolism, neurovascular coupling, and the production of functional neuroimaging signals. *Eur J Neurosci* 2011;33:577-88.
 91. Santello M, Volterra A. Synaptic modulation by astrocytes via Ca2+-dependent glutamate release. *Neuroscience* 2009;158:253-9.
 92. Walz W. Role of astrocytes in the clearance of excess extracellular potassium. *Neurochem Int* 2000;36:291-300.
 93. Piet R, Vargova L, Sykova E, et al. Physiological contribution of the astrocytic environment of neurons to intersynaptic crosstalk. *Proc Natl Acad Sci U S A* 2004;101:2151-5.
 94. Anderson MA, Burda JE, Ren Y, et al. Astrocyte scar formation aids central nervous system axon regeneration.

- Nature 2016;532:195-200.
95. Patti L, Raiteri L, Grilli M, et al. Evidence that $\alpha 7$ nicotinic receptor modulates glutamate release from mouse neocortical gliosomes. *Neurochem Int* 2007;51:1-7.
 96. Grybko M, Sharma G, Vijayaraghavan S. Functional distribution of nicotinic receptors in CA3 region of the hippocampus. *J Mol Neurosci* 2010;40:114-20.
 97. Teaktong T, Graham A, Court J, et al. Alzheimer's disease is associated with a selective increase in $\alpha 7$ nicotinic acetylcholine receptor immunoreactivity in astrocytes. *Glia* 2003;41:207-11.
 98. Yu WF, Guan ZZ, Bogdanovic N, et al. High selective expression of $\alpha 7$ nicotinic receptors on astrocytes in the brains of patients with sporadic Alzheimer's disease and patients carrying Swedish APP 670/671 mutation: a possible association with neuritic plaques. *Exp Neurol* 2005;192:215-25.
 99. Liu Y, Hu J, Wu J, et al. $\alpha 7$ nicotinic acetylcholine receptor-mediated neuroprotection against dopaminergic neuron loss in an MPTP mouse model via inhibition of astrocyte activation. *J Neuroinflammation* 2012;9:98.
 100. Talantova M, Sanz-Blasco S, Zhang X, et al. A β induces astrocytic glutamate release, extrasynaptic NMDA receptor activation, and synaptic loss. *Proc Natl Acad Sci U S A* 2013;110:E2518-27.
 101. Suzuki T, Hide I, Matsubara A, et al. Microglial $\alpha 7$ nicotinic acetylcholine receptors drive a phospholipase C/IP3 pathway and modulate the cell activation toward a neuroprotective role. *J Neurosci Res* 2006;83:1461-70.
 102. Ohnishi M, Katsuki H, Takagi M, et al. Long-term treatment with nicotine suppresses neurotoxicity of, and microglial activation by, thrombin in cortico-striatal slice cultures. *Eur J Pharmacol* 2009;602:288-93.
 103. Mencil M, Nash M, Jacobson C. Neuregulin upregulates microglial $\alpha 7$ nicotinic acetylcholine receptor expression in immortalized cell lines: implications for regulating neuroinflammation. *PLoS One* 2013;8:e70338.
 104. Parada E, Egea J, Buendia I, et al. The microglial $\alpha 7$ -acetylcholine nicotinic receptor is a key element in promoting neuroprotection by inducing heme oxygenase-1 via nuclear factor erythroid-2-related factor 2. *Antioxid Redox Signal* 2013;19:1135-48.
 105. Hua S, Ek CJ, Mallard C, et al. Perinatal hypoxia-ischemia reduces $\alpha 7$ nicotinic receptor expression and selective $\alpha 7$ nicotinic receptor stimulation suppresses inflammation and promotes microglial Mox phenotype. *Biomed Res Int* 2014;2014:718769.
 106. Jiang Y, Li L, Liu B, et al. Vagus nerve stimulation attenuates cerebral ischemia and reperfusion injury via endogenous cholinergic pathway in rat. *PLoS One* 2014;9:e102342.
 107. Lutz JA, Kulshrestha M, Rogers DT, et al. A nicotinic receptor-mediated anti-inflammatory effect of the flavonoid rhamnetin in BV2 microglia. *Fitoterapia* 2014;98:11-21.
 108. Thomsen MS, Mikkelsen JD. The $\alpha 7$ nicotinic acetylcholine receptor ligands methyllycaconitine, NS6740 and GTS-21 reduce lipopolysaccharide-induced TNF- α release from microglia. *J Neuroimmunol* 2012;251:65-72.
 109. Morioka N, Tokuhara M, Nakamura Y, et al. Primary cultures of rat cortical microglia treated with nicotine increases in the expression of excitatory amino acid transporter 1 (GLAST) via the activation of the $\alpha 7$ nicotinic acetylcholine receptor. *Neuroscience* 2014;258:374-84.
 110. Wang Y, Pereira EF, Maus AD, et al. Human bronchial epithelial and endothelial cells express $\alpha 7$ nicotinic acetylcholine receptors. *Mol Pharmacol* 2001;60:1201-9.
 111. Li XW, Wang H. Non-neuronal nicotinic $\alpha 7$ receptor, a new endothelial target for revascularization. *Life Sci* 2006;78:1863-70.
 112. Ng MK, Wu J, Chang E, et al. A central role for nicotinic cholinergic regulation of growth factor-induced endothelial cell migration. *Arterioscler Thromb Vasc Biol* 2007;27:106-12.
 113. Wu JC, Chruscinski A, De Jesus Perez VA, et al. Cholinergic modulation of angiogenesis: role of the $\alpha 7$ nicotinic acetylcholine receptor. *J Cell Biochem* 2009;108:433-46.
 114. Abbruscato TJ, Lopez SP, Mark KS, et al. Nicotine and cotinine modulate cerebral microvascular permeability and protein expression of ZO-1 through nicotinic acetylcholine receptors expressed on brain endothelial cells. *J Pharm Sci* 2002;91:2525-38.
 115. Kimura I, Dohgu S, Takata F, et al. Activation of the $\alpha 7$ nicotinic acetylcholine receptor upregulates blood-brain barrier function through increased claudin-5 and

- occludin expression in rat brain endothelial cells. *Neurosci Lett* 2019;694:9-13.
116. Cheng Q, Yakel JL. Activation of alpha7 nicotinic acetylcholine receptors increases intracellular cAMP levels via activation of AC1 in hippocampal neurons.

- Neuropharmacology* 2015;95:405-14.
117. Gallo V, Vineis P, Cancellieri M, et al. Exploring causality of the association between smoking and Parkinson's disease. *Int J Epidemiol* 2019;48:912-25.

Cite this article as: Xu ZQ, Zhang WJ, Su DF, Zhang GQ, Miao CY. Cellular responses and functions of $\alpha 7$ nicotinic acetylcholine receptor activation in the brain. *Ann Transl Med* 2021;9(6):509. doi: 10.21037/atm-21-273