

Overcoming the hurdles of using PET/CT for target volume delineation in curative intent radiotherapy of non-small cell lung cancer

Leila Tchelebi, Hani Ashamalla

Department of Radiation Oncology, New York Methodist Hospital, Brooklyn, NY 11215, USA

Correspondence to: Hani Ashamalla, MD. Department of Radiation Oncology, New York Methodist Hospital, 506 Sixth Street, Brooklyn, NY 11215, USA. Email: haa9002@nyp.org.

Submitted Jul 01, 2015. Accepted for publication Jul 03, 2015.

doi:10.3978/j.issn.2305-5839.2015.07.02

View this article at: <http://dx.doi.org/10.3978/j.issn.2305-5839.2015.07.02>

PET/CT imaging has proven to be an invaluable tool for target volume delineation in radiation treatment planning of non-small cell lung cancer. However, as with any technological advancement, it has its limitations. In their recent review article published in *Radiotherapy and Oncology*, Konert *et al.* (1) eloquently summarize many of the challenges encountered by radiation oncologists when using PET/CT-based planning in radical intent radiotherapy. The authors go further to describe evidence-based recommendations to provide clear guidance on the use of PET/CT in target volume delineation in order to overcome some of these challenges and standardize clinical practice.

¹⁸F-fluorodeoxyglucose (FDG) positron emission tomography (PET) capitalizes on the increased uptake of glucose in metabolically active tissues, including malignant tumors. In FDG-PET imaging, a radioactive analog of glucose is injected into the patient. As it decays, photons are ultimately produced and detected by the PET scanner. When acquired with a CT scan, the combined PET/CT can be registered with the CT acquired during patient simulation to identify FDG-avid areas for inclusion in the treatment volume. In cases where it may be difficult to distinguish tumor from atelectasis or pulmonary edema, PET/CT fusion with a planning CT is of particular benefit. PET/CT also has the potential to decrease inter-observer variability in target volume delineation, as sighted by Ashamalla *et al.* and others (1,2). Indeed, the advantages of using PET/CT for target volume delineation in not only early stage NSCLC, but in locally advanced disease and for other body sites as well, has been widely published (3-6).

Despite its feasibility and widespread application in radiation treatment planning, PET/CT can be challenging to use and a standardized method of applying it to target volume delineation is needed.

The first set of challenges encountered when attempting to use a PET/CT to contour a target volume as identified by Konert *et al.* (1) is the acquisition of the image. PET/CT scans obtained for diagnostic purposes are usually acquired in a different position than the CT scans used for treatment planning, making an accurate registration of the two images difficult. The authors caution against the co-registration of diagnostic PET/CT with planning CTs. Instead, they suggest either obtaining a second PET/CT once distant metastatic disease is ruled out, or, requesting that the diagnostic PET/CT be obtained in the treatment planning position. In an era in which controlling costs is paramount and obtaining insurance authorization can be challenging, the former option is generally not financially feasible. The latter option is not only more cost-effective, but, highlights the need for early coordination of patient care to minimize unnecessary expenditures, which our current healthcare system is emphasizing. The challenges of co-registration between PET/CT and planning CT, however, should not be minimized. Given that the goal of using PET/CT is to accurately delineate the target when its location on the planning CT is ambiguous, being able to accurately co-register these two images is crucial. Another means of facilitating accurate co-registration which the authors do not comment on is the use of an immobilization device during PET/CT acquisition, which our group has implemented in past publications (2,7,8). This same device

can then be used for the planning CT to further improve co-registration accuracy.

Next, the authors describe the challenges faced in image interpretation and respiratory motion management and describe the methods by which treatment planning volumes should be generated based on the PET/CT. They highlight the need for a multi-disciplinary approach to target volume delineation including consultation with a nuclear medicine physician. This is a commendable recommendation because it allows the radiation oncologist to benefit from the expertise of the nuclear medicine physician skilled in interpreting PET/CT and to clarify any ambiguities on the PET/CT itself. The authors also make recommendations for target volume delineation based on whether respiratory compensation is utilized, highlighting the fact that PET/CTs are acquired during free breathing and therefore inherently account for tumor motion by averaging the tumor location during the breathing cycle. In order to take advantage of this aspect of the PET, however, the importance of accurately co-registering the PET/CT and planning CT is once again emphasized. To conclude this section of their manuscript, the authors comment on the difficulty with standardizing PET contouring secondary to variations in the window/level (W/L) settings and biological factors which may affect the levels of background PET activity. One way to standardize the use of PET in target volume delineation is by utilizing the “halo” to define the GTV. We have previously defined an anatomic biologic contour(ABC)-based GTV as the region encompassed by the distinct “halo” around areas of maximal SUV uptake and shown that its use in target volume delineation significantly decreases inter-observer variability in radiation treatment planning (2,7,8). This “halo” not only aids in increasing concordance among observers, but, can also increase the accuracy of target volume delineation as shown by Lin *et al.* (9).

Another challenge posed by the authors is the identification of which lymph nodes to include in the delineation of the target volume given that lymph nodes harboring disease require inclusion within the treatment field. Identification of pathologic lymph nodes is complicated by their oftentimes lower standard-uptake value (SUV) as compared to the primary tumor. Use of PET/CT for predicting nodal involvement is associated with a variable false positive rate (10), especially in the case of small sub-centimeter lymph nodes (11). Some authors have attempted to identify an SUV cut-off in order to improve the accuracy of the test, however, the choice of which cut-off value to use remains a subject of debate and

the false positive rate cannot be completely eliminated (12,13). Another means of increasing the accuracy of PET/CT in identifying pathologic lymph nodes which has been proposed is using the ratio of lymph node to primary tumor SUV (14). This is a novel parameter which still needs to be validated in other studies but has shown promising results.

The authors conclude by addressing the future of PET fusion technology with a discussion of the combination between functional MR and PET information. Another compelling imaging modality which may become useful for target volume delineation is hypoxia imaging (15-17). Hypoxia PET combines a variety of 2-nitroimidazoles, labeled with fluor-18 (^{18}F), as the radiotracer. These compounds are able to identify regions of tumor hypoxia. Use of hypoxia PET for contouring target volumes will allow us to go one step further by varying radiation dose levels within the target volume to target radio-resistant hypoxic regions. While more research is needed before this can be applied to a clinical setting, it holds promise as an imaging modality to aid in target volume delineation.

In conclusion, the use of PET/CT for target volume delineation in curative intent radiation therapy of NSCLC has allowed for significant improvement in contouring accuracy and decrease in inter-observer variability. However, there are several practical limitations to its use that need to be strongly considered. Konert *et al.* (1) provide a comprehensive analysis of the challenges presented by the use of PET/CT in target delineation in NSCLC. Their recommendations to improve the application of PET are essential given that it is currently the standard method for delineating treatment volumes. Their publication also highlights the need for improving methods of imaging acquisition and target volume determination to guide practice as we move forward.

Acknowledgements

None.

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

References

1. Konert T, Vogel W, MacManus MP, et al. PET/CT imaging for target volume delineation in curative intent

- radiotherapy of non-small cell lung cancer: IAEA consensus report 2014. *Radiother Oncol* 2015;116:27-34.
2. Ashamalla H, Raffla S, Parikh K, et al. The contribution of integrated PET/CT to the evolving definition of treatment volumes in radiation treatment planning in lung cancer. *Int J Radiat Oncol Biol Phys* 2005;63:1016-23.
 3. Bradley J, Thorstad WL, Mutic S, et al. Impact of FDG-PET on radiation therapy volume delineation in non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2004;59:78-86.
 4. Riegel AC, Berson AM, Destian S, et al. Variability of gross tumor volume delineation in head-and-neck cancer using CT and PET/CT fusion. *Int J Radiat Oncol Biol Phys* 2006;65:726-32.
 5. Brianzoni E, Rossi G, Ancidei S, et al. Radiotherapy planning: PET/CT scanner performances in the definition of gross tumour volume and clinical target volume. *Eur J Nucl Med Mol Imaging* 2005;32:1392-9.
 6. Bassi MC, Turri L, Sacchetti G, et al. FDG-PET/CT imaging for staging and target volume delineation in preoperative conformal radiotherapy of rectal cancer. *Int J Radiat Oncol Biol Phys* 2008;70:1423-6.
 7. Tejwani A, Lavaf A, Parikh K, et al. The role of PET/CT in decreasing inter-observer variability in treatment planning and evaluation of response for cervical cancer. *Am J Nucl Med Mol Imaging* 2012;2:307-13.
 8. Ashamalla H, Guirgius A, Bieniek E, et al. The impact of positron emission tomography/computed tomography in edge delineation of gross tumor volume for head and neck cancers. *Int J Radiat Oncol Biol Phys* 2007;68:388-95.
 9. Lin S, Han B, Yu L, et al. Comparison of PET-CT images with the histopathological picture of a resectable primary tumor for delineating GTV in nonsmall cell lung cancer. *Nucl Med Commun* 2011;32:479-85.
 10. Schmidt-Hansen M, Baldwin DR, Hasler E, et al. PET-CT for assessing mediastinal lymph node involvement in patients with suspected resectable non-small cell lung cancer. *Cochrane Database Syst Rev* 2014;11:CD009519.
 11. Billé A, Pelosi E, Skanjeti A, et al. Preoperative intrathoracic lymph node staging in patients with non-small-cell lung cancer: accuracy of integrated positron emission tomography and computed tomography. *Eur J Cardiothorac Surg* 2009;36:440-5.
 12. Hellwig D, Graeter TP, Ukena D, et al. ¹⁸F-FDG PET for mediastinal staging of lung cancer: which SUV threshold makes sense? *J Nucl Med* 2007;48:1761-6.
 13. Lee BE, Redwine J, Foster C, et al. Mediastinoscopy might not be necessary in patients with non-small cell lung cancer with mediastinal lymph nodes having a maximum standardized uptake value of less than 5.3. *J Thorac Cardiovasc Surg* 2008;135:615-9.
 14. Mattes MD, Moshchinsky AB, Ahsanuddin S, et al. The Ratio of Lymph Node to Primary Tumor SUV on PET/CT Accurately Predicts Nodal Malignancy in Non-Small Cell Lung Cancer. *Clin Lung Cancer* 2015. pii: S1525-7304(15)00163-1.
 15. Servagi-Vernat S, Differding S, Hanin FX, et al. A prospective clinical study of 18F-FAZA PET-CT hypoxia imaging in head and neck squamous cell carcinoma before and during radiation therapy. *Eur J Nucl Med Mol Imaging* 2014;41:1544-52.
 16. Hu M, Xing L, Mu D, et al. Hypoxia imaging with 18F-fluoroerythronitroimidazole integrated PET/CT and immunohistochemical studies in non-small cell lung cancer. *Clin Nucl Med* 2013;38:591-6.
 17. van Elmpt W, Zegers CM, Das M, et al. Imaging techniques for tumour delineation and heterogeneity quantification of lung cancer: overview of current possibilities. *J Thorac Dis* 2014;6:319-27.

Cite this article as: Tchelebi L, Ashamalla H. Overcoming the hurdles of using PET/CT for target volume delineation in curative intent radiotherapy of non-small cell lung cancer. *Ann Transl Med* 2015;3(14):191. doi:10.3978/j.issn.2305-5839.2015.07.02