



Identifying and mitigating factors contributing to 30-day hospital readmission in high risk patient populations

Michael P. Rogers, Paul C. Kuo

Department of Surgery, University of South Florida Morsani College of Medicine, Tampa, FL, USA

Correspondence to: Paul C. Kuo, MD, MS, MBA. Professor of Surgery Department of Surgery Morsani College of Medicine University of South Florida. Email: paulkuo@usf.edu.

Comment on: Garg SK, Goyal H, Obaitan I, *et al.* Incidence and predictors of 30-day hospital readmissions for liver cirrhosis: insights from the United States National Readmissions Database. *Ann Transl Med* 2021;9:1052.

Submitted Sep 14, 2021. Accepted for publication Sep 26, 2021.

doi: 10.21037/atm-2021-11

View this article at: <https://dx.doi.org/10.21037/atm-2021-11>

Unanticipated hospital readmission may significantly impact patient quality of life, hospital system resource utilization, and healthcare expenditure. The cost of unplanned hospital readmission is estimated between \$20–40 billion dollars annually in the United States (1,2). Particular conditions have been identified to contribute disproportionately to readmission and include congestive heart failure (CHF), septicemia, and pneumonia (2). Accordingly, significant attempts have been made to identify and mitigate factors contributing to 30-day hospital readmission. Efforts by the Centers for Medicare and Medicaid Services (CMS) Hospital Readmissions Reduction Program (HRRP), Joint Commission on Accreditation of Healthcare Organizations (JCAHO) performance metrics, and readmission rates as a measure of quality and hospital financial performance have endeavored to reduce unnecessary readmissions to the benefit of patients and hospitals alike (3). Robust national datasets have allowed researchers to identify predictors of hospital readmission unique to individual diagnoses and procedures (4–7). Consequently, readmission rates declined from 21.5% in 2007 to 17.8% in 2015 for conditions targeted by CMS, and from 15.3% to 13.1% for non-targeted conditions (8). As reimbursement and quality metrics are increasingly tied to patient outcome (i.e., mortality, infection, and unplanned 30-day readmission), interest in further identifying contributing factors remains a priority.

Various strategies have been explored to abate unforeseen hospital readmission. Models determining optimal discharge timing, ideal hospital lengths of stay, location of patient discharge, and the temporal influence of

index hospitalization have all been investigated to further reduce this burden (6,7,9–11). While the complete cadre of factors comprising the healthcare macroenvironment and their influence on readmission has yet to be elucidated, several authors have made significant strides in identifying modifiable predictors for individualized disease processes. Kothari and colleagues' utilization of the Healthcare Cost and Utilization Project State Inpatient Database (HCUP SID) for Florida and California to determine the influence of patient discharge location on readmission after liver transplantation represents one such approach (9). The HCUP-SID is developed and maintained by the Agency for Healthcare Research and Quality (AHRQ) to inform nation, State, and community level decision making by providing data elements of inpatient hospital stays (12). Elements including principal diagnoses and procedures, admission and discharge status, patient demographics, expected payment source, total charges, and length of stay allow for thorough investigation and consideration of the various factors associated with hospital readmission. Kothari *et al.*'s analysis revealed similar findings to prior work in patients with a primary diagnosis of congestive heart failure and myocardial infarction. Patients discharged to inpatient rehabilitation and long-term acute care facilities following liver transplantation had lower risk of 30-day readmission when compared to patients discharged to home (9). This and similar analyses may inform decision making when considering discharge planning for individualized patients that may be at risk from unplanned hospital readmission. The strengths of these large datasets in providing

longitudinal follow-up, cost data, and discharge locations offers novel understanding of the patient, surgeon, and hospital-level factors that may influence readmission.

Mitigating hospital readmission in patients with liver cirrhosis is a topic of significant interest. Previous efforts at identifying and understanding knowledge gaps in the care of these patients have explored patient and hospital factors (13-15). Orman and colleagues' recent systematic review found increased model for end-stage liver disease (MELD) score frequently associated with higher readmission (14). Similarly, Wei *et al.* single state analysis identified clinical complexity as well as sociodemographic factors as strong predictors of readmission (13). These studies highlight ongoing work in this cohort with high 30-day readmission. Despite the burden of these admissions however, to date, studies have been limited in the breadth of included factors for analysis. Orman *et al.* comment on the wide heterogeneity of studies included for systematic review, including differences in inclusion/exclusion criteria between studies, cirrhosis-specific and non-specific factors analyzed, and the ability to accurately capture hospital readmission (14). Further, a lack of study of the social determinants of health in this population and the differences in methodology across studies has hindered progress. As a result, a paucity of information currently exists to adequately address readmission in this cohort.

Garg and colleagues' novel investigation of factors contributing to 30-day readmission for liver cirrhosis similarly informs physicians using a large robust dataset and has several important findings (5). Readmission for cirrhosis is high with resultant significant financial impact on healthcare systems (16,17). Indeed, hospitalization costs secondary to cirrhosis and its sequela have been reported to be even greater than that of congestive heart failure or chronic obstructive pulmonary disease (COPD) (18). Garg *et al.* utilized the largest nationally representative inpatient sample, the Nationwide Readmission Database, which provides unique, all-payer information on approximately 49% of all hospitalizations occurring in 27 geographically diverse states in the US (19). Including over 300,000 patients, the authors identified a 31.4% readmission rate, substantially higher than the reported national average readmission rates for any other medical condition (20). Interestingly, though the CMS HRRP does not include cirrhosis currently, this remarkably high readmission rate makes a strong case for including this diagnosis in the future. Perhaps expectantly, the authors found patients with co-morbid conditions including CHF, COPD, peripheral

vascular disease, and diabetes mellitus, a substantial proportion of those readmitted.

Similar to previous analyses, Garg *et al.* observed post-discharge care had a significant impact on hospital readmission in this cohort (5,7,21). In their study, patients with chronic liver disease had significantly less post-discharge care (i.e., skilled nursing facilities, long term acute care hospitals, etc.) than patients admitted for CHF or COPD, despite higher readmission rates. Accordingly, the authors advocate for the development and implementation of strategies to ensure longitudinal support of these at-risk patients (5). Given this influence on readmission, discharge disposition should likely be dictated by individual patient and healthcare system factors, including the status of the patient at discharge, presence of co-morbidities, ability to access to care once discharged, and whether the patient may require ongoing care not available as an outpatient.

The identification of modifiable factors for decreasing the incidence of readmission at the index hospitalization is important in making strides to reduce this burden. Patients undergoing esophagogastroduodenoscopy (EGD) in this study had lower rates of hospital readmission. The authors hypothesize this is consequent to the ability to identify and intervene on high-risk varices, thereby decreasing rates of readmission for bleeding (5). The thorough work-up of patients at the initial hospitalization is thus paramount to identify and appropriately stratify those at risk for complications or need for ongoing medical care. Patients with an increased Charlson comorbidity index, hepatorenal syndrome, hepatic encephalopathy, ascites, esophageal varices, non-alcoholic and biliary cirrhosis, or history of bariatric surgery deserve particular attention (5).

Reducing unplanned 30-day hospital readmission remains a priority for patients, physicians, healthcare systems, and payors alike. It is estimated that approximately one-third of readmissions in this cohort may be preventable (22). These events significantly impact patient quality of life, hospital system resource utilization, and overall healthcare cost. Identifying potential patient and hospital factors that may reduce these readmissions is therefore of significant consequence. Garg and colleagues are to be congratulated on their excellent contribution to the literature and in providing insight on the contributors to readmission in this previously understudied cohort using a large nationally representative database.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Annals of Translational Medicine*. The article did not undergo external peer review.

Conflicts of Interest: Both authors have completed the ICMJE uniform disclosure form (available at <https://dx.doi.org/10.21037/atm-2021-11>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- Jencks SF, Williams MV, Coleman EA. Rehospitalizations among patients in the Medicare fee-for-service program. *N Engl J Med* 2009;360:1418-28.
- Hines AL, Barrett ML, Jiang HJ, et al. Conditions With the Largest Number of Adult Hospital Readmissions by Payer, 2011: Statistical Brief #172. Healthcare Cost and Utilization Project (HCUP) Statistical Briefs. Rockville, MD 2011.
- Upadhyay S, Stephenson AL, Smith DG. Readmission Rates and Their Impact on Hospital Financial Performance: A Study of Washington Hospitals. *Inquiry* 2019;56:46958019860386.
- Shah RM, Zhang Q, Chatterjee S, et al. Incidence, Cost, and Risk Factors for Readmission After Coronary Artery Bypass Grafting. *Ann Thorac Surg* 2019;107:1782-9.
- Garg SK, Goyal H, Obaitan I, et al. Incidence and predictors of 30-day hospital readmissions for liver cirrhosis: insights from the United States National Readmissions Database. *Ann Transl Med* 2021;9:1052.
- Kothari AN, Qu LT, Gil LA, et al. Weekend readmissions associated with mortality following pancreatic resection for cancer. *Surg Oncol* 2020;34:218-22.
- Kothari AN, Loy VM, Brownlee SA, et al. Adverse Effect of Post-Discharge Care Fragmentation on Outcomes after Readmissions after Liver Transplantation. *J Am Coll Surg* 2017;225:62-7.
- Zuckerman RB, Sheingold SH, Orav EJ, et al. Readmissions, Observation, and the Hospital Readmissions Reduction Program. *N Engl J Med* 2016;374:1543-51.
- Kothari AN, Yau RM, Blackwell RH, et al. Inpatient Rehabilitation after Liver Transplantation Decreases Risk and Severity of 30-Day Readmissions. *J Am Coll Surg* 2016;223:164-171.e2.
- Cousin-Peterson E, Janjua HM, Barry TM, et al. Discharge timing: Does targeting an ideal length of stay for patients undergoing colectomy impact readmissions and costs of care? *Am J Surg* 2021;221:570-4.
- Andriotti T, Goralnick E, Jarman M, et al. The Optimal Length of Stay Associated With the Lowest Readmission Risk Following Surgery. *J Surg Res* 2019;239:292-9.
- Agency for Healthcare Research and Quality (9/13/21). Available online: www.hcup-us.ahrq.gov/sidoverview.jsp
- Wei M, Ford J, Li Q, et al. Hospital Cirrhosis Volume and Readmission in Patients with Cirrhosis in California. *Dig Dis Sci* 2018;63:2267-74.
- Orman ES, Ghabril M, Emmett TW, et al. Hospital Readmissions in Patients with Cirrhosis: A Systematic Review. *J Hosp Med* 2018. [Epub ahead of print]. doi: 10.12788/jhm.2967.
- Rosenblatt R, Cohen-Mekelburg S, Shen N, et al. Cirrhosis as a Comorbidity in Conditions Subject to the Hospital Readmissions Reduction Program. *Am J Gastroenterol* 2019;114:1488-95.
- Fagan KJ, Zhao EY, Horsfall LU, et al. Burden of decompensated cirrhosis and ascites on hospital services in a tertiary care facility: time for change? *Intern Med J* 2014;44:865-72.
- Seraj SM, Campbell EJ, Argyropoulos SK, et al. Hospital readmissions in decompensated cirrhotics: Factors pointing toward a prevention strategy. *World J Gastroenterol* 2017;23:6868-76.
- Di Pascoli M, Ceranto E, De Nardi P, et al. Hospitalizations Due to Cirrhosis: Clinical Aspects in a Large Cohort of Italian Patients and Cost Analysis Report. *Dig Dis* 2017;35:433-8.
- Nationwide Readmissions Database (NRD) (9/13/21). Healthcare Cost and Utilization Project (HCUP)). Available online: <https://www.hcup-us.ahrq.gov/>

- nrdoverview.jsp
20. Berry JG, Gay JC, Joynt Maddox K, et al. Age trends in 30 day hospital readmissions: US national retrospective analysis. *BMJ* 2018;360:k497.
 21. Pedersen PU, Ersgard KB, Soerensen TB, et al. Effectiveness of structured planned post discharge support

- to patients with chronic obstructive pulmonary disease for reducing readmission rates: a systematic review. *JBHI Database System Rev Implement Rep* 2017;15:2060-86.
22. Saberifiroozi M. Improving Quality of Care in Patients with Liver Cirrhosis. *Middle East J Dig Dis* 2017;9:189-200.

Cite this article as: Rogers MP, Kuo PC. Identifying and mitigating factors contributing to 30-day hospital readmission in high risk patient populations. *Ann Transl Med* 2021;9(21):1610. doi: 10.21037/atm-2021-11