## The REMARK checklist

Item to be reported		Reported on Page Number/Line Number	Reported on Section/Paragraph
INTR	DDUCTION		
1	State the marker examined, the study objectives, and any pre-specified hypotheses.	Page4/Line104-112	Introduction/Paragraph4
MATI	ERIALS AND METHODS	1	
Patie	nts		
2	Describe the characteristics (e.g., disease stage or co-morbidities) of the study patients, including their source and inclusion and exclusion criteria.	N/A	N/A
3	Describe treatments received and how chosen (e.g., randomized or rule-based).	N/A	N/A
Spec	men characteristics		
4	Describe type of biological material used (including control samples) and methods of preservation and storage.	Page4/Line120	Methods/Paragraph1
Assay	r methods		
5	Specify the assay method used and provide (or reference) a detailed protocol, including specific reagents or kits used, quality control procedures, reproducibility assessments, quantitation methods, and scoring and reporting protocols. Specify whether and how assays were performed blinded to the study endpoint.	Page4 -5/Line116-161	Methods/Paragraph1-5
Study	design		1
6	State the method of case selection, including whether prospective or retrospective and whether stratification or matching (e.g., by stage of disease or age) was used. Specify the time period from which cases were taken, the end of the follow-up period, and the median follow-up time.	N/A	N/A
7	Precisely define all clinical endpoints examined.	N/A	N/A
8	List all candidate variables initially examined or considered for inclusion in models.	N/A	N/A
9	Give rationale for sample size; if the study was designed to detect a specified effect size, give the target power and effect size.	N/A	N/A
Statis	tical analysis methods		1
10	Specify all statistical methods, including details of any variable selection procedures and other model-building issues, how model assumptions were verified, and how missing data were handled.	Page5-6/Line164-170	Methods/Paragraph6
11	Clarify how marker values were handled in the analyses; if relevant, describe methods used for cutpoint determination.	Page4/Line118-120	Methods/Paragraph1
			1

RESULTS Data				
Report distributions of basic demographic characteristics (at least age and sex), standard (disease-specific) prognostic variables, and tumor marker, including numbers of missing values.	N/A	N/A		
sis and presentation				
Show the relation of the marker to standard prognostic variables.	Page 6-8/Line194-252	Results/Paragraph3-7		
Present univariable analyses showing the relation between the marker and outcome, with the estimated effect (e.g., hazard ratio and survival probability). Preferably provide similar analyses for all other variables being analyzed. For the effect of a tumor marker on a time-to-event outcome, a Kaplan-Meier plot is recommended.	Page 6-8/Line194-252	Results/Paragraph3-7		
For key multivariable analyses, report estimated effects (e.g., hazard ratio) with confidence intervals for the marker and, at least for the final model, all other variables in the model.	N/A	N/A		
Among reported results, provide estimated effects with confidence intervals from an analysis in which the marker and standard prognostic variables are included, regardless of their statistical significance.	Page 7-8/Line226-252	Results/Paragraph6-7		
If done, report results of further investigations, such as checking assumptions, sensitivity analyses, and internal validation.	N/A	N/A		
USSION	1	1		
Interpret the results in the context of the pre-specified hypotheses and other relevant studies; include a discussion of limitations of the study.	Page 10-12/Line319-389	Discussion/Paragraph2-4		
Discuss implications for future research and clinical value.	Page 12/Line390-394	Discussion/Paragraph5		
	Describe the flow of patients through the study, including the number of patients included in each stage of the analysis (a diagram may be helpful) and reasons for dropout. Specifically, both overall and for each subgroup extensively examined report the numbers of patients and the number of events.  Report distributions of basic demographic characteristics (at least age and sex), standard (disease-specific) prognostic variables, and tumor marker, including numbers of missing values.  Sis and presentation  Show the relation of the marker to standard prognostic variables.  Present univariable analyses showing the relation between the marker and outcome, with the estimated effect (e.g., hazard ratio and survival probability). Preferably provide similar analyses for all other variables being analyzed. For the effect of a tumor marker on a time-to-event outcome, a Kaplan-Meier plot is recommended.  For key multivariable analyses, report estimated effects (e.g., hazard ratio) with confidence intervals for the marker and, at least for the final model, all other variables in the model.  Among reported results, provide estimated effects with confidence intervals from an analysis in which the marker and standard prognostic variables are included, regardless of their statistical significance.  If done, report results of further investigations, such as checking assumptions, sensitivity analyses, and internal validation.  USSION  Interpret the results in the context of the pre-specified hypotheses and other relevant studies; include a discussion of limitations of the study.	Describe the flow of patients through the study, including the number of patients included in each stage of the analysis (a diagram may be helpful) and reasons for dropout. Specifically, both overall and for each subgroup extensively examined report the numbers of patients and the number of events.  Report distributions of basic demographic characteristics (at least age and sex), standard (disease-specific) prognostic variables, and tumor marker, including numbers of missing values.  Sis and presentation  Show the relation of the marker to standard prognostic variables.  Page 6-8/Line194-252  Present univariable analyses showing the relation between the marker and outcome, with the estimated effect (e.g., hazard ratio and survival probability). Preferably provide similar analyses for all other variables being analyzed. For the effect of a tumor marker on a time-to-event outcome, a Kaplan-Meier plot is recommended.  For key multivariable analyses, report estimated effects (e.g., hazard ratio) with confidence intervals for the marker and, at least for the final model, all other variables in the model.  Among reported results, provide estimated effects with confidence intervals from an analysis in which the marker and standard prognostic variables are included, regardless of their statistical significance.  If done, report results of further investigations, such as checking assumptions, sensitivity analyses, and internal validation.  Page 10-12/Line319-389  Page 10-12/Line319-389		

From: McShane LM, Altman DG, Sauerbrei W, Taube SE, Gion M, Clark GM: Reporting recommendations for tumor marker prognostic studies (REMARK). J Natl Cancer Inst 2005; 97: 1180-1184.

Article information: https://dx.doi.org/10.21037/atm-21-6611
\*As the checklist was provided upon initial submission, the page number/line number reported may be changed due to copyediting and may not be referable in the published version. In this case, the section/paragraph may be used as an alternative reference.