

Construction of the prognostic model in non-metastatic renal cancer patients with venous tumor thrombus

Baohua Zhu[#], Yiren Yang[#], Zhao Huangfu[#], Wei Zhang, Aimin Jiang, Linhui Wang

Department of Urology, Changhai Hospital, Second Military Medical University, Shanghai, China

Contributions: (I) Conception and design: B Zhu, L Wang; (II) Administrative support: L Wang; (III) Provision of study materials or patients: B Zhu, Y Yang; (IV) Collection and assembly of data: B Zhu, Y Yang, Z Huangfu; (V) Data analysis and interpretation: B Zhu, Z Huangfu, Y Yang, A Jiang, W Zhang; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Linhui Wang, MD, PhD; Aimin Jiang, MD, PhD. Department of Urology, Changhai Hospital, Second Military Medical University, No. 168 Changhai Rd., Shanghai 200433, China. Email: wanglinhui@smmu.edu.cn; czjiangaimin@smmu.edu.cn.

Background: Venous system invasion is a prominent characteristic of local progression in renal cancer and treatment-naïve renal cancer patients with venous tumor thrombus (VTT) gained short natural course and poor prognosis. This study aimed to investigate the efficacy of the surgery and prognostic factors in non-metastatic renal cancer patients with VTT and to construct a nomogram prognostic model.

Methods: Clinical data of 114 non-metastatic renal cancer patients with VTT who underwent surgical treatment from January 2011 to September 2022 were retrospectively analyzed. In order to find independent risk factors of prognosis, survival analysis was performed via univariate and multivariate Cox regression models and Kaplan-Meier method. Nomogram prognostic model was established to calculate patients' risk scores. Receiver operating characteristic curve and decision curve analysis were conducted to evaluate the efficacy of the prognostic model.

Results: A total of 114 patients were included in this study and there were 48, 12, 25, 23, and 6 cases of grade 0–IV VTT. No perioperative death occurred. The 3-year probabilities of overall survival (OS) and 5-year probabilities of OS were 67% and 43.8%, respectively. Multivariate Cox regression analysis revealed that kidney tumor diameter, preoperative lactate dehydrogenase (LDH), and preoperative neutrophils were independent risk factors. Nomogram was constructed to predict prognosis in renal cancer patients with VTT based on above indicators and Mayo VTT grading. The area under the ROC curve of 1-, 2-, 3-, and 5-year OS of the patients were 0.82, 0.67, 0.57, and 0.55 respectively.

Conclusions: Surgical treatment enables renal cancer patients with VTT to gain a better prognosis. Kidney tumor diameter, preoperative LDH, and preoperative neutrophils were independent risk factors. The nomogram perfects the Mayo grading, and provides a reliable reference for evaluation of prognosis of renal cancer patients with VTT.

Keywords: Venous tumor thrombus (VTT); renal cancer; prognosis; nomogram model; surgical treatment

Submitted Jun 13, 2023. Accepted for publication Oct 20, 2023. Published online Nov 14, 2023. doi: 10.21037/tau-23-341

View this article at: https://dx.doi.org/10.21037/tau-23-341

Introduction

Renal cell carcinoma (RCC) is a malignant tumor originating from the kidney parenchyma, and its incidence accounts for 2–3% of adult malignancies (1), increasing by

1–2% annually (2). Venous system invasion is a prominent characteristic of local progression in renal cancer, affecting about 4% to 10% of all patients with RCC (3). Treatmentnaïve renal cancer patients with venous tumor thrombus (VTT) gained short natural course and poor prognosis,

^{*}These authors contributed equally to this work.

with median survival time of 5 months and 1-year cancerspecific survival (CSS) of 29% (4). A comprehensive review of a substantial patient series, which underwent systemic targeted therapy for in situ RCC tumor thrombi, indicated limited clinical efficacy in reducing the tumor thrombus level. Additionally, the study failed to uncover a noteworthy influence on the surgical approach among those who eventually required thrombectomy (5,6). Radical nephrectomy (RN) and VTT extraction have become the preferred treatment for non-metastatic renal carcinoma patients with VTT, of which the 5-year disease-free survival (DFS) is 39-60% (7-9). The surgery performed on these patients is characterized by its complexity, high-risk nature, and the potential for pulmonary arterial embolism due to detachment of cancer emboli. Such embolism poses a significant threat to patient survival. Furthermore, there exists a direct correlation between the grade of cancer emboli and the perioperative mortality rate, with highergrade emboli associated with an incremental increase in mortality risk (10-12). The great difficulty and risk of surgery require researchers to predict the prognosis of patients according to the different clinical and pathological characteristics of each patient, so as to better formulate treatment strategies. The objectives of this study were to investigate the efficacy of the surgery in non-metastatic renal cancer patients with VTT and to analyze the clinicopathologic elements for the establishment of the prognostic model, and thus providing a theoretical basis for the development of individualized therapy. We present

Highlight box

Key findings

 We first found that preoperative lactate dehydrogenase (LDH) and preoperative neutrophils might influence the prognosis of renal cell carcinoma patients with venous tumor thrombus (VTT).

What is known and what is new?

- The Mayo grading has long been regarded as the standard for assessing the prognosis of patients with renal cancer accompanied by VTT.
- We constructed a new nomogram including tumor diameter, preoperative LDH, and preoperative neutrophils to perfect the Mayo grading.

What is the implication, and what should change now?

 Now Mayo grading with our nomogram provides a more accurate assessment of the prognosis for patients with renal cancer accompanied. this article in accordance with the TRIPOD reporting checklist (available at https://tau.amegroups.com/article/view/10.21037/tau-23-341/rc).

Methods

Study cobort

Clinicopathologic and prognostic data of 114 non-metastatic renal cancer patients with VTT who underwent surgical treatment in Shanghai Changhai Hospital from January 2011 to September 2022 were retrospectively analyzed. The study protocol was designed in accordance with the Declaration of Helsinki (as revised in 2013) and approved by the Ethics Committee of Changhai Hospital (No. CHEC2021-191). Informed consent was waived by Ethics Committees due to the retrospective nature of the study.

Inclusion criteria: (I) non-metastatic renal carcinoma patients with VTT grading 0–IV; (II) patients who underwent RN and VTT extraction without VTT detachment and perioperative death; and (III) postoperative pathological confirmed RCC.

Exclusion criteria: (I) patients displaying evidence of metastasis based on preoperative imaging; (II) patients who underwent exploratory laparotomy but did not undergo complete surgical intervention; and (III) patients who were lost to follow-up during the study period.

Postoperative follow-up

The follow-up data of all the patients were obtained through telephone calls or outpatient visits. Last follow-up time was on 10th February, 2023. The primary endpoint was overall survival (OS), which was defined as the time from surgery to death or the last follow-up.

Statistical analyses

Continuous variables were presented as the mean \pm standard deviation, and categorical variables were summarized by frequencies and percentages. The chi-squared test or Fisher's exact test was applied to compare categorical variables, and the t-test or nonparametric Wilcoxon's test was used to compare continuous variables between groups. Among all patients enrolled in this work, two-sided P value <0.05 indicated statistical significance. Univariate and multivariate Cox regression analyses were used to determine the effect of

selected variables on OS. Nomogram model was constructed with four clinical parameters based on multivariate Cox regression analysis. A novel risk score system was developed in this study, incorporating four clinical parameters. The integration of these parameters resulted in the creation of a new risk system, which was represented using a nomogram. The construction of the nomogram was facilitated by the utilization of the R package rms. Calibration curve was also depicted to compare the real and predicted incidence. Receiver operating characteristic (ROC) curve of 1-, 2-, 3-, and 5-year OS of all patients was utilized to access the sensitivity and specificity of risk system, and area under the ROC curve (AUC) larger than 0.6 was considered as suitable, while lager than 0.75 was considered as well performed. AUC was visualized by using R package 'pROC' packages. To compare the clinical benefit of applying novel risk score system in clinical practice when comparing with other parameters, including lactate dehydrogenase (LDH), neutrophil, thrombus grading and tumor diameter, we applied DCA curve to assess prognostic benefit of risk score system, which was illustrated by R package ggDCA. Decision curve analysis (DCA) is a statistical method used to assess and contrast the clinical utility of different prediction models or diagnostic tests. It provides a framework to assess the net benefit of using a particular model or test in clinical decision-making. In DCA, an assortment of threshold probabilities is defined, representing the clinician's willingness to treat or intervene based on the predicted probability of an outcome. The calculation of the net benefit involves an assessment of advantages and disadvantages associated with each threshold probability. By plotting the net benefit against the threshold probability, the performance of different models or tests can be compared. A model or test with higher net benefit over a wide range of threshold probabilities indicates superior clinical utility. Furthermore, calibration curve, known as a calibration plot or reliability diagram, is a graphical representation that evaluates the performance and calibration of a predictive model, was adopted to evaluate how well the predicted probabilities from our risk score system with the actual observed outcomes and determine whether our model is overconfident or underconfident in its predictions.

Results

Baseline characteristics of included patients

Of all the 114 patients, 85 were male and 29 were female,

aging 58.5±12.3 years, body mass index (BMI) 23.5± 3.2 kg/m². The tumors were located on the left side in 56 cases and right side in 58 cases. The average diameter of the tumors was 8.52±3.79 cm. Ninety cases are clear cell RCC (ccRCC) while 24 were non-ccRCC [8 papillary RCC (pRCC), 3 collecting duct kidney cancer, 3 renal cancer associated with TFE3 gene fusions, 1 renal oncocytoma, 1 FH deficient RCC, 2 sarcomatoid RCC, 1 renal cell carcinoma unclassified, and 5 other types]. Grading 0-IV (the Mayo classification) VTT involved 48, 12, 25, 23, and 6 cases respectively. We can observe that compared to the group alive, patients in the group dead had higher values in proportion of open surgery, length of stay, preoperative neutrophil, and tumor diameter. More baseline data are provided in Table 1. More detailed clinical data of the patients are shown in Table S1.

Survival analysis and risk factors for prognosis

All the 114 patients were followed up and by the time of the last follow-up, 64 patients had died. Three- and five-year OS were 67% and 43.8% respectively.

Univariate Cox regression analysis was performed using gender, age, BMI, the presence of hypertension, the presence of diabetes, preoperative hemoglobin, preoperative creatinine, preoperative LDH, preoperative neutrophils, tumor side, pathological type, the presence of lymphatic metastasis, the presence of perirenal adipose tissue invasion, tumor thrombus grading, tumor diameter, Clavien grading, postoperative targeted therapy, five of which were included into multivariate Cox regression analysis (P<0.05). Multivariate Cox regression analysis indicated that tumor diameter, preoperative LDH and preoperative neutrophils were independent risk factors for survival (*Table 2*).

Construction and validation of nomogram model

Considering the generality of VTT Mayo grading, the new nomogram model for non-metastasis renal cancer patients with VTT was established using tumor diameter, preoperative LDH, preoperative neutrophils and Mayo grading (*Figure 1*). The prognosis of high-risk group (*Figure 2A*, risk score >136.34) is poorer than that of lowrisk group (*Figure 2B*).

Under the nomogram model, the AUC of 1-, 2-, 3-, and 5-year OS of patients was 0.82, 0.67, 0.57, and 0.55, respectively (*Figure 3*). In the bootstrapped DCA analysis conducted on patients with thrombus, the risk score

Table 1 Baseline data of 114 included patients

Characteristics	Group alive (n=54)	Group dead (n=60)	P value
Gender			0.108
Female	10 (8.8)	19 (16.7)	
Male	44 (38.6)	41 (36.0)	
Hypertension			0.164
No	33 (28.9)	44 (38.6)	
Yes	21 (18.4)	16 (14.0)	
Diabetes			0.620
No	46 (40.4)	53 (46.5)	
Yes	8 (7.0)	7 (6.1)	
Tumor side			0.859
Left	27 (23.7)	29 (25.4)	
Right	27 (23.7)	31 (27.2)	
Surgical operative methods			0.009
Laparoscopic surgery	23 (20.2)	12 (10.5)	
Open surgery	31 (27.2)	48 (42.1)	
Tumor thrombus grading			0.941
Grade 0	24 (21.1)	24 (21.1)	
Grade I	5 (4.4)	7 (6.1)	
Grade II	12 (10.5)	13 (11.4)	
Grade III	11 (9.6)	12 (10.5)	
Grade IV	2 (1.8)	4 (3.5)	
Clavien-Dindo grading			0.593
Grade I	27 (23.7)	27 (23.7)	
Grade II/III	27 (23.7)	33 (28.9)	
ASA scoring			0.661
1	2 (1.8)	3 (2.6)	
2	44 (38.6)	44 (38.6)	
3	8 (7.0)	11 (9.6)	
4	0 (0.0)	2 (1.8)	
Lymphatic metastasis			0.394
Metastasis	6 (5.3)	10 (8.8)	
Non-metastasis	48 (42.1)	50 (43.9)	
Perirenal adipose tissue invasion			0.359
Invasion	18 (15.8)	25 (21.9)	
Non-invasion	36 (31.6)	35 (30.7)	
Pathological type			0.529
ccRCC	44 (38.6)	46 (40.4)	
Non-ccRCC	10 (8.8)	14 (12.3)	

Table 1 (continued)

Table 1 (continued)

Characteristics	Group alive (n=54)	Group dead (n=60)	P value		
Age (years)	58 [53, 65]	60 [51.75, 69]	0.388		
BMI (kg/m²)	24.21±3.09	22.9±3.24	0.030		
Operation time (hours)	3 [2.3, 3.77]	3 [2.45, 4.05]	0.408		
Length of stay (days)	10 [7, 13.75]	14.5 [10, 17]	<0.001		
Intraoperative blood loss (mL)	850 [200, 1,575]	1,200 [500, 2,025]	0.208		
Preoperative LDH (U/L)	173 [152.25, 191.5]	169.5 [148, 248.75]	0.658		
Preoperative creatinine (µmol/L)	84.5 [75, 98.5]	86.5 [69.75, 101.5]	0.827		
Preoperative neutrophils (×10 ⁹ /L)	3.7 [3, 4.97]	4.95 [3.58, 5.73]	0.019		
Preoperative hemoglobin (g/L)	126 [107, 137]	114 [96.75, 130.25]	0.176		
Tumor diameter (mm)	70 [50, 90]	80 [60, 110]	0.033		
Postoperative targeted therapy			0.353		
No	25 (21.9)	33 (28.9)			
Yes	29 (25.4)	27 (23.7)			

Data are presented as n (%), median [IQR], or mean \pm SD. ASA, American Society of Anesthesiologists; ccRCC, clear cell renal cell carcinoma; BMI, body mass index; LDH, lactate dehydrogenase; IQR, interquartile range; SD, standard deviation.

Table 2 Univariate and multivariate Cox regression analysis

Mariakian	Univariate Cox regre	ssion analysis	Multivariate Cox regression analysis				
Variation -	HR (95% CI)	P value	HR (95% CI)	P value			
Gender (male vs. female)	0.64 (0.37–1.11)	0.109					
Age	1.01 (0.99–1.04)	0.244					
BMI	0.93 (0.85–1.01)	0.077	0.94 (0.87–1.04)	0.117			
Tumor side (left vs. right)	0.99 (0.60–1.65)	0.978					
Hypertension (yes vs. no)	0.97 (0.55–1.73)	0.928					
Diabetes (yes vs. no)	0.88 (0.39–1.95)	0.746					
Tumor thrombus grading	1.21 (1.01–1.46)	0.039	1.12 (0.90–1.40)	0.310			
Pathological type (non-ccRCC vs. ccRCC)	1.50 (0.82–1.76)	0.189					
Preoperative hemoglobin	0.98 (0.97–1.00)	0.004					
Preoperative creatinine	1.00 (0.99–1.01)	0.715					
Preoperative neutrophil	1.28 (1.06–1.54)	0.010	1.29 (1.07–1.55)	0.007			
Preoperative LDH	1.00 (1.00–1.01)	<0.001	1.00 (1.00–1.01)	0.008			
Lymphatic metastasis	1.78 (0.90–3.54)	0.098					
Perirenal adipose tissue invasion	1.70 (0.99–2.92)	0.052					
Clavien-Dindo grading (grade I vs. grade II/III)	1.29 (0.77–2.14)	0.335					
Tumor diameter	1.02 (1.01–1.02)	<0.001	1.01 (1.00–1.03)	0.011			
Postoperative targeted therapy	0.97 (0.58–1.62)	0.894					

HR, hazard ratio; CI, confidence interval; BMI, body mass index; ccRCC, clear cell renal cell carcinoma; LDH, lactate dehydrogenase.




Figure 1 Establishment of a nomogram predicting 1-, 3-, and 5-year OS based on the independent prognostic factors in patients with tumor thrombus grading. Firstly, each individual subject's value was located on the corresponding axis of the nomogram. Secondly, a vertical line was drawn upward from this point to intersect with the "Points" axis, thereby generating a specific point. This process was repeated for all variables of interest. In the next step, the points obtained for each variable were summed up, and the total sum was located on the "Total points" line. Finally, by projecting this point onto the 1-, 3-, and 5-year OS lines, the corresponding probabilities of survival at each time point were determined. LDH, lactate dehydrogenase; OS, overall survival.

exhibited a significantly higher net benefit in accurately predicting OS, particularly in the 5-year period. This was in comparison to preoperative LDH, neutrophil count, tumor thrombus grading, and tumor diameter (*Figure 4A*). Moreover, when examining the bootstrapped calibration plots of the risk score, it can be observed that there were no adverse deviations between the predicted risk and observed risk for 1-, 2-, 3-, and 5-year OS across the entire range (*Figure 4B*).

Discussion

RCC with VTT is always a conundrum in urology. Treatment-naïve renal cancer patients with VTT gained short natural course with median survival time of 5 months and 1-year CSS of 29% (4). In recent years, with the advancements in surgical techniques and laparoscopic devices, the success rate of RN and VTT extraction has been greatly enhanced, and 5-year OS of patients after surgery was 39–60%, significantly better than those with

no surgery, which shows that surgery is the most efficient method (13-15). This study showed that 3- and 5-year OS was 67% and 43.8% respectively, and in fact, similar studies were also reported before. In the realm of prognostic models for patients with RCC and VTT, a unified standard is currently lacking, as reported in the literature. Additionally, certain models within this domain incorporate non-conventional diagnostic variables as predictive factors, significantly diminishing their clinical applicability. Meanwhile, survival models for RCC patients with VTT, based on patient data and exhibiting practicality, do possess limitations such as suboptimal calibration and small sample sizes within specific subgroups of predictive variables (16-19). Therefore, this study sought to analyze the data of 114 patients with non-metastatic RCC accompanied by VTT who were treated at Changhai Hospital. The primary objective was to determine the independent prognostic factors for postoperative OS in this specific patient cohort and develop a more robust and effective personalized survival model for them.




Figure 2 Prognostic value of risk score based on nomogram. (A) The optimal cutoff value of the risk score from nomogram. (B) KM curves between low and high-risk subgroups based on optimal cutoff value. KM, Kaplan-Meier.

Tumor diameter influences surgical difficulties and patients' OS, which are the focus of clinicians all the time. Previous studies demonstrated that the malignancy and invasion of renal cancer increased with the tumor size (20). Yang *et al.* proved that smaller ccRCC has less malignant and better prognosis than bigger one (21). Similar clinical research mentioned that large tumor diameter and renal

vein invasion predicted worse DSS and progression-free survival (PFS) (22). Coincidentally, this study indicated that with the tumor diameter increasing, RCC patients with VTT had worse prognosis. In fact, larger primary tumor closely adhere to surrounding tissues so more tissues are stripped to expose the kidney, leading to longer operation time, more blood loss and higher probability




Figure 3 ROC curves of the nomogram at 1-, 2-, 3-, and 5-year OS in our cohort. AUC, area under the ROC curve; ROC, receiver operating characteristic; OS, overall survival.

of postoperative complications (17), which contributes to preoperative risk analysis and long-term prognosis evaluation. Abel et al. reported the prognostic data of 636 non-metastasis renal cancer with VTT, finding the correlation between tumor diameter and recurrence rate, and constructed a nomogram to predict tumor postoperative recurrence rate including six factors (23). The discovery that tumor diameter has an impact on patient prognosis highlights several important considerations: Firstly, preoperative imaging examinations play a critical role in accurately delineating tumor borders. Secondly, meticulous surgical techniques during the procedure are essential for discerning tumor boundaries. Lastly, postoperative followup should be given special attention for patients with larger tumor diameters, and when necessary, preventive medication should be considered.

The impact of VTT grading on prognosis still remains controversial. In our study, univariate Cox regression




Figure 4 Performance of risk score in our cohort. (A) Comparison of the DCA of the nomogram and other parameters including preoperative LDH level, preoperative neutrophil count, tumor thrombus grading and tumor diameter for 1-, 2-, 3-, and 5-year OS in our cohort. (B) Calibration plots of the nomogram for 1-, 2-, 3-, and 5-year OS in our cohort. OS, overall survival; LDH, lactate dehydrogenase; DCA, decision curve analysis.

analysis suggested that VTT grading was related to the prognosis of patients, but multivariate Cox regression analysis indicated no correlation. This phenomenon may be because the sample numbers were relatively small and grade 0 VTT accounts for a relatively large proportion. This observation could potentially be attributed to several factors. Firstly, the relatively small sample size and the significant proportion of grade 0 VTT cases might have influenced the results. Additionally, advancements in surgical techniques, mean duration of follow-up, and the specific clinicopathologic factors investigated alongside tumor thrombus levels could have contributed to these discrepancies. It is worth noting that significant factors predicting OS may vary depending on the duration of follow-up, which can lead to conflicting conclusions regarding the impact of tumor thrombus extension on survival outcomes. A research from Shengjing Hospital of China Medical University revealed no differences in CSS between renal vein group and inferior vena cava (IVC) group (24). Another study subdivided 142 patients with VTT into high-Mayo group and low-Mayo group, indicating that VTT grading did not affect long-term survival (9). However, high Mayo grading was associated with postoperative 12-month death risk. By contrast, several studies showed that the prognosis of patients with renal VTT was better than patient with IVC tumor thrombus (25,26). Martínez-Salamanca et al. analyzed 1122 patients with different grade VTT, indicating that 5-year OS of VTT in renal vein, above the renal vein but below the diaphragm, above the diaphragm was respectively 43.2%, 37%, 22% (27). Unexpectedly, they reported that VTT grading was the risk factor for prognosis of patients with VTT. We are aware that Mayo grading is subdivided by tumor thrombus position in vein, and different grade tumor thrombus means different surgical strategies and difficulties. So, the correlation between VTT grading and prognosis still deserves further investigation.

Moreover, researchers raised some prognostic factors based on laboratory measures including preoperative hemoglobin, preoperative serum alkaline phosphatase (ALP), preoperative alanine transaminase (ALT), preoperative albumin, and so on (26,28). This study also indicated that preoperative LDH and preoperative neutrophils were independent risk factors for OS. LDH, one of the most important enzymes of glycolysis, is a glycolytic enzyme that catalyzes the conversion of pyruvate to lactic acid. In tumor cells, glycolysis is extremely elevated regardless of oxygen availability, which is known as the

Warburg effect. Low LDH expression inhabits tumor invasion and metastasis by reducing the proliferative capacity and resistance of chemotherapy (29). Recent years, a multitude of studies demonstrated that serum LDH level was associated with prognosis (30,31). Our research found that high-level preoperative LDH was related to poor prognosis. The level of serum LDH is a convenient and economic prognostic indicator. However, the reason of serum LDH promotion in renal cancer patients with VTT is still not clear. Large multi-center studies are needed to confirm these findings. These years, some systemic inflammatory and immune response indicators such as C-reactive protein and neutrophil to lymphocyte ratio (NLR) were considered to predict prognosis (32). Our nomogram included preoperative neutrophils, which was one of independent risk factor of OS. Thus, with the preoperative neutrophils increasing, the prognosis of patients are poorer. This situation may be in contacted in the change of tumor immune microenvironment (TIME). Ghatalia et al. (33) reported that neutrophil augmented renal cancer postoperative recurrence by analyzing tumorinfiltrating immune cell and located ccRCC recurrence rate. The above simple preoperative blood biochemical indicators can be easily analyzed and may help select high-risk patients. Consistent with previous research, our study has yielded further evidence supporting the use of preoperative LDH and preoperative neutrophil as independent prognostic indicators in non-metastatic renal cancer patients with VTT. Moreover, incorporating these parameters into prognostic nomograms has demonstrated improved predictive accuracy and clinical utility. These findings highlight the potential of preoperative LDH and preoperative neutrophils as valuable tools for risk stratification and treatment decision-making in this patient population.

Lymph node dissection plays a pivotal role in the surgical management of malignant tumors in the genitourinary system. However, the value of lymph node dissection in RN has been a subject of ongoing debate. At our institution, lymph node dissection is performed selectively, based on evidence of lymph node enlargement on preoperative imaging studies. Only when tumor cells are identified in the resected lymph node specimens during surgery are patients classified as N1. According to our established criteria, our study findings did not demonstrate a significant association between lymph node metastasis and patient survival. While some studies have reported potential oncological benefits of lymph node dissection (34,35), others argue that it does not

confer any survival advantages for kidney cancer patients (36,37). Clinical practice involving regional lymph node dissection is often driven by the desire to determine the local extent of the tumor and potentially guide decisions regarding adjuvant therapy. However, it remains uncertain whether lymph node dissection itself provides inherent survival benefits. Karmali et al. (38) analyzed sentinel lymph nodes in renal cancer patients, and found that RCC is more inclined to hematogenous dissemination rather than lymphatic spread. Patients with regional lymph node metastasis typically already exhibit distant metastasis, with only a small percentage (2-5%) presenting solely with lymph node involvement and no distant spread. Consequently, lymph node metastasis may not necessarily function as an independent prognostic factor, and the inferior prognosis observed in lymph node-positive patients could potentially be attributed to the presence of preexisting microscopic distant metastasis.

Several researchers have made analyses on the prognosis of renal cancer patients with VTT, and some studies have suggested that risk factors abound such as lymphatic invasion, perirenal adipose tissue invasion, tumor diameter, tumor necrosis, preoperative hemoglobin, pathological type, tumor thrombus grading (9,39-41). Meanwhile, these influential factors remain many controversies, needing to be further explored. This study analyzed clinical data of 114 non-metastasis renal cancer patients with VTT, and performed Cox regression analyses on potential factors. Results indicated that tumor diameter, preoperative LDH and preoperative neutrophils were independent risk factors for OS. A new nomogram model was constructed based on the above data and Mayo grading to quantified and analyzed prognosis, helping clinicians to guide treatment and discuss decisions with patients.

Compared with the traditional tumor-node-metastasis (TNM) staging system, nomograms based on results of multivariate analysis were proved to better predict prognosis in various cancer types (42,43). Multivariate Cox regression analysis indicated that tumor diameter, preoperative LDH and preoperative neutrophils were independent risk factors for OS in non-metastasis renal cancer with VTT and these three factors binding Mayo grading effectively predict prognosis. The nomogram is verified to be effective via Kaplan-Meier (KM) plots, ROC and DCA curve. The AUC of 1- and 2-year OS is 0.82 and 0.67, but long-term OS is only 0.57 and 0.55 considering the limited number of samples and different treatment method after surgery in this study. Most importantly, the indicator that our nomogram

included are easy to gain, embodying the practicality, availability and economy of the nomogram.

The present study is subject to several limitations: Firstly, it should be noted that only cases with complete clinical and follow-up data were included in this study, which may introduce a certain degree of selection bias. Secondly, the survival model established in this study may not be generalizable to patients with RCC accompanied by VTT who exhibit extremely abnormal laboratory parameters. Additionally, due to the relatively low number of patients in our cohort who had tumor thrombus invading the vein wall, they were not included in the study. Lastly, it is important to acknowledge that the survival model established based on our dataset has yet to undergo validation using patient data from other institutions. Therefore, further validation of the model is warranted through large-scale, multi-center, and prospective studies. This will help to enhance the robustness and generalizability of the findings and ensure the reliability of the survival model in clinical practice.

When considering the collective impact of various factors on the OS of non-metastatic renal cancer patients with VTT, several independent risk factors stand out. These include renal tumor diameter, preoperative LDH, and preoperative neutrophil. By taking these factors into account, clinicians can gain valuable insights into the prognosis of patients and make informed decisions regarding their treatment. To enhance the accuracy of prognosis prediction, a nomogram model can be developed. This model combines the aforementioned risk factors with the traditional Mayo grading system, allowing for a comprehensive and quantitative analysis of prognosis. The nomogram serves as a visual tool that enables clinicians to estimate the likelihood of survival and identify patients at a higher risk of adverse outcomes. By utilizing this nomogram model, clinicians can more effectively identify highrisk patients who require early intervention and tailored treatment strategies. This proactive approach not only improves the individualized care provided to patients but also enhances overall treatment outcomes. Ultimately, the integration of the nomogram model with traditional Mayo grading empowers healthcare professionals to make more precise prognostic assessments and deliver optimal care to renal cancer patients with VTT.

Conclusions

In a word, surgical intervention improves the prognosis of renal cancer patients with VTT. Independent risk

factors include kidney tumor diameter, preoperative LDH levels, and preoperative neutrophil count. The nomogram enhances the Mayo grading system and serves as a dependable tool for prognostic evaluation in renal cancer patients with VTT.

Acknowledgments

Funding: None.

Footnote

Reporting Checklist: The authors have completed the TRIPOD reporting checklist. Available at https://tau.amegroups.com/article/view/10.21037/tau-23-341/rc

Data Sharing Statement: Available at https://tau.amegroups.com/article/view/10.21037/tau-23-341/dss

Peer Review File: Available at https://tau.amegroups.com/article/view/10.21037/tau-23-341/prf

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at https://tau.amegroups.com/article/view/10.21037/tau-23-341/coif). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013). This study was approved by the Ethics Committee of Changhai Hospital (No. CHEC2021-191). Informed consent was waived by Ethics Committees due to the retrospective nature of the study.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the noncommercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: https://creativecommons.org/licenses/by-nc-nd/4.0/.

References

- 1. Siegel RL, Miller KD, Wagle NS, et al. Cancer statistics, 2023. CA Cancer J Clin 2023;73:17-48.
- 2. Klatte T, Rossi SH, Stewart GD. Prognostic factors and prognostic models for renal cell carcinoma: a literature review. World J Urol 2018;36:1943-52.
- Lardas M, Stewart F, Scrimgeour D, et al. Systematic Review of Surgical Management of Nonmetastatic Renal Cell Carcinoma with Vena Caval Thrombus. Eur Urol 2016:70:265–80.
- 4. Reese AC, Whitson JM, Meng MV. Natural history of untreated renal cell carcinoma with venous tumor thrombus, Urol Oncol 2013;31:1305-9.
- Cost NG, Delacroix SE Jr, Sleeper JP, et al. The impact of targeted molecular therapies on the level of renal cell carcinoma vena caval tumor thrombus. Eur Urol 2011;59:912-8.
- Di Silverio F, Sciarra A, Parente U, et al. Neoadjuvant therapy with sorafenib in advanced renal cell carcinoma with vena cava extension submitted to radical nephrectomy. Urol Int 2008:80:451-3.
- Chen Z, Yang F, Ge L, et al. Outcomes of renal cell carcinoma with associated venous tumor thrombus: experience from a large cohort and short time span in a single center. BMC Cancer 2021;21:766.
- 8. Shiff B, Breau RH, Mallick R, et al. Prognostic significance of extent of venous tumor thrombus in patients with non-metastatic renal cell carcinoma: Results from a Canadian multi-institutional collaborative. Urol Oncol 2021;39:836.e19-27.
- Zapała Ł, Sharma S, Kunc M, et al. Analysis of Clinicopathological Factors Influencing Survival in Patients with Renal Cell Carcinoma and Venous Tumor Thrombus. J Clin Med 2021;10:3852.
- Tabbara MM, González J, Martucci M, et al. Current Approaches in Surgical and Immunotherapy-Based Management of Renal Cell Carcinoma with Tumor Thrombus. Biomedicines 2023;11:204.
- Karnes RJ, Blute ML. Surgery insight: management of renal cell carcinoma with associated inferior vena cava thrombus. Nat Clin Pract Urol 2008;5:329-39.
- 12. Zhao X, Yan Y, Dong JH, et al. Influence of Deep Invasive Tumor Thrombus on the Surgical Complexity and Prognosis of Patients With Non-Metastatic Renal Cell Carcinoma Combined With Venous Tumor Thrombus.

- Front Oncol 2022;12:833780.
- Wang B, Li H, Ma X, et al. Robot-assisted Laparoscopic Inferior Vena Cava Thrombectomy: Different Sides Require Different Techniques. Eur Urol 2016;69:1112-9.
- Wang B, Li H, Huang Q, et al. Robot-assisted Retrohepatic Inferior Vena Cava Thrombectomy: First or Second Porta Hepatis as an Important Boundary Landmark. Eur Urol 2018;74:512-20.
- 15. Tian X, Hong P, Liu Z, et al. En bloc retroperitoneal laparoscopic radical nephrectomy with inferior vena cava thrombectomy for renal cell carcinoma with level 0 to II venous tumor thrombus: A single-center experience. Cancer 2020;126 Suppl 9:2073-8.
- Qu L, Chen H, Chen Q, et al. Development and validation of a prognostic model incorporating tumor thrombus grading for nonmetastatic clear cell renal cell carcinoma with tumor thrombus: A multicohort study. MedComm (2020) 2023;4:e300.
- Wang K, Liu Z, Hong P, et al. T.H.R.O.B.V.S. Score A Comprehensive Model to Predict the Surgical Complexity of Renal Cell Carcinoma With Tumor Thrombus. Front Oncol 2022;12:900550.
- Lue K, Russell CM, Fisher J, et al. Predictors of Postoperative Complications in Patients Who Undergo Radical Nephrectomy and IVC Thrombectomy: A Large Contemporary Tertiary Center Analysis. Clin Genitourin Cancer 2016;14:89-95.
- Campi R, Barzaghi P, Pecoraro A, et al. Contemporary techniques and outcomes of surgery for locally advanced renal cell carcinoma with focus on inferior vena cava thrombectomy: The value of a multidisciplinary team. Asian J Urol 2022;9:272-81.
- Bhindi B, Thompson RH, Lohse CM, et al. The Probability of Aggressive Versus Indolent Histology Based on Renal Tumor Size: Implications for Surveillance and Treatment. Eur Urol 2018;74:489-97.
- 21. Yang C, Shuch B, Serrano M, et al. Adverse Histopathologic Characteristics in Small Clear Cell Renal Cell Carcinomas Have Negative Impact on Prognosis: A Study of 631 Cases With Clinical Follow-up. Am J Surg Pathol 2019;43:1413-20.
- Hötker AM, Karlo CA, Zheng J, et al. Clear Cell Renal Cell Carcinoma: Associations Between CT Features and Patient Survival. AJR Am J Roentgenol 2016;206:1023-30.
- Abel EJ, Masterson TA, Karam JA, et al. Predictive Nomogram for Recurrence following Surgery for Nonmetastatic Renal Cell Cancer with Tumor Thrombus. J Urol 2017;198:810-6.

- 24. Chen X, Li S, Xu Z, et al. Clinical and oncological outcomes in Chinese patients with renal cell carcinoma and venous tumor thrombus extension: single-center experience. World J Surg Oncol 2015;13:14.
- 25. Abel EJ, Thompson RH, Margulis V, et al. Perioperative outcomes following surgical resection of renal cell carcinoma with inferior vena cava thrombus extending above the hepatic veins: a contemporary multicenter experience. Eur Urol 2014;66:584-92.
- Abel EJ, Margulis V, Bauman TM, et al. Risk factors for recurrence after surgery in non-metastatic RCC with thrombus: a contemporary multicentre analysis. BJU Int 2016;117:E87-94.
- Martínez-Salamanca JI, Huang WC, Millán I, et al.
 Prognostic impact of the 2009 UICC/AJCC TNM staging
 system for renal cell carcinoma with venous extension. Eur
 Urol 2011;59:120-7.
- 28. Haddad AQ, Leibovich BC, Abel EJ, et al. Preoperative multivariable prognostic models for prediction of survival and major complications following surgical resection of renal cell carcinoma with suprahepatic caval tumor thrombus. Urol Oncol 2015;33:388.e1-9.
- Sheng SL, Liu JJ, Dai YH, et al. Knockdown of lactate dehydrogenase A suppresses tumor growth and metastasis of human hepatocellular carcinoma. FEBS J 2012;279:3898-910.
- 30. Matrana MR, Bathala T, Campbell MT, et al. Outcomes of unselected patients with metastatic clear-cell renal cell carcinoma treated with first-line pazopanib therapy followed by vascular endothelial growth factor receptor tyrosine kinase inhibitors or mammalian target of rapamycin inhibitors: a single institution experience. BJU Int 2016;118:264-71.
- 31. Obata J, Tanaka N, Mizuno R, et al. Plasma fibrinogen level: an independent prognostic factor for disease-free survival and cancer-specific survival in patients with localised renal cell carcinoma. BJU Int 2016;118:598-603.
- Parosanu AI, Pirlog CF, Slavu CO, et al. The Prognostic Value of Neutrophil-to-Lymphocyte Ratio in Patients with Metastatic Renal Cell Carcinoma. Curr Oncol 2023;30:2457-64.
- Ghatalia P, Gordetsky J, Kuo F, et al. Prognostic impact of immune gene expression signature and tumor infiltrating immune cells in localized clear cell renal cell carcinoma. J Immunother Cancer 2019;7:139.
- Capitanio U, Suardi N, Matloob R, et al. Extent of lymph node dissection at nephrectomy affects cancerspecific survival and metastatic progression in specific sub-

- categories of patients with renal cell carcinoma (RCC). BJU Int 2014;114:210-5.
- 35. Capitanio U, Becker F, Blute ML, et al. Lymph node dissection in renal cell carcinoma. Eur Urol 2011;60:1212-20.
- 36. Marchioni M, Bandini M, Pompe RS, et al. The impact of lymph node dissection and positive lymph nodes on cancer-specific mortality in contemporary pT(2-3) nonmetastatic renal cell carcinoma treated with radical nephrectomy. BJU Int 2018;121:383-92.
- Bhindi B, Wallis CJD, Boorjian SA, et al. The role of lymph node dissection in the management of renal cell carcinoma: a systematic review and meta-analysis. BJU Int 2018;121:684-98.
- Karmali RJ, Suami H, Wood CG, et al. Lymphatic drainage in renal cell carcinoma: back to the basics. BJU Int 2014;114:806-17.
- 39. Cao C, Bi X, Liang J, et al. Long-term survival and prognostic factors for locally advanced renal cell

Cite this article as: Zhu B, Yang Y, Huangfu Z, Zhang W, Jiang A, Wang L. Construction of the prognostic model in non-metastatic renal cancer patients with venous tumor thrombus. Transl Androl Urol 2023;12(11):1645-1657. doi: 10.21037/tau-23-341

- carcinoma with renal vein tumor thrombus. BMC Cancer 2019;19:144.
- 40. Tilki D, Hu B, Nguyen HG, et al. Impact of synchronous metastasis distribution on cancer specific survival in renal cell carcinoma after radical nephrectomy with tumor thrombectomy. J Urol 2015;193:436-42.
- 41. Shang B, Guo L, Shen R, et al. Prognostic Significance of NLR About NETosis and Lymphocytes Perturbations in Localized Renal Cell Carcinoma With Tumor Thrombus. Front Oncol 2021;11:771545.
- 42. Ruan R, Chen S, Tao Y, et al. Retrospective analysis of predictive factors for lymph node metastasis in superficial esophageal squamous cell carcinoma. Sci Rep 2021;11:16544.
- 43. Chen X, Guo T. Development of a nomogram for predicting survival of patients with ovarian serous cystadenocarcinoma after based on SEER database. Zhejiang Da Xue Xue Bao Yi Xue Ban 2021;50:369-74.

Supplem	•			•																				
-	More detailed c	_ength	ASA	Intraoperative		Age	BMI	Hypertension	Diabetes	Preoperative	Preoperative	Preoperative				Tumor	Clavien-Dindo	Lymphatic metastasis	Perirenal adipose	Pathological type		Postoperative	Outcome	OS
number		(days)	scoring (1, 2, 3, 4, 5)		(female 0, male 1)		(kg/m²)	(no 0, yes 1)	(no 0, yes 1)	LDH (U/L)	(µmol/L)	neutrophils (×109/L)	(g/L)	(left 0, right 1)	surgery 1)	thrombus	grading (grade I 0, grade II/III 1)	(non-metastasis 0, metastasis 1)	tissue invasion (non- invasion 0, invasion 1)		(mm)	targeted therapy (no 0, yes 1)	(alive 0, dead 1)	(months)
1	4.7 4.1	16 12	2 2	2,000 1,000	1 0	61 71	23.1 29.1	0 1	0	139 153	94 86	4 5.8	155 94	1	0 1	III 0	1 0	0	0	0	90 60	0	1	91 21
3	5.1 4.4	17 22	2	200 2,400	0 1	61 62	19.3 27	1 1	0	202 383	49 124	2.9 6	86 158	0 1	0	IV 0	0	0	0 1	0 1	50 60	1 0	0	95 35
5	1.6	7	2	50	1	58	24.3	1	1	173	64	5.6	137	0	1	0	0	0	1	0	55	0	0	64
6 7	3.3 3.3	12 15	2 1	4,500 2,000	1 1	65 21	21.5 23.1	0	0	175 217	76 84	5 3.3	135 123	0 1	0	III II	1	0	1 0	1 1	50 145	0 1	0	60 57
8	3.9	7	2	1,200	1	59	24.7	1	0	176	79	6.1	136	0	0	0	1	1	1	1	73	0	0	56
9 10	2.3 4.5	8	2 2	50 1,600	1	60 33	24.2 24.4	0	0	178 176	99 77	3.6 2.9	145 129	1	1 0	O III	0 1	0	0	0 1	60 100	0 1	0	50 24
11 12	3.0 3.0	15 14	3	4,500 800	1	54 67	27.1 27.8	1	1	259 287	125 120	7.7 3.5	97 119	0	0	III III	1	0	0	0	55 100	0	0	31 6
13	2.6	11	2	1,000	0	69	23.6	1	0	353	129	5.3	73	1	0	11	0	0	1	0	100	1	1	38
14 15	3.2 1.8	18 10	2	1,100 1,000	1	65 49	23.3 21.8	0	0	189 163	84 55	4.8 4.5	155 150	0	1	0	0	0	1	0	70 110	0	0	52 44
16	2.6	8	2	3,500	1	70	21.8	0	0	202	101	5.4	110	1	0	II	0	0	0	0	70	0	1	27
17 18	4.9 2.2	12 8	2	800 3,000	1 1	48 52	24.6 24.7	0	0	185 141	60 123	2.9 3.7	135 93	0	0	l III	1	0	0	1 0	65 75	1	0	25 24
19	2.5	17	2	3,400	1	56	28.2	0	0	156	89	5.3	111	0	0	II	1	0	0	0	40	0	0	23
20 21	2.5 2.0	15 9	2	2,500 200	1 1	58 45	22.7 21.7	0	0	188 204	110 69	8 6	72 111	0 1	0	IV 0	1 0	1 0	0 1	0	160 75	0 1	1	9 16
22	3.7	12	2	100	1	57	28.4	0	0	198	86	3.2	161	1	1	I	0	0	0	0	70	1	0	68
23 24	3.6 4.0	7 15	2 3	300 50	1	51 84	19.8 30	0 1	0 1	177 305	72 105	4.9 6.3	145 103	1	0	0	0 1	0	0	0	100 80	1	0 1	99 20
25	4.3	11	2	100	1	43	22.6	0	0	139	77	4.8	142	1	1	0	0	0	0	0	45	1	0	114
26 27	2.0 5?	21 17	2 2	1,000 50	1	51 57	29.9 23.5	0	1	165 379	133 56	3 3.8	137 117	0	1	0	1 0	0	0	0	30 60	1 0	1	120 25
28	3.8	7 10	2	750 500	1	53 63	27	1 0	0	170	83	2.5 2.8	128 106	1	1	II III	0	0	0	0	62 105	0	0	19 20
29 30	2.3 3.2	21	2 3	2,300	1	65	24.2 20.3	0	0	145 259	72 125	3	121	1	0	III	1	1	1	1	95	0	1	4
31 32	2.5 2.4	7 16	2	300 600	0	28 61	17.8 23	0	0	166 205	66 87	4.6 5.7	77 97	1	0	0 II	1	1	1 0	1 0	100 110	0	0	53 43
33	3.0	29	2	1,200	0	73	27.9	0	0	197	92	5.6	102	0	0	II	1	0	1	0	140	0	1	48
34 35	2.5 5.0	13 17	2	1,300 2,000	1 1	67 65	21.2 23.2	1 0	0	208 178	77 82	1.2 5.8	90 118	1 0	0 1	III IV	1 1	0	0 1	0 1	40 70	0 1	0	49 12
36	3.5	16	2	800	1	77	21.4	0	0	148	134	3.9	132	0	0	III	1	0	0	0	60	1	1	69
37 38	3.5 3.2	14 9	2	2,000 2,100	1 1	52 58	27.9 22.4	1	0	257 159	145 109	7.1 4.2	126 104	0 1	0	II III	0	1 0	1 1	0	90 85	1 1	0 1	19 6
39	3.3	7	2	200	0	54	28	0	0	165	66	3.8	129	0	1	0	0	0	0	0	60	0	0	24
40 41	2.3 3.3	9	3 3	50 900	1 0	79 69	20.8 21.1	1 0	0	81 164	89 73	2.9 3	81 157	1	0	O II	0 1	0	0	0	95 130	1	0	6 28
42 43	3.3 2.8	28 16	2	2,600 100	1	85 53	24.2 31.2	0	0	169 149	127 97	2.8 3.7	89 135	1	0	II O	1	0	0	0	62 45	1	0	34 33
44	2.8	8	2	1,400	0	62	18.1	0	0	165	87	6.8	92	1	0	III	1	0	1	0	95	1	0	35
45 46	3.9 3.1	5 10	2	200 1,500	1 1	54 57	24.3 22.8	1 0	0	192 268	75 104	4.7 4.4	110 121	1 1	1 0	0 III	0	0	1	1 0	81 77	0 1	0	41 42
47	2.0	7	2	100	0	68	23.5	0	0	233	53	2.7	147	0	1	0	0	0	0	0	55	0	0	45
48 49	7.8 6.4	11 21	1 4	2,500 4,500	1 0	36 72	29.1 27.8	0 1	0	211 257	118 73	3.6 3.2	120 127	0 1	1 0	II IV	1 1	0	0 1	0	90 120	1 1	0 1	59 43
50	2.0	15	2	1,400	1	69	22.1	0	0	162	70	2.1	148	1	0	III	1	0	1	0	80	1	1	16
51 52	4.3 6.8	10 30	3 2	200 1,200	1 0	64 72	28.3 20.4	1	1	150 159	85 68	4.5 5.7	120 88	0 1	1 1	O III	0 1	0	0 1	0	70 80	0	0 1	119 49
53	5.0	10	3	1,200	1	54	17.6	0	0	131	75	5.29	127	1	0	0	1	0	0	0	80	1	1	29
54 55	4.0 4.0	17 15	4 2	700 500	1	54 66	18.6 22	0	0	95 153	93 138	6.4 2.4	89 138	1	0	0	0	0	0	0	90 45	1	0	56 117
56 57	6.5 6.0	10 21	1 2	1,300 3,000	1	21 39	20 24.8	0	0	198 190	90 92	3.88 4.1	119 102	1	0	l III	1	1	0	1 0	120 80	0	1	48 120
58	2.3	7	2	1,400	0	74	22.9	0	0	180	90	3.7	122	1	0	1	1	0	0	0	35	0	0	44
59 60	4.0 2.6	14 18	3	2,000 1,000	1 0	63 40	18.6 21.2	0	0	128 153	66 65	8.3 3.1	84 129	1 1	0	l I	0	1 0	0 1	0	130 80	1 1	1 1	33 40
61	2.8	16	2	3,000	1	65	24.5	0	1	121	95	3	92	0	0	I	1	0	1	0	90	0	1	36
62 63	5.7 3.7	25 8	2	5,500 3,000	1 1	51 69	24.2 18	0	0	112 135	123 137	5.6 5.7	99 88	1 0	0 0	III III	0	0	0	0 1	170 80	1 0	1	5 18
64	5.6	20	2	800	0	58	21.2	0	0	168	62	5.4	92	0	0	III	1	0	0	0	27	1	1	118
66	3.0 3.5	15 17	2 2	1,000 2,200	0	76 46	19.8 24	1 0	0	113 204	103 81	7.7 5	86 113	0	0	II II	0	1 0	1	0	60 110	0	1	21 24
67	8.5 3.4	18 10	2	800 50	0	64 59	18.4	0	0	128	52	4.8 3.4	96	1	0	II	1	0	1	0	100 40	0	1	13
69	2.5	22	2 3	100	1	75	25.8 25.7	0 1	0	147 171	71 77	5.2	158 142	0	1	0	1	0	1	0	40	0	1	41
70 71	2.5 5.6	14 13	2	3,100 4,000	1 0	63 58	25.7 22.2	1	0	168 171	83 66	3.6 5.8	106 115	1 0	0	II 0	1	0	0	0	60 110	0	1	19 13
72	3.7	12	2	200	1	41	19.2	0	0	484	118	4	106	0	0	0	0	0	0	1	140	0	1	30
73 74	3.9 3.1	20 16	2	1,000 500	0	71 37	24.4 18.1	1 0	0	368 182	87 58	8.1 5.1	83 111	1 1	0	I 0	0 1	0	1 0	1 0	130 70	0	1	20 33
75	2.5	10	2	100	1	68	21	0	0	175	106	3.3	90	1	1	0	0	0	0	0	80	0	0	91
76 77	6.3 3.0	20 19	2	4,000 400	1	64 63	24.8 23.2	0 1	0	154 268	85 117	3.7 3.1	107 208	0	0	III I	1 0	1 0	1 0	0	50 130	1 0	1	8 15
78	4.3	15	2	4,800	1	68	24.3	1	0	161	94	4.4	119	1	0	II 	1	0	0	1	85	0	0	71
79 80	5.8 4.0	16 12	2	2,000 20	1	53 59	20.4	0	0	206 149	109 54	5.4 4.6	116 96	0	0 1	0	1	0	1	0	75 85	0	0	20 68
81 82	3.0 2.1	8 18	2	300 200	1	54 53	19.3 20.9	0	0	150 211	75 76	4.4 2.8	144 120	1	0	l O	0	0	0	0	30 105	0	0	62 60
83	1.8	8	2	230	0	46	25	0	1	148	52	1.1	101	1	0	0	0	0	0	0	48	0	1	40
84 85	2.6 3.1	7 15	2	200 4,000	1 1	61 52	20.8 24.7	0	0	185 152	84 58	5.5 7.4	125 93	0	1 0	0 II	0 1	0	0	0	70 80	1 1	0	32 5
86	2.2	15	3	2,500	0	55	16.8	0	0	226	75	3.3	57	0	0	0	1	1	1	1	130	1	1	11
87 88	3.2 2.5	11 7	2	400 500	0 1	66 58	19.4 26	0	0	246 148	75 88	4.3 3	109 126	0	1 1	II 0	1 0	0	0	0	65 55	1 0	1	28 44
89	2.5	9	1	1,200	1	33	30.6	0	0	283	101	4.9	129	1	0	III	1	0	0	1	60	0	1	26
90 91	4.1 2.0	15 7	2	2,000 100	0 1	39 51	21.2 26.7	0 1	0 1	913 142	54 55	2.2 6.3	106 164	1 0	0 1	IV 0	1 0	0	1 0	0	205 35	0	1 0	10 29
92	1.8	6	2	20	1	66	24.3	1	0	143	78	6.2	128	1	1	0	0	0	0	0	50	0	0	23
93 94	4.3 2.5	7 8	2	400 4,500	1	60 65	24.2 27.4	1	0	164 184	115 93	5 3.6	140 133	0 1	1 0	0 III	0 1	1 0	0	1 0	100 80	0 1	0	21 2
95	2.3	15	3	3,000	0	67	21.6	1	0	212	65	2.3	124	0	0	IV	1	0	0	0	50	1	0	4
96 97	3.0 2.7	11 10	2	1,500 1,200	1	65 77	18.9 24.9	1 0	0	108 173	97 109	3.3 3.1	75 75	1	1 0	II II	1	0	1	0	90 90	0	0	5 5
98	2.3	13	3	3,000	1	53	23.5	0	0	108	100	6.5	67	0	0	III	1	0	1	0	135	0	0	13
99 100	3 1	13 13	3 1	1,400 500	1	52 59	28 22.5	1 0	0	257 158	145 94	7.05 3.09	126 145	0	0	0	0	1 0	1 0	0	90 45	1 1	0 1	19 100
101	2	14 8	2	50	1	74 87	22.6	1	1	164 342	117 75	5.57 5.67	131	0	1	0	0	0	0	0	90 60	1	1	48
102 103	2	8 13	2	100 1,200	1	87 59	20.3 27.5	0 1	0	342 550	75 83	5.67 4.7	127 130	0	1 0	U I	υ 1	0	0	U 1	60 48	υ 1	1	40 50
104 105	2	14 12	2	1,200 1,200	1	60 58	25 24.8	0	0	101 138	67 74	6.35 4.57	137 126	0	0	0	1	0	0	1	55 110	1	1	78 96
105	2	6	2	1,200 200	1	58 56	26.8	0	0	138 145	74 75	4.57 5.96	126	1	0	0	0	0	0	0	110 70	0	0	96 102
107 108	2	12 8	2	1,600 80	1	43 51	22.1 26.8	0	0	220 163	89 65	6.25 4.68	130 163	0	0 1	0	1	0	1	1 0	38 70	1 1	1	42 70
109	2	6	2	1,500	0	67	25.2	1	0	194	75	3.51	164	0	0	0	1	1	1	0	113	1	0	72
110 111	2	8	2	2,000 100	1	44 58	25.9 24.2	0 1	0	150 152	88 85	5.94 2.73	137 129	0	0 1	0	0	0	1 0	0	90 50	1 0	1	60 45
112	3	8	3	1,300	1	43	23.2	0	0	277	87	2.6	141	1	0	II	1	1	0	0	60	1	1	40
113 114	3 2	7 10	3	200 1,000	1 1	60 73	18 27.3	0 1	0	142 166	98 89	2.34 2.95	140 135	1	0 1	0 II	0	1 0	0 1	0	75 50	1	0	80 76
ASA, Ame	rican Society o	of Anesth	nesiologists;	BMI, body mas	ss index; L	DH, lactat	te dehydro	genase; ccRC0	C, clear cell r	enal cell carc	inoma; OS, ov	erall survival.	_	_										