



Characterizing the pharmacokinetics and pharmacodynamics of immunosuppressant medicines and patient outcomes in elderly renal transplant patients

Amelia R. Cossart¹, W. Neil Cottrell¹, Scott B. Campbell², Nicole M. Isbel², Christine E. Staatz¹

¹School of Pharmacy, University of Queensland, Brisbane, Australia; ²Department of Nephrology, University of Queensland at the Princess Alexandra Hospital, Brisbane, Australia

Contributions: (I) Conception and design: All authors; (II) Administrative support: None; (III) Provision of study material or patients: None; (IV) Collection and assembly of data: AR Cossart, WN Cottrell, CE Staatz; (V) Data analysis and interpretation: All authors; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Amelia R. Cossart. School of Pharmacy, University of Queensland, 20 Cornwall Street, Woolloongabba QLD 4102, Australia. Email: amelia.cossart@uqconnect.edu.au.

Abstract: This review examines what is currently known about the pharmacokinetics and pharmacodynamics of commonly prescribed immunosuppressant medicines, tacrolimus, cyclosporine, mycophenolate and prednisolone, in elderly renal transplant recipients, and reported patient outcomes in this cohort. Renal transplantation is increasing rapidly in the elderly, however, currently, long-term patient outcomes are relatively poor compared to younger adults. Some studies have suggested that elderly recipients may have higher dose-adjusted exposure and/or lower clearance of the calcineurin inhibitors tacrolimus and cyclosporine; with one study reporting up to 50% reduction in tacrolimus exposure in the elderly. Elderly transplant recipients do not appear to have higher dosage-adjusted exposure to mycophenolic acid (MPA). The effects of ageing on the pharmacokinetics of prednisolone are unknown. Only one study has examined how aging effects drug target enzymes, reporting no difference in baseline inosine 5'-monophosphate dehydrogenase (IMPDH) activity and MPA-induced IMPDH activity in elderly compared to younger adult renal transplant recipients. In elderly transplant recipients, immunosenescence likely lowers the risk of acute rejection, but increases the risk of drug-related adverse effects. Currently, the three main causes of death in elderly renal transplant recipients are cardiovascular disease, infection and malignancy. One study has showed that renal transplant recipients aged over 65 years are seven times more likely to die with a functioning graft compared with young adults (aged 18–29 years). This suggests that an optimal balance between immunosuppressant medicine efficacy and toxicity is not achieved in elderly recipients, and further studies are needed to foster long-term graft and patient survival. Lower maintenance immunosuppressant targets in elderly recipients may decrease patient susceptibility to drug side effects, however, further studies are required and appropriate targets need to be established.

Keywords: Immunosuppressant medicines; elderly; renal transplantation; pharmacokinetics; pharmacodynamics; patient outcomes

Submitted Oct 10, 2018. Accepted for publication Oct 19, 2018.

doi: 10.21037/tau.2018.10.16

View this article at: <http://dx.doi.org/10.21037/tau.2018.10.16>

Introduction

Renal transplantation is first-line treatment for patients with end stage kidney disease (ESKD), regardless of age (1,2). When compared to dialysis, renal transplantation increases life expectancy, as well as improves physical and social functioning (quality of life), mental stability, vitality, overall well-being and future perspective (2-5). The fastest growing age group with ESKD is patients aged over 65 years (4,6). In the last decade, there has been a noticeable increase in the number of elderly patients receiving a renal transplant. In the USA, the number of transplants performed in patients aged over 65 years increased from 10% to 17% of the total population between 2002 and 2012 (7). In Australia, 14% of renal transplants performed in 2015 were in patients aged 65 years of age and above (8). In Europe, a 2012 Renal Transplantation Registry report stated that transplant prevalence was 22% in the over 65 years age group and 20% in the over 75 years age group (9). This rise in the incidence of renal transplantation in the elderly can be attributed to an aging population (10), improvements in transplant outcomes (11), and the introduction of expanded criteria donor (ECD) kidneys (12).

Although age is no longer considered an absolute barrier to transplantation, long-term graft and patient survival is not improving in elderly renal transplant recipients, as compared to younger adults (1,6,10,13). The three most frequent causes of death in elderly recipients are infection, malignancy and cardiovascular events (14-16), all of which can be attributed partly to immunosuppressant medications. Studies show that older recipients have a lower risk of acute rejection due to reduced immune reactivity (15) and the elderly are the most likely age group to die with a functioning graft (1,17,18). The risk of acute rejection increases with ECD kidneys, and although elderly patients are considered at lower risk of rejection compared to younger adults, they are still at risk (19,20). Additionally, age-related comorbidities negatively influence patient outcomes in elderly recipients (13). Consequently, elderly renal transplant recipients are at greater risk of both morbidity and mortality when compared to younger recipients (21,22).

The intricate balance between under- and over-immunosuppression becomes increasingly complex in elderly recipients, due to changes in drug pharmacokinetics and pharmacodynamics (23-25). Responses seen in the elderly can be unexpected and exaggerated when compared to younger adults with similar characteristics

(gender, weight, height, ethnicity) (25). Larger variability in response may be due to changes in physiology and disease, however, as studies conducted in the elderly are limited, it is difficult to draw definitive conclusions (25). Obtaining a greater understanding of immunosuppressant drug pharmacokinetics and pharmacodynamics in elderly renal transplant recipients should assist with therapeutic decisions in this unique and vulnerable sub-population. Therefore, the purpose of this review is to critique and summarize the available literature on the pharmacokinetics and pharmacodynamics of immunosuppressant medicines in elderly renal transplant recipients.

ECD kidneys

Elderly kidney transplant recipients, who receive ECD kidneys, incur additional barriers to graft survival, which compound the already significant effects of physiological aging. ECD kidneys have a shortened lifespan, with both one and five-year graft survival significantly lower than with standard donor kidneys (26). Allocation of grafts with shortened lifespans (ECD kidneys) to patients with shorter life expectancies (the elderly) is considered acceptable as the survival and quality-of-life benefits of old-for-old donor matching still outweighs long-term dialysis (14,15,17,20,27-29).

The decline in immune reactivity observed in elderly recipients is complicated by increased immunogenicity of older donor grafts (15). This has resulted in more potent early stage immune responses observed in elderly transplant recipients with ECD kidneys, which can lead to episodes of acute rejection (21) and delayed graft function (27). In one retrospective study involving 1,307 participants, it was shown that receiving an ECD kidney significantly increased the risk of acute rejection, but had no effect on overall patient survival in patients aged 60 years and over (30).

Age-related pharmacokinetic (PK) changes

A number of physiological changes associated with advancing age can affect drug absorption, distribution, metabolism and excretion in healthy adults. Decreased surface area of the intestinal epithelium, reduced splanchnic blood flow, altered gastric motility (31), delayed gastric emptying and increased gastric pH (but reduced gastric acid secretion) (31,32), can potentially lead to reduced drug absorption in the elderly. However, studies have shown that the bioavailability of drugs absorbed through permeation of the gastrointestinal tract remains largely unchanged in

the elderly; with absorption of drugs via carrier-mediated transport processes most likely to be impeded (24). The absorption of basic drugs may actually improve, as the rise in gastric pH, caused by a decline in gastric parietal cells, improves absorption (24). Overall, it is assumed that there is a slight decline in drug absorption in the elderly; however, this is patient and drug specific.

Elderly patients tend to have an increased percentage of body fat, and a reduction in lean body muscle mass and total body water, compared to younger adults (24,33). These body composition changes generally occur in all elderly patients, including 'thin' body types (34), resulting in a volume of distribution increase for lipophilic drugs and decrease for hydrophilic drugs (31). A drug's concentration in the plasma is inversely related to its volume of distribution, leading to a greater plasma concentration in elderly patients compared to younger adults for hydrophilic drugs. Elderly patients can also have altered drug binding to plasma proteins (albumin, alpha-1-acid glycoprotein and lipoproteins) (32,35). A reduction in serum albumin in the elderly can lead to a decrease in total drug concentration and an increase in free drug fraction, with no corresponding change in unbound (free) drug concentration. Changes in drug protein binding may be an important consideration when interpreting therapeutic drug monitoring (TDM) results for drugs where total rather than unbound drug concentration is measured (25,35).

There are a number of transporters involved in the movement of drugs across biological membranes within the body. Solute carrier (SLC) transporters mediate the uptake of drugs from the blood, transporting them across the basolateral membrane, and into cells (proximal tubule of the kidney or into hepatocytes), or the gut (into enterocytes) (36), whilst ATP-binding cassette transporters (ABC transporters) are involved in the cellular efflux of drugs and their metabolites (37). As many transporters are multi-specific, drug-drug interactions that affect drug membrane transportation can lead to altered pharmacokinetics and drug effects (37). There is some research that suggests that protein carrier activity is influenced by age, with a study by Fu, Csanaky and Curtis in 2012 showing that the mRNA levels of xenobiotic-processing genes (includes SLCs) was significantly lower in aged mice. Similarly, a study by van Assema and researchers showed that aging reduced P-glycoprotein levels in the blood-brain barrier in healthy adults (38). When considering transporter effects in the elderly, the presence and impact of multiple comorbidities or concomitant medications can

affect these transporters in extremely complex ways, making it difficult to predict the final outcome.

Advancing age leads to a decline in liver size, which can cause a reduction in hepatic blood flow (approximately 40%), and reduced drug transfer across the hepatic endothelium (23,39). More specifically, drug metabolism is directly attributed to the drug's extraction rate (high or low) by the liver, as haemoperfusion declines in the elderly; drugs with a high extraction rate have an age-related decline in clearance. Conversely, clearance of drugs with a low extraction rate remain largely unchanged, as metabolism is less dependent on hepatic blood flow than activity of membrane bound metabolising enzymes such as cytochrome P450 (CYPs) (32). Studies have shown that although it is unlikely CYP3A enzyme number or affinity reduces with age (40), drug metabolism via Phase-I CYP3A-mediated reactions may decline with advancing age (39). This is likely due to inter- morphological and circulatory changes (40), and can be variable between patients (32). Phase-II conjugation reactions (primarily glucuronidation) and the uridine-5'-diphospho glucuronyl transferase (UGT) system (in the liver) are generally less influenced by age (32,39).

In terms of drug elimination, glomerular filtration rate (GFR) declines at a rate of approximately 1 mL/min/1.73 m² of body surface area, from 20–30 years of age onwards (31,34). As a result, renal drug excretion is often significantly reduced in the elderly, regardless of comorbidities. Little is known about whether biliary excretion and enterohepatic recirculation of drugs is affected by age.

A potential decline in liver, and likely decline in renal function in the elderly can cause a reduction in drug clearance, compared with younger adults. Drug clearance is the principle determinant of total drug exposure (31). Steady-state plasma drug concentration achieved for a given drug dose may be higher in elderly patients, which in turn, may increase the risk of toxicity (31). Lipophilic drugs generally have increased distribution in elderly recipients and an increased half-life. Hence drug concentrations may decline at a slower rate in the elderly (as fat acts as a reservoir for the drug) (35), increasing the likelihood of drug accumulation with multiple doses (31).

Age-related pharmacodynamic changes

Drug response is dependent on a range of factors including homeostatic mechanisms, receptor density and affinity, and signal transduction pathways (25,40,41). The entire cascade of events, from drug administration to patient response,

must be considered when examining the pharmacodynamics of a drug (41). Advancing biological age leads to structural and functional changes, including decreased physiological fitness and a marked decline in the robustness of the immune system (immunosenescence) (32). The decline in functional reserve in the elderly is distinctly different between individuals (42).

With aging, there is a progressive reduction in the body's homeostatic mechanisms. This means that in the elderly, it takes more time after administration of a pharmacological intervention (medicine) for counter-regulatory mechanisms to return the body to its original steady-state (32). Additionally, receptor number and responsiveness declines with age, however, this does not lead to a reduction in drug sensitivity or effectiveness. Instead, with advancing age, despite it taking longer for a drug to reach maximum effect (E_{max}), the reaction is significantly stronger, and the overall incidence of adverse effects is greater (32,33). Elderly patients may be more likely to suffer the side effects of medicines when dosed to the same target levels as younger adults.

Immunosenescence

Immunosenescence describes the reduced immune reactivity observed in elderly recipients due to: a decline in naïve T-cells (immune cells), the accumulation of memory T-cells, and alteration of B-cells, causing a decrease in antibody response (15). Immunosenescence can affect all immunological compartments and causes a shift in both the regulation and function of the total immune system (20). The primary mechanism behind this phenomenon is involution of the thymus (shrinkage of the gland responsible for the development of peripheral CD4+ and CD8+ T-cells with age), leading to a progressive reduction in the naïve T-cell count (43). Reduction in naïve T-cell number corresponds to the accumulation of memory T-cells (21,43). Inadequate output of naïve T-cells prevents homeostatic equilibrium of the peripheral T-cell pool being restored, resulting in proliferation of existing peripheral T-cells (homeostatic proliferation). However, this new population of T-cells, comprised mostly of memory T-cells, has a reduced ability to respond to new immunological challenges (21,43).

There is a paucity of evidence surrounding the impact of age on the function of B-cells (44). The literature demonstrates that advancing age affects a decline in naïve B-cells (corresponding to an increase in memory T-cells), as well as diminished B-cell antibody response (15)

[compounded by the reduction in T-cells which are integral to B-cell activation (45)]. Unlike T-cells, B-cell homeostasis is maintained in the periphery through a reduction in the turnover of mature B-cells (20). Despite this, the decline in B-cells causes a change in antibody specificity, and a reduction in plasma cells in the bone marrow (20,43).

Clinically, the effects of immunosenescence in elderly renal transplant patients increases the risk of infections, malignancies, autoimmune disorders, atherosclerosis and neuro-degeneration (20,46). In transplant recipients, immunosenescence likely lowers the risk of acute rejection (47), but increases the risk of drug-related adverse effects, particularly infections and malignancy (48). However, at present, the impact of immunosenescence on the immune system response is only partly understood (49,50). Consequently, it is still unclear how best to dose immunosuppressant drugs in elderly renal transplant recipients.

Immunosuppressant medicines

The most commonly prescribed maintenance immunosuppressive protocol for kidney transplant recipients is triple therapy with a calcineurin inhibitor (cyclosporine or tacrolimus), an antimetabolite (mycophenolate or azathioprine) and a corticosteroid (prednisolone or prednisone) (51,52). This combination of drugs is prescribed concurrently, with each drug primarily targeting a different part of the immune system's response. Tacrolimus, mycophenolate and corticosteroids is the most commonly prescribed combination, regardless of age (53,54). It is thought that intuitively, clinicians may lower immunosuppressant medicine doses in elderly patients due to their reduced immunocompetence and increased susceptibility to side effects and infection, however, the literature has yet to confirm how widespread this practice is, and how therapeutic adjustments are made (55). Currently, the dose of tacrolimus or cyclosporine is adjusted according to drug concentrations measured during TDM, whereas antimetabolites and corticosteroids are prescribed under a fixed dosing regimen in most transplant centres (51).

Calcineurin inhibitors (tacrolimus and cyclosporine)

Calcineurin inhibitors (CNI), cyclosporine and tacrolimus, block calcineurin enzymatic activity. Calcineurin phosphorylates the nuclear factors of activated T lymphocytes (T-cells) which results in the production of interleukin-2 (56).

Interleukin-2 increases the proliferation and activity of T-cells, which are essential for amplification of the immune response. Blocking calcineurin prevents interleukin-2 transcription and T-cell receptor engagement (42,57), suppressing the immune system (42). It has been demonstrated in clinical trials that the rate of acute rejection at one-year post-transplant is significantly greater in patients prescribed cyclosporine compared to tacrolimus ($P=0.001$) (57). Hence, tacrolimus is now the more commonly prescribed CNI, and forms the foundation of 90% patients' initial and maintenance immunosuppressive regimes (7,42).

Blockage of the calcineurin pathway affects non-immune cells, including neurons, skeletal and cardiac myocytes. Off-target effects include enhanced expression of transforming growth factor- β and development of interstitial fibrosis (a feature of nephrotoxicity), and are thought to contribute to the toxicity observed with clinical use of CNIs (42). Both cyclosporine and tacrolimus can cause nephrotoxicity and neurotoxicity (58). However, hypertension and cosmetic complications such as hypertrichosis, gingival hyperplasia and gynaecomastia are more frequently observed in patients taking cyclosporine, whilst glucose intolerance, new-onset diabetes mellitus, alopecia, and diarrhoea impact up to 75% of patients taking tacrolimus (58).

Calcineurin inhibitors have a narrow therapeutic index and display large intra- and inter-individual pharmacokinetic variability (59-61). Over-immunosuppression (too high doses) increases the risk of side effects including neurotoxicity and nephrotoxicity, whilst under-immunosuppression increases the risk of graft rejection (59). Elderly recipients have been shown to be more susceptible to CNI side effects (27), and hence, prescribing equivalent concentrations to younger recipients may increase their risk of toxicity. Currently, specific protocols regarding the best choice and dose of CNIs in elderly recipients have not been validated (1,54).

Tacrolimus is rapidly absorbed in the gastrointestinal tract with maximum drug concentrations generally reached within 1–2 hours. It is highly lipophilic, but poorly water soluble, and passes across the intestinal wall by passive diffusion (29,45). Oral bioavailability is generally poor, with a mean value of approximately 25%, however this can vary substantially between individuals (reported range 5–95%) (42,62). Drug absorption is likely reduced by P-glycoprotein expression in the gastro-intestinal tract, as well as pre-systemic metabolism by cytochrome P450 3A (CYP3A) (62). Tacrolimus binds extensively to erythrocytes (approximately 99%), and in plasma it mainly binds to

alpha-1-acid glycoprotein and albumin (62,63). For this reason, tacrolimus concentrations are typically measured in whole blood rather than plasma (42). Tacrolimus undergoes extensive metabolism via the liver and CYP3A system into more than 15 active metabolites, with >95% eliminated by the biliary route (64) and <1% unchanged drug excreted in the urine and faeces (51). Large variability in the pharmacokinetics of tacrolimus has commonly been attributed to a number of factors including cytochrome P450 genotype, drug-drug interactions, patient haematocrit, patient weight, time post-transplant and patient hepatic function (65).

Cyclosporine is highly lipophilic and only slightly water-soluble. Oral absorption is slow, and overall, cyclosporine has low oral bioavailability (median: 21%) (42), with oral absorption ranging between 10–89% (66). Drug absorption is likely reduced by P-glycoprotein expression in the gastro-intestinal tract, as well as pre-systemic metabolism by CYP3A (60). Similar to tacrolimus, cyclosporine is extensively metabolized by hepatic CYP3A into >25 metabolites and excreted predominantly by the biliary system (90%). Renal excretion accounts for only 6% of drug elimination. Cyclosporine also binds extensively to red blood cells and therefore, whole blood concentrations are typically used to measure drug concentration (42). Large variability in the pharmacokinetics of cyclosporine has been commonly attributed to a number of factors including drug-drug interactions, patient haematocrit, patient weight, time post-transplant and patient hepatic function (66).

Antimetabolites (mycophenolate)

Antimetabolites (mycophenolate and azathioprine) are critical components of maintenance immunosuppressive therapy regimens as they lower the risk of acute rejection by blocking T- and B-cell proliferation (67). Mycophenolate is the typical antimetabolite of choice, as it more effectively lowers the acute rejection risk, when compared to azathioprine (58,68). Mycophenolate is available in two forms: mycophenolate mofetil (MMF; Cellcept[®]) and enteric-coated mycophenolate sodium (Myfortic[®]) which are roughly bio-equivalent (69). Both forms are rapidly converted to the active drug, mycophenolic acid (MPA), in the body. MPA reversibly inhibits the enzyme, inosine 5'-monophosphate dehydrogenase (IMPDH). There are two isoforms of IMPDH, both of which are sensitive to inhibition by MPA (42). IMPDH plays a role in the synthesis of guanine nucleotides, which are critical to T-

and B-lymphocyte proliferation and function (42). Hence, blocking the de-novo purine synthesis pathway of guanine stops the proliferation of T- and B-cells, and the immune system is suppressed (69,70).

Mycophenolate is generally prescribed at a fixed dose regardless of age and has been shown to effectively reduce the incidence of acute rejection and improve long-term graft survival when compared to azathioprine as an adjunct immunosuppressant therapy (69). Side effects are most commonly related to the gastrointestinal tract—mycophenolate frequently causes diarrhoea (30% patients), nausea, bloating, dyspepsia and vomiting (20% patients). More serious side effects include leukopenia, anaemia and thrombocytopenia (58).

Mycophenolate mofetil is rapidly absorbed in the upper gastrointestinal tract and hydrolyzed into MPA and hydroxyethyl morpholine (an inactive metabolite that is rapidly metabolized and then excreted in the urine) (42). Peak concentrations are achieved within 1–2 hours following oral administration. Enteric-coated mycophenolate sodium is more slowly absorbed, with peak drug concentrations occurring within 1.5–2.75 hours after administration (42). Mycophenolate mofetil and enteric-coated mycophenolic sodium formulations both have high oral bioavailability, approximately 80–90%. Drug distribution primarily occurs within the plasma compartment of the blood, with 97–99% of MPA bound to albumin. MPA metabolism is extensive and mostly occurs via the UGT system (71). The mostly inactive metabolite MPAG is then transported from liver cells into the bile. Once biliary excretion has occurred, MPA-7-O-glucuronide (MPAG) may be converted back into MPA by glucosidase produced by the intestinal flora, resulting in MPA reabsorption. This enterohepatic cycle of excretion and reabsorption can lead to a secondary peak within the concentration-time profile (42). Final elimination is by active tubular secretion of MPAG via the kidneys (72). MPA displays non-linear pharmacokinetics, with complex and large pharmacokinetic variability (73) partly attributed to patient serum albumin levels, renal function, genetics, drug-drug interactions (45,71,72,74).

Prednisolone and prednisone

Prednisolone and prednisone are corticosteroids that can interconvert. Prednisolone is the active drug moiety; prednisone is a pro-drug that undergoes reversible metabolism to the active metabolite (75). Corticosteroids are

commonly included in initial immunosuppressant regimens at high doses to lower the risk of acute rejection (76), with doses tapered in the months following transplantation to lower doses (often <10 mg/day) (77). Prednisolone is usually given once daily in the morning to mimic the body's natural diurnal rhythm of cortisol peak and trough concentrations (77).

Prednisolone blocks the immune system by depleting circulating T-cells through the inhibition of interleukin-2 signalling and production. Additionally, it inhibits transcription of the inflammatory cytokine interleukin-1, causing a secondary anti-inflammatory effect, which in turn, detrimentally affects wound healing (77,78). Side effects associated with long-term steroid use are common (54) and can include cosmetic changes (cushingoid appearance), growth retardation, osteoporosis, impaired wound healing, glucose intolerance and hyperlipidaemia (58,79). Although TDM is not necessary (80), it is crucial that doses be adjusted according to an individual's response (managing/minimising side effects).

Studies examining immunosuppressant protocols in elderly recipients

Despite the possibility of physiological and metabolic changes with advancing age, few studies have examined the impact of a lower or altered immunosuppressant dosing protocol on graft outcomes in older adults. In 2009, Badowski and researchers lowered the immunosuppressant targets for a cohort of patients aged over 60 years. In this study, 101 subjects were allocated an initial regimen of mycophenolate mofetil 2 g/day and tacrolimus, target level 10–12 ng/mL; and 88 subjects 1 g/day and 8–10 ng/mL respectively (55). Dose adjustments were made when required, and overall, they found that a reduction in both tacrolimus target drug concentrations and maintenance mycophenolate dosing improved long-term graft survival, without a corresponding increase in the risk of acute rejection ($P=0.006$) (55). Despite these findings, there have been no follow-up studies at different clinics, and it remains largely unknown whether immunosuppressant doses or targets should be adjusted in elderly recipients.

Additionally, there is some contention about the inclusion of corticosteroids in immunosuppressive regimens in the elderly due to long-term toxicity (76). A review by Pascual *et al.* in 2012 showed that steroid withdrawal two weeks after transplantation was not associated with increased mortality or graft loss when patients were prescribed

dual therapy with a CNI and mycophenolate (81). Le Meur *et al.* also concluded in 2015 that steroid avoidance or early withdrawal may improve patient outcomes in individuals with a low immunological risk (the elderly) (15). Resultantly, the US Registry's current guidelines state that a post-transplant steroid-free regimen does not increase the risk of intermediate-term clinical outcomes (81), and a 2012 Annual Report in the USA showed that approximately 30% of all renal transplant recipients are prescribed steroid-free regimens (7). However, a more recent Cochrane review of steroid usage in adult kidney transplant recipients by Haller *et al.* in 2016 reported that steroid avoidance or withdrawal after transplantation increased the risk of acute rejection (82) and concluded that steroids should form part of maintenance immunosuppression (82). To date no clinical trials examining the impact of steroid withdrawal have been specifically done in elderly patients (45).

Studies examining immunosuppressant PK in elderly recipients

Only six studies have been explicitly designed to examine the pharmacokinetics of immunosuppressant medicines in elderly renal transplant recipients. Findings have been summarized in *Table 1* and are critiqued below.

A prospective pilot study in 2008 by Falck *et al.* examined the pharmacokinetics of cyclosporine in 11 elderly and 14 younger adults. Elderly patients (>65 years) achieved similar 2-hour post-dose target levels as younger adults, with lower weight-adjusted doses (4.3 ± 0.8 vs. 6.1 ± 2.1 mg/kg/day; $P=0.025$). This led to the suggestion that drug clearance decreases with advancing age. Additionally, the intracellular concentration of cyclosporine in T-lymphocytes was measured, and it was found that the elderly had higher intracellular cyclosporine concentrations, despite similar whole blood concentrations. The authors suggested that in elderly patients a significantly larger proportion of the whole blood cyclosporine concentration may be located at the site of action (within the T-lymphocyte) indicating that in this population it might be safe to aim for an even lower target level than current guidelines (83).

In 2012, a retrospective, multi-centre study by Jacobson *et al.* examining 374 elderly renal transplant recipients over the first 6 months after transplantation, found that elderly patients (mean: 68.5 years) had higher dose-normalized tacrolimus trough concentrations than younger adults (84). After dose and body weight normalisation, tacrolimus concentrations were greater than 50% higher in elderly

recipients, compared to younger adults (84). The study was retrospective, but involved large patient numbers (1,809 in total across all age groups). One study limitation was that only tacrolimus trough values were examined, rather than total drug exposure across the dosing interval.

A prospective study by David-Neto *et al.* [2017] involving 44 elderly patients and 31 younger adults examined tacrolimus exposure over time. Elderly recipients achieved higher target exposure and lower estimated total body clearance, with a lower normalized tacrolimus dose (85). The authors attribute this finding to a lower apparent clearance of tacrolimus in elderly patients. In this study, the average age of elderly patients was 65 years (85).

A prospective, point prevalence study conducted by Miura *et al.* in 2009 was the first to examine multiple immunosuppressant medicines in elderly patients, and showed that age had no impact on dose and weight adjusted pharmacokinetic parameters of tacrolimus, prednisolone and mycophenolate (C_{max} , C_0 and AUC_{0-12h}) (86). However, only 12 of the 110 patients studied were elderly, which was defined as being 60–64 years of age (86). The study was undertaken at a single time-point, four weeks post-transplantation, hence, intra-patient variability and changes over time-post-transplantation were not considered.

Wang *et al.* [2007] were the first to examine the impact of aging on the pharmacokinetics of mycophenolate mofetil. MPA exposure, based on the area under the concentration-time curve (AUC), was significantly lower in elderly patients despite receiving the same MMF doses. There was however, no significant difference in peak drug concentration between the elderly and younger adult patient groups (87). This finding suggests that elderly patients have higher apparent clearance of mycophenolate, which contradicts our general understanding of the effects of aging on the pharmacokinetics of a drug. The authors instead suggest that elderly patients have more complex drug absorption processes (87). Overall, the sample size was relatively small, with 24 elderly and 24 younger adult patients examined. The mean age of the elderly patients was 65 years. As the study was a point-prevalence study, with each patients sampled between 10–12 weeks after transplantation, intra-patient variability could not be accounted for, which may explain the unusual finding.

A final study by Tang *et al.* in 2016, compared the effects of aging on the pharmacokinetics of mycophenolate in younger adults (54 patients) and the elderly (26 patients) (70). Patients were repetitively sampled over the first six months post-transplantation, with all patients given the same dose

Table 1 A summary of studies examining the pharmacokinetic parameters of immunosuppressant medicines in elderly renal transplant patients

Study origin/design	Immuno-suppressant studied	Sample size	Definition of "Elderly"; population studied	PK approach	Results/conclusions
Falck <i>et al.</i> Norway; prospective pilot study (83)	Cyclosporine	18–64 years: n=14; >65 years: n=11	>65 years; mean: 73±3	Whole blood cyclosporine AUC _{0-12h} stable post-Tx (30–40 days) ^A and intracellular concentration in T-lymphocytes AUC _{0-6h} ^A	Elderly received lower doses to achieve same cyclosporine whole blood 2-hour post-dose target as younger patients (4.3±0.8 vs. 6.1±2.1 mg/day/kg; P=0.025), and had lower whole blood clearance (22.5±5.2 vs. 30.2±10.4 L/h; P=0.032) and low intracellular cyclosporine clearance (0.951×10 ⁶ ±0.32×10 ⁶ vs. 1.72×10 ⁶ ±0.71×10 ⁶ cells/hr; P=0.0029)
Jacobson <i>et al.</i> USA; longitudinal study (84)	Tacrolimus	18–34 years: n=348; 35–64 years: n=1,831; 65–84 years: n=374	>65 years; median: 68.5	Whole blood tacrolimus trough level (bi-weekly weeks 1–8 and bi-monthly months 3–6) ^B	Elderly had higher tacrolimus troughs than younger adults (129.8 vs. 77.1 ng/mL·mg/kg; D/BW ^D -adjustment) and received lower doses by 1–2 mg/day; age and CYP3A5*1 genotype effects tacrolimus troughs
David-Neto <i>et al.</i> Brazil; longitudinal study (85)	Tacrolimus	<60 years: n=31; >60 years: n=44	>60 years; mean: 65±3	Whole blood tacrolimus AUC _{0-12h} (7, 30, 60, and 180 days post-Tx) ^{B,E}	Elderly had lower weight-adjusted tacrolimus doses and higher tacrolimus exposure (AUC _{adjusted} day 7: 2,286±1,372 vs. 1,369±582 ng·hr·kg/mL; P=0.001); Elderly require a lower tacrolimus dose, compared to younger adults (day 7: 79±27 vs. 119±35 µg/kg/day; P<0.01)
Miura <i>et al.</i> Japan; point prevalence study (86)	Tacrolimus, MPA and prednisolone	20–39 years: n=41; 40–59 years: n=57; 60–64 years: n=12	>60 years; mean: 63±3	Whole blood tacrolimus ^C AUC _{0-12h} , plasma MPA AUC _{0-12h} , plasma prednisolone AUC _{0-24h} , plasma MPA AUC _{0-12h} (estimate of enterohepatic recirculation) ^A	Age had no impact on D/BW ^D adjusted PK of tacrolimus or dose-adjusted parameters of prednisolone and MPA
Wang <i>et al.</i> China; point prevalence study (87)	MPA	18–55 years: n=24; >60 years: n=24	>60 years; mean: 66±4	Plasma MPA AUC _{0-12h} ^{A,B}	Mean MPA AUC _{0-12h} significantly lower in elderly patients (elderly: 22.2±9.0 µg·h/mL vs. adults: 32.8±8.8 µg·h/mL; P=0.016); no difference in: pre-dose, peak concentrations, or peak times, between groups
Tang <i>et al.</i> The Netherlands; longitudinal study (70)	MPA	19–58 years: n=54; 60–76 years: n=26	>65 years; mean: 66±5	Plasma MPA AUC _{0-12h} (day 6); Week 3, 7 and 20—limited sampling MPA AUCs (3 samples) ^{A,B}	Age did not significantly affect the PK of MPA; no age-related adjustments to dosing should be made

^A, drug levels measured by a validated high-performance liquid chromatography method; ^B, AUC calculated using linear trapezoidal method; ^C, whole blood concentrations determined by microparticle Enzyme Immunoassay (MEIA) IMx[®] method; ^D, D/BW = dose/body weight; ^E, only 22 elderly patients performed the analysis at 60 and 180 days post-transplant. AUC, area-under the concentration-time curve; CYP3A5, cytochrome P450; MPA, mycophenolic acid; PK, pharmacokinetics; Tx, transplant.

of mycophenolate mofetil. Researchers concluded that aging does not significantly alter mycophenolate exposure (AUC_{0-12h}), which differs to the findings by Wang *et al.* (87) and Miura *et al.* (86). Authors suggested this may be due to the pharmacokinetics of mycophenolate being investigated repetitively both early and later post-transplantation. The mean age of the elderly recipients was 65 years (±4.9 years). Due to the necessary addition of tacrolimus as co-immunosuppressant therapy, the effects of ageing on tacrolimus were also examined. Tang *et al.*'s tacrolimus results mirrored David-Neto *et al.*'s (85) with both

studies reporting that age significantly affects tacrolimus pharmacokinetics, with elderly patients having higher weight-normalized tacrolimus concentrations compared to younger adults (70).

Overall, these studies suggest that the doses of cyclosporine and tacrolimus required to achieve optimum immunosuppression in elderly patients may need to be lower than doses in younger adults, whilst mycophenolate doses should remain unchanged. No studies to date have looked at the effects of aging on MPA metabolites or free drug levels. Studies investigating the effects of

Table 2 A summary of studies examining the pharmacodynamics effects of immunosuppressant medicines in elderly renal transplant recipients

Study origin/design	Immuno-suppressant studied	Sample size	Inclusion criteria	Definition of "Elderly"; population studied	PD approach	Results/conclusions
Tang <i>et al.</i> The Netherlands; longitudinal study (70)	MPA	19–58 years: n=54; 60–76 years: n=26	Uncomplicated recovery to a kidney transplant, regimen included MPA	>65 years; mean: 66±5	IMPDH activity—baseline, day 6, weeks 3, 7 and 20 post-transplant, measured using area under the effect (IMPDH activity)—time curve (AEC _{0–12h}) ^A	Baseline IMPDH activity showed large inter-individual variability. No significant differences in IMPDH activity at baseline or any time point post-transplant between age groups

^A, calculated using a linear trapezoidal model. IMPDH, inosine 5'-monophosphate dehydrogenase; MPA, mycophenolic acid.

aging on the pharmacokinetics of corticosteroids are also lacking. Unfortunately, few studies have included many patients over 70 years of age and hence it is still largely unknown how the physiological process of ageing affects the pharmacokinetic parameters of immunosuppressant medicines in this cohort (27).

Additionally, a number of population pharmacokinetic studies have been performed examining the pharmacokinetics of cyclosporine, tacrolimus and MPA in adult kidney transplant recipients [see Han 2013, Staatz 2011, Brooks 2016, and Kiang and Ensom, for reviews of these studies (65,66,71,74)]. These studies have not specifically examined the effect of being 'elderly' on the pharmacokinetics of immunosuppressant medicines, but rather looked at patient age in general, and as such, did not necessarily include a large number of elderly patients. Consequently, only some of these studies have identified patient age as an important covariate (88–92). As these studies generally look at multiple covariate factors, some of the effects of aging may also be explained by other patient factors, which can change with age (e.g., patient weight, renal function, haematocrit).

Studies examining immunosuppressant PD in elderly recipients

Studies examining immunosuppressant pharmacodynamics in elderly renal transplant recipients are lacking (Table 2). One study by Tang *et al.* [2016] examined baseline inosine 5'-monophosphate dehydrogenase (IMPDH) activity and MPA-induced IMPDH activity in 54 younger adults (mean age: 44 years) and 26 elderly (mean age: 65 years) renal transplant recipients. Patients were repetitively sampled for the first six months after transplantation. Overall, researchers found no difference in IMPDH activity pre-transplantation between the two age groups and there

was no significant difference in AEC_{0–12h} [area under the effect-time curve), EC₅₀ (concentration that yields half the maximum response (31)] or E_{max} (70). This led researchers to conclude that age does not affect the pharmacodynamics of MPA (70). Clearly, further studies are required.

Transplant outcomes

Elderly renal transplant patients are at greatest risk of both morbidity and mortality (Table 3) (100). Elderly recipients are more susceptible to drug-related adverse effects, particular cardiovascular disease and infection (27,101), as well as more likely to suffer increased co-morbidity, both of which have been shown to increase post-transplant complications (27) and negatively impact patient survival (47,96).

Elderly recipients are more likely to die from infectious or cardiovascular-related causes due to side effects of immunosuppressant medicines, rather than graft failure (27,32,100). A study by Ojo *et al.* showed that renal transplant recipients aged over 65 years were seven times more likely to die with a functioning graft compared with young adults (aged 18–29 years) (46,100,102). This suggests that an optimal balance between immunosuppressant medicine efficacy and toxicity is not achieved in elderly recipients and further studies are needed to foster long-term graft and patient survival.

The risk of infection-related mortality has been shown to increase exponentially with age, due to reduced immune reactivity (29,103,104). Meier-Kriesche *et al.* reported in 2000 that elderly renal transplant recipients have a five-fold increase in their overall risk of death due to infection, compared with recipients aged between 30–39 years (105). Risk of infection, both opportunistic and non-opportunistic, is also increased with pharmacological over-immunosuppression, as infection is a side effect of immunosuppressant medicines (103). The literature suggests that lower maintenance immunosuppressant

Table 3 A summary of studies examining transplant outcomes in elderly versus younger adult renal transplant recipients

Study origin/design	Sample size	Definition of "Elderly"; population studied	Transplant outcomes	Results/conclusions
Schulak <i>et al.</i> USA; prospective study (93)	TOTAL: n=29. Low risk patients—no diabetes or cardiovascular disease	≥60 years; mean: 62±2	Overall 1 year patient survival in the elderly—82% Graft survival in the elderly* at: 1 year 84% low risk patients vs. 58% high risk patients Patient survival in the elderly* at: 1 year 91% low risk patients vs. 67% high risk patients	High risk patient survival was worse (P=0.055); therefore, comorbidity pre-transplant screening is important. Patient death was the major cause of graft loss in elderly recipients
Doyle <i>et al.</i> USA; retrospective study (94)	TOTAL: n=1,846: ≥60 years: n=206; 18–59 years: n=1,640	≥60 years; mean: 64±3	Graft survival* at (P<0.0001): 1 year 83% younger adults vs. 73% elderly; 5 years 70% younger adults vs. 56% elderly Patient survival* at (P<0.0001): 1 year 96% younger adults vs. 85% elderly; 5 years 86% younger adults vs. 69% elderly Graft loss due to death—45% younger adults vs. 61% elderly (P>0.05)	Patient (P<0.001) and graft (P<0.001) survival decreased in elderly compared to younger adults. Pre-transplant history of malignancy, vascular disease and smoking history increased risk of graft loss and patient death in elderly (P<0.001)
Otero-Ravina <i>et al.</i> Spain; retrospective study (95)	TOTAL: n=621: <60 years: n=137; >60 years: n=484	> 60 years; mean: 64±3	Graft survival* at: (P=0.0012): 1 year 82% younger adults vs. 73% elderly, and; 5 years 70% younger adults vs. 56% elderly Patient survival * at: (P<0.0001): 1 years 96% younger adults vs. 85% elderly; 5 years 86% younger adults vs. 69 elderly Graft loss in elderly—47% due to patient death	Mortality and graft survival greater in younger adults compared to elderly recipients (P<0.05)
Huang <i>et al.</i> USA; retrospective study (96)	Total: n=31,179: 60–69 years: n=24,877; 70–79 years: n=6,103; 80+: n=199	>60 years; median: 64	Graft survival* at (P<0.0001): 2 years: 60–69 (89%), 70–79 (86%), 80+ (73%) Graft loss: greater in 80 years than 60–69 years (HR 1.78, 95% CI: 1.42–2.23)	Increased mortality risk in >80 years group compared with 60–69 year group (P<0.0001)
Boesmueller <i>et al.</i> Austria; retrospective study (46)	Total: n=83: <70 years: n=64; ≥70 years: n=19	>70 years; mean: 72.7	Graft survival* at (P>0.05): 1 year 94.4% younger adults vs. 95% elderly; 5 years 79% younger adults vs. 52% elderly Patient survival* at (P=0.028): 1 year 94.4% younger adults vs. 95% elderly; 5 years 82.6% younger adults vs. 67% elderly	Major cause of graft loss in both groups was death with a functioning graft Patient survival reduced in elderly, compared with younger adults (P=0.028). Morbidity in elderly related to hemodynamic, oncologic and infectious events
Impedovo <i>et al.</i> USA; retrospective study (97)	Total: n=761: >60 years: n=69; <60 years: n=692	>60 years; mean: 60±11	Graft survival at: 1 year 97% younger adults vs. 91.9% elderly; 3 years 95.4% younger adults vs. 79.0% elderly; 5 years 94.2% younger adults vs. 79% elderly No difference in death censored graft survival at 1, 3, and 5 years (P=0.08) between groups	Elderly had worse survival compared to younger adults (P=0.0005). Common causes of death in the elderly were myocardial infarction, other cardiovascular complications and tumours
Hatamizadeh <i>et al.</i> USA; retrospective study (98)	Total: n=15,667: 65–69 years: n=10,101; 70–74 years: n=4,271; 75+: n=1,295. Reference group, 15–64 years: n=129,803	>65 years; mean: 67±1	Mortality risk increase: 1.8 folds 65–74 group vs. 2.0 folds elderly group (reference was baseline comparison) Patient survival—poor predictors: obesity (BMI >30 kg/m ²) and diabetes	Transplantation attenuates mortality risk, and comorbidities are not associated with poorer outcomes in elderly
Karim <i>et al.</i> UK; retrospective study (13)	Total: n=19,103: <50 years: n=11,421; 50–59 years: n=4,195; 60–69 years: n=2,887; 70–79 years: n=589; 80+ years: n=11	>70 years	Mortality risk increased with age*: <50 (5.8%), 50–59 (14.2%), 60–69 (22.0%), 70–79 (31.9%), and 80+ (45.5%) Most common causes of death in elderly were cardiac (21.2%), infection (21.2%) and malignancy (20.2%)	Overall risk of death increased with each additional decade >50 years (P<0.001)
Wu <i>et al.</i> USA; cohort study (99)	TOTAL: n=1,102: 60–86 years: n=266	>65 years	Most common co-morbidities—diabetes 30.3%, heart failure 11.9% CCI score ≥5 more likely in older patients Patient death significantly higher with CCI score ≥5 (HR 2.67; 95% CI: 1.75–4.08; P<0.001)	Increased comorbidity number (≥5) negatively affects patient survival (P<0.001).

*, calculated by Kaplan-Meier methods. BMI, body mass index; CCI, Charlson Comorbidity Index; CI, confidence interval; HR, hazard ratio.

targets in elderly recipients may decrease patient susceptibility to side effects, which in turn, would lower the risk of infection (14), however, studies in patients aged above 70 years have yet to establish this (19,100,106).

Co-morbidities increase with advancing age, irrespective of transplantation. A study by Wu *et al.* [2005] examined the impact of patient co-morbidities on acute rejection and patient and graft survival in a cohort of 1,102 adult renal transplant patients (Table 3) (99). The study did not specifically examine elderly patients, however, 266 patients aged greater than 60 years (median age: 67 years) were included (99). Wu and researchers showed a trend toward an increasing number of co-morbidities over time, with significantly more patients having greater than five co-morbidities at the study's conclusion ($P < 0.05$). The results also showed that patients with greater than five co-morbidities were at significantly greater risk of death both immediately and more than three months post-transplantation, compared to patients with less than five co-morbidities ($P < 0.05$) (99). With respect to age, researchers showed that elderly patients had the lowest survival rate, with survival decreasing with increasing age ($P < 0.05$). Overall, this study serves to show the detrimental effect of co-morbidity number on both acute and long-term patient and graft outcomes (99).

Conclusions

As the number of elderly patients undergoing renal transplantation continues to rise rapidly (107), it is crucial that current immunosuppressant prescribing practices be more clearly examined and defined. It is imperative that the effects of aging on the pharmacokinetics and pharmacodynamics of immunosuppressant medicines be established so that the most safe and efficacious immunosuppressive regimens are prescribed.

From the limited studies done to date, it appears that elderly transplant recipients have higher tacrolimus exposure (up to 50% greater), despite receiving lower adjusted doses, whilst mycophenolate exposure remains unchanged. These preliminary findings support the assumption that CYP associated metabolism is more likely to be affected by aging than metabolism by UGTs (108). Future studies should examine MPAG exposure in the elderly, as this is likely to increase with reduced patient

renal function. MPAG competes with MPA for protein binding sites, and a reduction in renal function may increase MPAG levels in the elderly. Resultantly, higher MPAG levels could potentially be influencing mycophenolate dosing, if adjustments were made according to total drug exposure. Further studies are also required to ascertain the effect of ageing on intracellular drug levels in lymphocytes.

Based on the results of a single study, there appears to be no difference in the effect of mycophenolate on IMPDH activity between elderly and younger adult recipients, however, further studies are required to corroborate and confirm this finding. Future studies should also examine CNIs, in terms of their maximum effect (E_{max}) on calcineurin inhibition, as currently, elderly recipients are most likely to suffer CNI side effects, but it is not completely understood as to why. What is clear is that the elderly are at greatest risk of immunosuppressant side effects, with the three main causes of death being infection, malignancy and cardiovascular disease attributed to immunosuppressant medicine usage (27,45,100,101). This suggests that the pharmacodynamic effects of immunosuppressant medicines are influenced by age. It is difficult to know with certainty whether a difference in baseline reserve or difference in drug sensitivity (31) (or both) is responsible for the change in patient responsiveness to immunosuppressant medicines in elderly renal transplant recipients.

Like any transplant group, the elderly are a heterogeneous sub-population and this influences patient responsiveness to drugs. Because of this unpredictability, elderly renal transplant recipients may not only require less immunosuppression but also specific, individualised changes, as immunosenescence affects some immunological targets more so than others (45). The complexity surrounding appropriate immunosuppressant dosing is compounded by the usage of combination immunosuppressant drug treatments. Currently, it is unknown whether elderly recipients justly require the same level of immunosuppression to support graft outcomes as younger adults, nor which immunosuppressant agents or target concentrations best achieve optimal patient and graft outcomes (106,109). Further studies are required to characterise these age-related changes (110).

Acknowledgments

None.

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

References

- Hod T, Goldfarb-Rumyantzev AS. Clinical issues in renal transplantation in the elderly. *Clin Transplant* 2015;29:167-75.
- Oniscu GC, Brown H, Forsythe JL. How great is the survival advantage of transplantation over dialysis in elderly patients? *Nephrol Dial Transplant* 2004;19:945-51.
- Wolfe RA, Ashby VB, Milford EL, et al. Comparison of mortality in all patients on dialysis, patients on dialysis awaiting transplantation, and recipients of a first cadaveric transplant. *N Engl J Med* 1999;341:1725-30.
- Cooper M, Forland CL. The elderly as recipients of living donor kidneys, how old is too old? *Curr Opin Organ Transplant* 2011;16:250-5.
- Huang E, Segev DL, Rabb H. Kidney transplantation in the elderly. *Semin Nephrol* 2009;29:621-35.
- Segall L, Nistor I, Pascual J, et al. Criteria for and Appropriateness of Renal Transplantation in Elderly Patients With End-Stage Renal Disease: A Literature Review and Position Statement on Behalf of the European Renal Association-European Dialysis and Transplant Association Descartes Working Group and European Renal Best Practice. *Transplantation* 2016;100:e55-65.
- U.S. Department of Health and Human Services. OPTN/SRTR 2012 Annual Data Report: Kidney 2012 Annual Data Report. USA, 2012:11-44.
- ANZDATA Registry. The 39th Annual ANZDATA Report Chapter 8: Transplantation 2016.
- Pippias M, Stel VS, Abad Diez JM, et al. Renal replacement therapy in Europe: a summary of the 2012 ERA-EDTA Registry Annual Report. *Clin Kidney J* 2015;8:248-61.
- Maggiore U, Abramowicz D, Budde K. Renal transplantation in the elderly. *Transplant Rev (Orlando)* 2015;29:191-2.
- Morris P, Knechtle SJ. Results of Renal Transplantation. In: Patzer RE, Knechtle SJ. editors. *Kidney Transplantation - Principles and Practice*. UK: Elsevier, 2014:676-97.
- Aubert O, Kamar N, Vernerey D, et al. Long term outcomes of transplantation using kidneys from expanded criteria donors: prospective, population based cohort study. *BMJ* 2015;351:h3557.
- Karim A, Farrugia D, Cheshire J, et al. Recipient age and risk for mortality after kidney transplantation in England. *Transplantation* 2014;97:832-8.
- Macrae J, Friedman AL, Friedman EA, et al. Live and deceased donor kidney transplantation in patients aged 75 years and older in the United States. *Int Urol Nephrol* 2005;37:641-8.
- Le Meur Y. What immunosuppression should be used for old-to-old recipients? *Transplant Rev (Orlando)* 2015;29:231-6.
- Ponticelli C, Podesta MA, Graziani G. Renal transplantation in elderly patients. How to select the candidates to the waiting list? *Transplant Rev (Orlando)* 2014;28:188-92.
- Rao PS, Merion RM, Ashby VB, et al. Renal transplantation in elderly patients older than 70 years of age: results from the Scientific Registry of Transplant Recipients. *Transplantation* 2007;83:1069-74.
- Frei U, Noeldeke J, Machold-Fabrizii V, et al. Prospective age-matching in elderly kidney transplant recipients--a 5-year analysis of the Eurotransplant Senior Program. *Am J Transplant* 2008;8:50-7.
- Sutherland AI, IJzermans JN, Forsythe JL, et al. Kidney and liver transplantation in the elderly. *Br J Surg* 2016;103:e62-72.
- Heinbokel T, Elkhali A, Liu G, et al. Immunosenescence and organ transplantation. *Transplant Rev (Orlando)* 2013;27:65-75.
- Klinger M, Banasik M. Immunological characteristics of the elderly allograft recipient. *Transplant Rev (Orlando)* 2015;29:219-23.
- Abecassis M, Bridges ND, Clancy CJ, et al. Solid-organ transplantation in older adults: current status and future research. *Am J Transplant* 2012;12:2608-22.
- McLachlan AJ, Hilmer SN, Le Couteur DG. Variability in response to medicines in older people: phenotypic and genotypic factors. *Clin Pharmacol Ther* 2009;85:431-3.
- de Oliveira Baldoni A, Drumond Chequer FM, Ferraz ERA, et al. Elderly and drugs: risks and necessity of rational use. *Brazilian Journal of Pharmaceutical Sciences* 2010;46:617-32.
- Hutchison LC, O'Brien CE. Changes in Pharmacokinetics and Pharmacodynamics in the Elderly Patient. *Journal of Pharmacy Practice* 2016;20:4-12.
- Pascual J, Zamora J, Pirsch JD. A systematic review of kidney transplantation from expanded criteria donors. *Am J Kidney Dis* 2008;52:553-86.

27. Shi YY, Hesselink DA, van Gelder T. Pharmacokinetics and pharmacodynamics of immunosuppressive drugs in elderly kidney transplant recipients. *Transplant Rev (Orlando)* 2015;29:224-30.
28. Johnson DW, Herzig K, Purdie D, et al. A comparison of the effects of dialysis and renal transplantation on the survival of older uremic patients. *Transplantation* 2000;69:794-9.
29. Meier-Kriesche HU, Ojo AO, Hanson JA, et al. Exponentially increased risk of infectious death in older renal transplant recipients. *Kidney Int* 2001;59:1539-43.
30. Lim WH, Dogra G, Chadban SJ, et al. Lack of impact of donor age on patient survival for renal transplant recipients ≥ 60 years. *Transpl Int* 2012;25:401-8.
31. Jacobson S. Effects of Pharmacokinetic and Pharmacodynamic Changes in the Elderly. *Psychiatry Times* 2013;26-9.
32. Turnheim K. When drug therapy gets old: pharmacokinetics and pharmacodynamics in the elderly. *Experimental Gerontology* 2003;38:843-53.
33. Aymanns C, Keller F, Maus S, et al. Review on pharmacokinetics and pharmacodynamics and the aging kidney. *Clin J Am Soc Nephrol* 2010;5:314-27.
34. Colloca G, Santoro M, Gambassi G. Age-related physiologic changes and perioperative management of elderly patients. *Surg Oncol* 2010;19:124-30.
35. Birkett DJ. *Pharmacokinetics Made Easy* 2nd ed. pocket guide ed. Australia: McGraw Hill Education, 2010.
36. Petzinger E, Geyer J. Drug transporters in pharmacokinetics. *Naunyn Schmiedebergs Arch Pharmacol* 2006;372:465-75.
37. Stieger B, Hagenbuch B. Recent advances in understanding hepatic drug transport. *F1000 Research* 2016;5:2465.
38. van Assema DM, Lubberink M, Boellaard R, et al. P-glycoprotein function at the blood-brain barrier: effects of age and gender. *Mol Imaging Biol* 2012;14:771-6.
39. McLean AJ, Le Couteur DG. Aging biology and geriatric clinical pharmacology. *Pharmacol Rev* 2004;56:163-84.
40. Corsonello A, Pedone C, Antonelli Incalzi R. Age-Related Pharmacokinetic and Pharmacodynamic Related Risk of Adverse Drug Reactions. *Current Medicinal Chemistry* 2010;17:571-84.
41. Hämmerlein A, Derendorf H, Lowenthal DT. Pharmacokinetic and Pharmacodynamic Changes in the Elderly: Clinical Implications. *Clin Pharmacokin* 1998;35:49-64.
42. Milone MC. Chapter 1 - Overview of the pharmacology and toxicology of immunosuppressant agents that require therapeutic drug monitoring. *Personalized Immunosuppression in Transplantation* 2016.
43. Albright JF, Albright JW. *Aging of Adaptive/Acquired Immunity. Aging, Immunity, and Infection*. Totowa, NJ: Humana Press, 2003:135-211.
44. Gibson KL, Wu YC, Barnett Y, et al. B-cell diversity decreases in old age and is correlated with poor health status. *Aging Cell* 2009;8:18-25.
45. Krenzien F, ElKhal A, Quante M, et al. A Rationale for Age-Adapted Immunosuppression in Organ Transplantation. *Transplantation* 2015;99:2258-68.
46. Boesmueller C, Biebl M, Scheidl S, et al. Long-term outcome in kidney transplant recipients over 70 years in the Eurotransplant Senior Kidney Transplant Program: a single center experience. *Transplantation* 2011;92:210-6.
47. Peeters LEJ, Andrews LM, Hesselink DA, et al. Personalized immunosuppression in elderly renal transplant recipients. *Pharmacol Res* 2018;130:303-7.
48. Kuypers DRJ. *Immunotherapy in Elderly Transplant Recipients: A Guide to Clinically Significant Drug Interactions*. *Drugs Aging* 2009;26:715-37.
49. Seyda M, Quante M, Uehara H, et al. Immunosenescence in renal transplantation: a changing balance of innate and adaptive immunity. *Curr Opin Organ Transplant* 2015;20:417-23.
50. Halleck F, Khadzhyrov D, Liefeldt L, et al. Immunologic outcome in elderly kidney transplant recipients: is it time for HLA-DR matching? *Nephrol Dial Transplant* 2016;31:2143-9.
51. Barraclough KA, Staatz CE, Johnson DW, et al. Kidney transplant outcomes are related to tacrolimus, mycophenolic acid and prednisolone exposure in the first week. *Transpl Int* 2012;25:1182-93.
52. Chadban SJ, Barraclough KA, Campbell SB, et al. KHA-CARI guideline: KHA-CARI adaptation of the KDIGO Clinical Practice Guideline for the Care of Kidney Transplant Recipients. *Nephrology (Carlton)* 2012;17:204-14.
53. *Renal transplantation and the older adult patient* [Internet]. Wolters Kluwer Health 2016 [cited 30/11/2017]. Available online: <https://www.uptodate.com/contents/renal-transplantation-and-the-older-adult-patient#H1>
54. Chadban SJ, Barraclough KA, Campbell SB, et al. The Kidney Health Australia-Caring for Australasians with Renal Impairment (KHA-CARI) Guidelines. *Nephrology* 2012;17:204-14.
55. Badowski M, Gurk-Turner C, Cangro C, et al. The impact of reduced immunosuppression on graft outcomes

- in elderly renal transplant recipients. *Clin Transplant* 2009;23:930-7.
56. Karpe KM, Talaulikar GS, Walters GD. Calcineurin inhibitor withdrawal or tapering for kidney transplant recipients. *Cochrane Database Syst Rev* 2017;7:CD006750.
 57. Morris P, Knechtle SJ. Calcineurin Inhibitors. In: Mejia JC, Basu A, Shapiro R. editors. *Kidney Transplantation - Principles and Practice*. UK: Elsevier, 2014:231-49.
 58. Muntean A, Lucan M. Immunosuppression in Kidney Transplantation. *Clujul Medical* 2013;86:177-80.
 59. Kim IW, Moon YJ, Ji E, et al. Clinical and genetic factors affecting tacrolimus trough levels and drug-related outcomes in Korean kidney transplant recipients. *Eur J Clin Pharmacol* 2012;68:657-69.
 60. Okada A, Ushigome H, Kanamori M, et al. Population pharmacokinetics of cyclosporine A in Japanese renal transplant patients: comprehensive analysis in a single center. *Eur J Clin Pharmacol* 2017;73:1111-9.
 61. Lloberas N, Elens L, Llaudo I, et al. The combination of CYP3A4*22 and CYP3A5*3 single-nucleotide polymorphisms determines tacrolimus dose requirement after kidney transplantation. *Pharmacogenet Genomics* 2017;27:313-22.
 62. Andreu F, Colom H, Elens L, et al. A New CYP3A5*3 and CYP3A4*22 Cluster Influencing Tacrolimus Target Concentrations: A Population Approach. *Clin Pharmacokinet* 2017;56:963-75.
 63. Staatz CE, Tett SE. Pharmacokinetic Considerations Relating to Tacrolimus Dosing in the Elderly. *Drugs Aging* 2005;22:541-57.
 64. Staatz CE, Willis C, Taylor PJ, et al. Population pharmacokinetics of tacrolimus in adult kidney transplant recipients. *Clin Pharmacol Ther* 2002;72:660-9.
 65. Brooks E, Tett SE, Isbel NM, et al. Population Pharmacokinetic Modelling and Bayesian Estimation of Tacrolimus Exposure: Is this Clinically Useful for Dosage Prediction Yet? *Clin Pharmacokinet* 2016;55:1295-335.
 66. Han K, Pillai VC, Venkataramanan R. Population pharmacokinetics of cyclosporine in transplant recipients. *AAPS J* 2013;15:901-12.
 67. McKay DB, Steinberg SM. *Kidney Transplantation: A Guide to the Care of Kidney Transplant Recipients*. New York: Springer, 2010.
 68. Johnson DW, Nicol DL, Purdie DM, et al. Is Mycophenolate Mofetil less safe than Azathioprine in Elderly Renal Transplant Recipients? *Transplantation* 2002;73:1158-63.
 69. Salvadori M, Holzer H, Mattos AD, et al. Enteric-Coated Mycophenolate Sodium is Therapeutically Equivalent to Mycophenolate Mofetil in de novo Renal Transplant Patients. *American Journal of Transplantation* 2004;4:231-6.
 70. Tang JT, de Winter BC, Hesselink DA, et al. The pharmacokinetics and pharmacodynamics of mycophenolate mofetil in younger and elderly renal transplant recipients. *Br J Clin Pharmacol* 2017;83:812-22.
 71. Kiang TKL, Ensom MHH. Population Pharmacokinetics of Mycophenolic Acid: An Update. *Clin Pharmacokinet* 2018;57:547-58.
 72. Božina N, Lalic Z, Nad-Skegro S, et al. Steady-state pharmacokinetics of mycophenolic acid in renal transplant patients: exploratory analysis of the effects of cyclosporine, recipients' and donors' ABCC2 gene variants, and their interactions. *Eur J Clin Pharmacol* 2017;73:1129-40.
 73. Staatz CE, Duffull SB, Kiberd B, et al. Population pharmacokinetics of mycophenolic acid during the first week after renal transplantation. *Eur J Clin Pharmacol* 2005;61:507-16.
 74. Staatz CE, Tett SE. Maximum A Posteriori Bayesian Estimation of Mycophenolic Acid Area Under the Concentration-Time Curve: Is This Clinically Useful for Dosage Prediction Yet? *Clin Pharmacokinet* 2011;50:759-72.
 75. Bergmann TK, Barraclough KA, Lee KJ, et al. Clinical pharmacokinetics and pharmacodynamics of prednisolone and prednisone in solid organ transplantation. *Clin Pharmacokinet* 2012;51:711-41.
 76. Haller MC, Kammer M, Kainz A, et al. Steroid withdrawal after renal transplantation: a retrospective cohort study. *BMC Med* 2017;15:8.
 77. Morris P, Knechtle SJ. Steroids. In: Knight SR, editor. *Kidney Transplantation – Principles and Practice*. 7 ed. UK: Elsevier, 2014:221-30.
 78. Glucocorticoid effects on the immune system [Internet]. Wolters Kluwer Health 2017. Available online: <https://www.uptodate.com/contents/glucocorticoid-effects-on-the-immune-system>
 79. Morton JM, Williamson S, Kear LM, et al. Therapeutic drug monitoring of prednisolone after lung transplantation. *J Heart Lung Transplant* 2006;25:557-63.
 80. Potter JM, McWhinney BC, Sampson L, et al. Area-Under-the-Curve Monitoring of Prednisolone for Dose Optimization in a Stable Renal Transplant Population. *Ther Drug Monit* 2004;26:408-14.

81. Pascual J, Royuela A, Galeano C, et al. Very early steroid withdrawal or complete avoidance for kidney transplant recipients: a systematic review. *Nephrol Dial Transplant* 2012;27:825-32.
82. Haller MC, Royuela A, Nagler EV, et al. Steroid avoidance or withdrawal for kidney transplant recipients. *Cochrane Database Syst Rev* 2016;(8):CD005632.
83. Falck P, Asberg A, Byberg KT, et al. Reduced elimination of cyclosporine A in elderly (>65 years) kidney transplant recipients. *Transplantation* 2008;86:1379-83.
84. Jacobson PA, Schladt D, Oetting WS, et al. Lower calcineurin inhibitor doses in older compared to younger kidney transplant recipients yield similar troughs. *Am J Transplant* 2012;12:3326-36.
85. David-Neto E, Romano P, Kamada Triboni AH, et al. Longitudinal Pharmacokinetics of Tacrolimus in Elderly Compared With Younger Recipients in the First 6 Months After Renal Transplantation. *Transplantation* 2017;101:1365-72.
86. Miura M, Satoh S, Kagaya H, et al. No impact of age on dose-adjusted pharmacokinetics of tacrolimus, mycophenolic acid and prednisolone 1 month after renal transplantation. *Eur J Clin Pharmacol* 2009;65:1047-53.
87. Wang CX, Meng FH, Chen LZ, et al. Population pharmacokinetics of mycophenolic acid in senile Chinese kidney transplant recipients. *Transplant Proc* 2007;39:1392-5.
88. Falck P, Midtvedt K, Vân Lê TT, et al. A Population Pharmacokinetic Model of Cyclosporin Applicable for Assisting Dose Management of Kidney Transplant Recipients. *Clin Pharmacokinet* 2009;48:615-23.
89. Wu KH, Cui YM, Guo JF, et al. Population pharmacokinetics of cyclosporine in clinical renal transplant patients. *Drug Metab Dispos* 2005;33:1268-75.
90. Ogasawara K, Chitnis SD, Gohh RY, et al. Multidrug resistance-associated protein 2 (MRP2/ABCC2) haplotypes significantly affect the pharmacokinetics of tacrolimus in kidney transplant recipients. *Clin Pharmacokinet* 2013;52:751-62.
91. Størset E, Holford N, Midtvedt K, et al. Importance of hematocrit for a tacrolimus target concentration strategy. *Eur J Clin Pharmacol* 2014;70:65-77.
92. Veličković-Radovanović RM, Janković SM, Milovanović JR, et al. Variability of mycophenolic acid elimination in the renal transplant recipients - population pharmacokinetic approach. *Ren Fail* 2015;37:652-8.
93. Schulak JA, Hricik DE. Kidney transplantation in the elderly. *Geriatr Nephrol Urol* 1991;1:105-12.
94. Doyle SE, Matas AJ, Gillingham K, et al. Predicting clinical outcome in the elderly renal transplant recipient. *Kidney Int* 2000;57:2144-50.
95. Otero-Raviña F, Rodríguez-Martínez M, Gude F, et al. Renal transplantation in the elderly: does patient age determine the results? *Age Ageing* 2005;34:583-7.
96. Huang E, Poommipanit N, Sampaio MS, et al. Intermediate-term outcomes associated with kidney transplantation in recipients 80 years and older: an analysis of the OPTN/UNOS database. *Transplantation* 2010;90:974-9.
97. Impedovo SV, Ditunno P, Ricapito V, et al. Advanced age is not an exclusion criterion for kidney transplantation. *Transplant Proc* 2013;45:2650-3.
98. Hatamizadeh P, Molnar MZ, Streja E, et al. Recipient-related predictors of kidney transplantation outcomes in the elderly. *Clin Transplant* 2013;27:436-43.
99. Wu C, Evans I, Joseph R, et al. Comorbid conditions in kidney transplantation: association with graft and patient survival. *J Am Soc Nephrol* 2005;16:3437-44.
100. Dreyer GJ, Hemke AC, Reinders ME, et al. Transplanting the elderly: Balancing aging with histocompatibility. *Transplant Rev (Orlando)* 2015;29:205-11.
101. Cameron JS. Renal transplantation in the elderly. *Int Urol Nephrol* 2000;32:193-201.
102. Ojo AO, Hanson JA, Wolfe RA, et al. Long-term survival in renal transplant recipients with graft function. *Kidney Int* 2000;57:307-13.
103. Lehner LJ, Staack O, Halleck F, et al. Need for optimized immunosuppression in elderly kidney transplant recipients. *Transplant Rev (Orlando)* 2015;29:237-9.
104. Welzl K, Kern G, Mayer G, et al. Effect of different immunosuppressive drugs on immune cells from young and old healthy persons. *Gerontology* 2014;60:229-38.
105. Meier-Kriesche HU, Ojo A, Hanson J, et al. Increased Immunosuppressive Vulnerability in Elderly Renal Transplant Recipients. *Transplantation* 2000;69:885-9.
106. Danovitch GM, Gill J, Bunnapradist S. Immunosuppression of the elderly kidney transplant recipient. *Transplantation* 2007;84:285-91.
107. ANZDATA Registry. The 39th Annual ANZDATA Report Chapter 2: Prevalence of End Stage Kidney Disease 2016.
108. McLachlan AJ, Pont LG. Drug metabolism in older people--a key consideration in achieving optimal outcomes with medicines. *J Gerontol A Biol Sci Med Sci* 2012;67:175-80.
109. Andrés A, Budde K, Clavien PA, et al. A randomized trial comparing renal function in older kidney transplant

patients following delayed versus immediate tacrolimus administration. *Transplantation* 2009;88:1101-8.
110. Staatz CE, Goodman LK, Tett SE. Effect of CYP3A

and ABCB1 Single Nucleotide Polymorphisms on the Pharmacokinetics and Pharmacodynamics of Calcineurin Inhibitors: Part II. *Clin Pharmacokinet* 2010;49:207-21.

Cite this article as: Cossart AR, Cottrell WN, Campbell SB, Isbel NM, Staatz CE. Characterizing the pharmacokinetics and pharmacodynamics of immunosuppressant medicines and patient outcomes in elderly renal transplant patients. *Transl Androl Urol* 2019;8(Suppl 2):S198-S213. doi: 10.21037/tau.2018.10.16