The REMARK checklist

| Item | to be reported | Reported on Page Number/Line Number | Reported on Section/Paragraph | | |
|--------|--|---|----------------------------------|--|--|
| INTR | INTRODUCTION | | | | |
| 1 | State the marker examined, the study objectives, and any pre-specified hypotheses. | Page 4/Line 106-114 | Introduction/Paragraph 3 | | |
| MATE | ERIALS AND METHODS | | | | |
| Patie | nts | | | | |
| 2 | Describe the characteristics (e.g., disease stage or co-morbidities) of the study patients, including their source and inclusion and exclusion criteria. | Page 4/Line 118-128 | Methods/Paragraph 1 | | |
| 3 | Describe treatments received and how chosen (e.g., randomized or rule-based). | Page 4/Line 118-128 | Methods/Paragraph 1 | | |
| Speci | men characteristics | | | | |
| 4 | Describe type of biological material used (including control samples) and methods of preservation and storage. | Page 4/Line 118-128 | Methods/Paragraph 1 | | |
| Assay | v methods | | | | |
| 5 | Specify the assay method used and provide (or reference) a detailed protocol, including specific reagents or kits used, quality control procedures, reproducibility assessments, quantitation methods, and scoring and reporting protocols. Specify whether and how assays were performed blinded to the study endpoint. | Page 6/Line 181-Page 7/Line 203 | Methods/Paragraph 5 | | |
| Study | design | 1 | 1 | | |
| 6 | State the method of case selection, including whether prospective or retrospective and whether stratification or matching (e.g., by stage of disease or age) was used. Specify the time period from which cases were taken, the end of the follow-up period, and the median follow-up time. | Page 4/Line 118-128 | Methods/Paragraph 1 | | |
| 7 | Precisely define all clinical endpoints examined. | Page 4/Line 122-126 | Methods/Paragraph 1 | | |
| 8 | List all candidate variables initially examined or considered for inclusion in models. | Page 4/Line 118-128 | Methods/Paragraph 1 | | |
| 9 | Give rationale for sample size; if the study was designed to detect a specified effect size, give the target power and effect size. | Page 4/Line 118-128 | Methods/Paragraph 1 | | |
| Statis | tical analysis methods | • | • | | |
| 10 | Specify all statistical methods, including details of any variable selection procedures and other model-building issues, how model assumptions were verified, and how missing data were handled. | Page 6/Line 182-Page 7/Line 203 | Methods/Paragraph 5 | | |
| 11 | Clarify how marker values were handled in the analyses; if relevant, describe methods used for cutpoint determination. | Page 7/Line 202-203 | Methods/Paragraph 5 | | |

| RESULTS | | | | | |
|---------|---|-----------------------------------|--------------------------|--|--|
| Data | | | | | |
| 12 | Describe the flow of patients through the study, including the number of patients included in each stage of the analysis (a diagram may be helpful) and reasons for dropout. Specifically, both overall and for each subgroup extensively examined report the numbers of patients and the number of events. | Page 7/Line207-214 | Results/Paragraph 1 | | |
| 13 | Report distributions of basic demographic characteristics (at least age and sex), standard (disease-specific) prognostic variables, and tumor marker, including numbers of missing values. | Page 8/Line 232-233 | Results/Paragraph 3 | | |
| Analy | sis and presentation | | | | |
| 14 | Show the relation of the marker to standard prognostic variables. | Page 8/Line 233-237 | Results/Paragraph 3 | | |
| 15 | Present univariable analyses showing the relation between the marker and outcome, with the estimated effect (e.g., hazard ratio and survival probability). Preferably provide similar analyses for all other variables being analyzed. For the effect of a tumor marker on a time-to-event outcome, a Kaplan-Meier plot is recommended. | Page8/Line 260-Page9/Line 285 | Results/Paragraph 6-8 | | |
| 16 | For key multivariable analyses, report estimated effects (e.g., hazard ratio) with confidence intervals for the marker and, at least for the final model, all other variables in the model. | Page 9/Line 282-285 | Results/Paragraph 8 | | |
| 17 | Among reported results, provide estimated effects with confidence intervals from an analysis in which the marker and standard prognostic variables are included, regardless of their statistical significance. | Page8/Line 233-237 | Results/Paragraph 3 | | |
| 18 | If done, report results of further investigations, such as checking assumptions, sensitivity analyses, and internal validation. | Page 8/Line 285-288 | Results/Paragraph 8 | | |
| DISC | NOISSION | 1 | 1 | | |
| 19 | Interpret the results in the context of the pre-specified hypotheses and other relevant studies; include a discussion of limitations of the study. | Page 10/Line 324-Page 13/Line 402 | Discussion/Paragraph 1-7 | | |
| 20 | Discuss implications for future research and clinical value. | Page 13/Line 403-408 | Discussion/Paragraph 8 | | |

From: McShane LM, Altman DG, Sauerbrei W, Taube SE, Gion M, Clark GM: Reporting recommendations for tumor marker prognostic studies (REMARK). J Natl Cancer Inst 2005; 97: 1180-1184.

Article information: https://dx.doi.org/10.21037/tau-21-1130

*As the checklist was provided upon initial submission, the page number/line number reported may be changed due to copy editing and may not be referable in the published version. In this case, the section/paragraph may be used as an alternative reference.