



The very low magnetic resonance imaging apparent diffusion coefficient (ADC) measure of abscess is likely due to pus's specific T2 relaxation time

Yi Xiang J. Wáng

Department of Imaging and Interventional Radiology, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong SAR, China

Correspondence to: Yi Xiang J. Wáng, MMed, PhD. Department of Imaging and Interventional Radiology, Faculty of Medicine, The Chinese University of Hong Kong, 30-32 Ngan Shing Street, Shatin, New Territories, Hong Kong SAR, China. Email: yixiang_wang@cuhk.edu.hk

Submitted Sep 24, 2023. Accepted for publication Oct 7, 2023. Published online Oct 13, 2023.

doi: 10.21037/qims-23-1363

View this article at: <https://dx.doi.org/10.21037/qims-23-1363>

It is well known that abscess fluid (i.e., pus) tends to demonstrate a very low magnetic resonance imaging (MRI)-derived apparent diffusion coefficient (ADC) regardless of the location of the abscess (1-8). At least in the brain, it may appear counterintuitive that abscess pus, being fluid or semi-fluid, has an ADC measure lower than those of white/grey matters (*Figure 1*).

Recently we noted that, regardless of whether $b=0$ data are included for the ADC calculation, the T2 relaxation time (T2 time) of a tissue or an *in vivo* substance is strongly associated with ADC measure in many scenarios (9-13) (*Figure S1*). While ADC measure is affected by many factors, a T2 time of around 70 ms (e.g., from 60 to 80 ms) at 3 T or (or its equivalent values at other magnetic fields) may be associated with the lowest ADC measure (*Figure 2*). It may be further inferred that, if a tissue (or an *in vivo* substance) has a T2 time of slightly less than half that of body water (such as gallbladder water: 172 ms at 3 T), then this tissue (or *in vivo* substance) will have a low ADC measure.

Following the discussion above, we looked at whether abscess pus has a T2 time of slightly less than half that of body water. We were able to identify four studies that reported the T2 times of abscess pus and a body fluid. We took the ratios of '*abscess pus T2 time/body water T2 time*', and

the results are shown in *Figure 3* (16-21). It is shown that abscess pus has a T2 time of about half that of body water. Wall *et al.* reported a T2 time for abscess of 81 ms, and T2 time for muscle, liver and urine of 29, 45 and 166 ms, respectively. The T2 times of the later three are consistent with the other reports at 1.5 or 3.0 T (12,22,23). There is a notion that T2 time doesn't change much over the range of field strengths used for routine clinical MRI (0.2 to 3.0 T) (23). Additionally, Subhawong *et al.* (24) described a case series of soft tissue masses, which included one abscess case and three cases of ganglion cyst. T2 signal ratio was measured as '*lesion T2 weighting signal intensity/muscle T2 weighting signal intensity*'. The abscess had a T2 signal ratio of 1.48, while the three ganglion cysts had a mean T2 signal ratio of 3.42, with the abscess' value being 0.43 of those of the ganglion cysts (abscess ADC: 0.63 mm²/s, ganglion cyst ADC mean: 2.49 mm²/s). The specific T2 relaxation time of pyogenic abscess fluid, according to our viewpoint, contributes to very low ADC measured by MRI. We argue that abscess pus may not have truly restricted diffusion compared with many other *in vivo* solid tissues. Of course, in real practice, abscess pus composition may vary, and so does its ADC measure (1,4). The discussion in this letter refers to the common scenarios.

Now we look at *Figure 1* again. Brain tissues are




Figure 1 MRI of a 37-year-old woman with pyogenic abscess in the left frontal lobe (A: T2-weighted image; B: diffusion-weighted image; C: ADC map; D: contrast-enhanced T1-weighted image). The central component of the lesion shows high signal on T2-weighted image, high signal on diffusion-weighted image (asterisk in B) and low value on ADC map (C, lower than the brain tissues). Arrow in (D) denotes a ring enhancement after contrast agent administration. It appears unreasonable that the abscess content has a so low ADC measure. Adapted from Feraco *et al.* (1). MRI, magnetic resonance imaging; ADC, apparent diffusion coefficient.




Figure 2 Relationship between T2 time and ADC at 3 T. This graph is modified from Wáng and Ma (12). For data sources also see Wáng and Ma (12). The vertical dotted line at X-axis 172 ms denotes that 172 ms is the T2 time of *in vivo* body water at 3 T [gallbladder water as an example (14,15)]. The vertical dotted line at X-axis 70 ms denotes that a T2 time around 70 ms at 3 T is associated with very low ADC measure. ADC, apparent diffusion coefficient; ms, millisecond.

generally noted to have a short T2 time with grey/dark signal on T2 weighted image. For 3T data, Wansapura *et al.* (25) described that the average T2 values for occipital and frontal gray matter are 41.6 and 51.8 ms, respectively, and average T2 values for occipital and frontal white matter are 48.4 and 44.7 ms, respectively. According to *Figure 3*, abscess pus is roughly estimated to have a T2 time of around 80 ms at 3T. Therefore, according to *Figure 2*, an increase of T2 time from grey/white matter values toward 80 ms would be associated with a lower ADC measure. The abscess high signal shown on *Figure 1B* (a high *b*-value diffusion-weighted image) likely does not reflect restricted fluid diffusivity, instead reflects the specific T2 time of the abscess fluid.




Figure 3 Ratios of ‘abscess pus T2 time/body water T2 time’ from four studies. 3-T ref is the ratio of 70 ms (the T2 time associated with the lowest ADC according to *Figure 2*)/172 ms (*in vivo* body water T2 time). Wall *et al.* (16) and Brown *et al.* (17): various abscess fluid T2 time/urine T2 time; Cieszanowski *et al.* (18): liver abscess fluid/liver cyst fluid. Yilmaz *et al.* (19): odontogenic jaw abscess fluid/odontogenic jaw cyst fluid. T2 time measurement is affected by magnetic field strength and homogeneity, pulse sequence, and data acquisition parameters, etc., therefore the ratio of ‘abscess pus T2 time/body water T2 time’ in the same study is presented instead of absolute T2 time value. Note that the composition of a body fluid may affect its T2 time (20,21). ADC, apparent diffusion coefficient; ref, reference; ms, millisecond.

Acknowledgments

Funding: None.

Footnote

Conflicts of Interest: The author has completed the ICMJE uniform disclosure form (available at <https://qims>).

amegroups.com/article/view/10.21037/qims-23-1363/coif). YXJW serves as the Editor-in-Chief of *Quantitative Imaging in Medicine and Surgery*. The author has no other conflicts of interest to declare.

Ethical Statement: The authors is accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Feraco P, Donner D, Gagliardo C, Leonardi I, Piccinini S, Del Poggio A, Franciosi R, Petralia B, van den Hauwe L. Cerebral abscesses imaging: A practical approach. *J Popul Ther Clin Pharmacol* 2020;27:e11-24.
2. Erdogan C, Hakyemez B, Yildirim N, Parlak M. Brain abscess and cystic brain tumor: discrimination with dynamic susceptibility contrast perfusion-weighted MRI. *J Comput Assist Tomogr* 2005;29:663-7.
3. Chang SC, Lai PH, Chen WL, Weng HH, Ho JT, Wang JS, Chang CY, Pan HB, Yang CF. Diffusion-weighted MRI features of brain abscess and cystic or necrotic brain tumors: comparison with conventional MRI. *Clin Imaging* 2002;26:227-36.
4. Lotan E, Hoffmann C, Fardman A, Ziv-Baran T, Komisar O, Harnof S. Postoperative versus Spontaneous Intracranial Abscess: Diagnostic Value of the Apparent Diffusion Coefficient for Accurate Assessment. *Radiology* 2016;281:168-74.
5. Chou CP, Chiou SH, Levenson RB, Huang JS, Yang TL, Yu CC, Chiang AJ, Pan HB. Differentiation between pelvic abscesses and pelvic tumors with diffusion-weighted MR imaging: a preliminary study. *Clin Imaging* 2012;36:532-8.
6. Oruç E, Yıldırım N, Topal NB, Kılıçturgay S, Akgöz S, Savcı G. The role of diffusion-weighted MRI in the classification of liver hydatid cysts and differentiation of simple cysts and abscesses from hydatid cysts. *Diagn Interv Radiol* 2010;16:279-87.
7. Oto A, Schmid-Tannwald C, Agrawal G, Kayhan A, Lakadamyali H, Orrin S, Sethi I, Sammet S, Fan X. Diffusion-weighted MR imaging of abdominopelvic abscesses. *Emerg Radiol* 2011;18:515-24.
8. Harish S, Chiavaras MM, Kotnis N, Rebello R. MR imaging of skeletal soft tissue infection: utility of diffusion-weighted imaging in detecting abscess formation. *Skeletal Radiol* 2011;40:285-94.
9. Yu WL, Xiao BH, Ma FZ, Zheng CJ, Tang SN, Wáng YXJ. Underestimation of the spleen perfusion fraction by intravoxel incoherent motion MRI. *NMR Biomed* 2023;36:e4987.
10. Wáng YXJ, Zhao KX, Ma FZ, et al. The contribution of T2 relaxation time to MRI-derived apparent diffusion coefficient (ADC) quantification and its potential clinical implications. *Quant Imaging Med Surg* 2023;13:7410-6.
11. Wang YXJ. Complicated relationships between tissue T2 relaxation time and in vivo tissue diffusion measures, depending on the ranges of T2 value. arXiv:2306.10657. [submitted on 19 Jun 2023]. Available online: <https://doi.org/10.48550/arXiv.2306.10657>
12. Wáng YXJ, Ma FZ. A tri-phasic relationship between T2 relaxation time and magnetic resonance imaging (MRI)-derived apparent diffusion coefficient (ADC). *Quant Imaging Med Surg* 2023. doi: 10.21037/qims-23-1342.
13. Oh J, Cha S, Aiken AH, Han ET, Crane JC, Stainsby JA, Wright GA, Dillon WP, Nelson SJ. Quantitative apparent diffusion coefficients and T2 relaxation times in characterizing contrast enhancing brain tumors and regions of peritumoral edema. *J Magn Reson Imaging* 2005;21:701-8.
14. Mohajeri S, Ijare OB, Bezabeh T, King SB, Thomas MA, Minuk G, Lipschitz J, Kirkpatrick I, Smith M, Smith IC. In vivo 1H MRS of human gallbladder bile at 3 T in one and two dimensions: detection and quantification of major biliary lipids. *NMR Biomed* 2014;27:1192-202.
15. Gajdošík M, Chmelík M, Halilbasic E, Pflieger L, Klepochová R, Trauner M, Trattng S, Krššák M. In Vivo (1) H MR Spectroscopy of Biliary Components of Human Gallbladder at 7T. *J Magn Reson Imaging* 2021;53:98-107.
16. Wall SD, Fisher MR, Amparo EG, Hricak H, Higgins CB. Magnetic resonance imaging in the evaluation of abscesses. *AJR Am J Roentgenol* 1985;144:1217-21.
17. Brown JJ, vanSonnenberg E, Gerber KH, Strich G, Wittich GR, Slutsky RA. Magnetic resonance relaxation

- times of percutaneously obtained normal and abnormal body fluids. *Radiology* 1985;154:727-31.
18. Cieszanowski A, Anysz-Grodzicka A, Szeszkowski W, Kaczynski B, Maj E, Gornicka B, Grodzicki M, Grudzinski IP, Stadnik A, Krawczyk M, Rowinski O. Characterization of focal liver lesions using quantitative techniques: comparison of apparent diffusion coefficient values and T2 relaxation times. *Eur Radiol* 2012;22:2514-24.
 19. Yilmaz UN, Yaman F, Atilgan SS. MR T1 and T2 relaxations in cysts and abscesses measured by 1.5 T MRI. *Dentomaxillofac Radiol* 2012;41:385-91.
 20. Daoust A, Dodd S, Nair G, Bouraoud N, Jacobson S, Walbridge S, Reich DS, Koretsky A. Transverse relaxation of cerebrospinal fluid depends on glucose concentration. *Magn Reson Imaging* 2017;44:72-81.
 21. Yoshimura S, Tanaka H, Kawabata S, Kozawa J, Takahashi H, Hidaka Y, Hotta M, Kashiwagi N, Tomiyama N. Effect of urinary glucose concentration and pH on signal intensity in magnetic resonance images. *Jpn J Radiol* 2022;40:930-8.
 22. de Bazelaire CM, Duhamel GD, Rofsky NM, Alsop DC. MR imaging relaxation times of abdominal and pelvic tissues measured in vivo at 3.0 T: preliminary results. *Radiology* 2004;230:652-9.
 23. Bottomley PA, Foster TH, Argersinger RE, Pfeifer LM. A review of normal tissue hydrogen NMR relaxation times and relaxation mechanisms from 1-100 MHz: dependence on tissue type, NMR frequency, temperature, species, excision, and age. *Med Phys* 1984;11:425-48.
 24. Subhawong TK, Durand DJ, Thawait GK, Jacobs MA, Fayad LM. Characterization of soft tissue masses: can quantitative diffusion weighted imaging reliably distinguish cysts from solid masses? *Skeletal Radiol* 2013;42:1583-92.
 25. Wansapura JP, Holland SK, Dunn RS, Ball WS Jr. NMR relaxation times in the human brain at 3.0 tesla. *J Magn Reson Imaging* 1999;9:531-8.

Cite this article as: Wáng YXJ. The very low magnetic resonance imaging apparent diffusion coefficient (ADC) measure of abscess is likely due to pus's specific T2 relaxation time. *Quant Imaging Med Surg* 2023;13(12):8881-8885. doi: 10.21037/qims-23-1363




Figure S1 Almost perfect linear relationship between brain glioma T2 time and ADC. T2 time ranges from 71 to 283 ms. Low ADC is noted for glioma with T2 time of around 80 ms (however, there is no data with T2 time shorter than 60 ms). Data are extracted from the study of Oh *et al.* (13), acquired using a 1.5-T scanner and with ADC based on b -value of 0 and $1,000 \text{ s}/\text{mm}^2$. ADC, apparent diffusion coefficient; ms, millisecond.