



Patients' death: "Everybody hurts... sometimes"*

Rosa Tesoro¹, Monica Escher²

¹Department of Psychiatry, University Hospitals of Geneva, Geneva, Switzerland; ²Department of Rehabilitation and Geriatrics, Division of Palliative Medicine, University Hospitals of Geneva, Geneva, Switzerland

Correspondence to: Rosa Tesoro, MD. Department of Psychiatry, University Hospitals of Geneva, Geneva, Switzerland. Email: rosa.tesoro@hcuge.ch.

Comment on: Wiemann B, Ketteler E, Fahy B. Surgeon and medical student response to patient death. *Ann Palliat Med* 2023;12:70-80.

Keywords: Patient death; burnout; mental health; palliative care; surgeons

Submitted Mar 01, 2023. Accepted for publication Apr 30, 2023. Published online May 08, 2023.

doi: 10.21037/apm-23-257

View this article at: <https://dx.doi.org/10.21037/apm-23-257>

Patients die. This is a reality physicians will face all along their career. What happens when a surgeon takes his sterile gloves off after a patient's death and goes back home? What are his feelings? How does he cope?

Brianne Wiemann, Erika Ketteler and Bridget Fahy from New Mexico tried to find answers to these questions in their article published this month (1). Surgeons and medical students surveyed by our colleagues reported emotional distress manifested as anxiety and burnout. One out of 4 younger respondents tried not to think about the patient's death. The most common coping strategies were seeking comfort from loved ones and extracurricular activities. Emotional burden was heightened for COVID-19 related deaths since support from colleagues was limited and concerns for one's own and loved ones' safety were present. Moreover, almost half of the respondents indicated they had not received any training on how to deal with a patient's death.

The COVID-19 pandemic drew attention to the impact patient death can have on physicians' mental health. Physicians have a higher rate of burnout, depression, and suicide than the general population (2), but some physician groups may be at higher risk than others. Surgeons are described as having their own "moral economy". Not offering surgery is experienced as giving up on a patient, and this view is strengthened by the expectations from patients, families, and society (3). Surgeon professional identity includes an acute sense of being responsible for patient outcomes (4), while a focused-on-surgery attitude

leads to mainly consider "on the table death" and to operate when the benefits or the postoperative quality of life are uncertain (5). Such attitudes ill prepare surgeons to face patient death, all the more so that it is a comparatively rare event (6,7). Surgeons' typical traits are described as main reasons to choose this specialty, but also as important stress factors: commitment to patients, long hours work, high volume and performance level, decisions about life and death. Since surgical education traditionally focus on the care by a surgical procedure, death is perceived as a surgical failure (8). Surgeons could experience some difficulties discussing with patients with incurable diseases to avoid culpability feelings (9). They negotiate with patients a package of post-operative care prior to undertaking high-risk surgical procedures to shift responsibility of poor outcomes on patient's decision for treatment limitation (10) as a defence from a strong sense of responsibility for surgical outcomes. Pressure to succeed has already been described in sociological studies that focused on the image of a surgeon who feels himself as the only responsible of the patient's life.

Physicians regularly exposed to patient death come to integrate it within the scope of their clinical activity and seem to better come to terms with it (11).

When medical students are exposed to patients' death, senior personnel's empathy make the process easier. It seems important to not forget empathy with experience: looking for support and sharing experiences with colleagues still remains an important coping mechanism. Residents and fellows from the Mayo International Health Program were

* Everybody Hurts, R.E.M. 1992.

interviewed about their participation to International Health Electives in low or lower-middle-income countries (12). They reported a lack of preparation for exposure to death and a lack of closure. Even though this study explored an atypical experience, institutions should pay attention to these two aspects in undergraduate, postgraduate and continuing medical education (12).

Medical students and junior physicians represent a vulnerable group as far as mental health is concerned. Experiencing the death of a patient provokes various emotions in medical students, who report feeling upset, sad, and helpless (2,13). Although they are capable of positively transforming the experience into a professional and a life lesson, they express the need for support from senior physicians. The first clinical years are marked by an increased risk of developing depression. Prevalence rates among residents range between 20.9% and 43.2% (14). Most importantly, working conditions have been identified as a risk factor. Long working hours and imbalance between professional and family life compounded by study and examinations requirements are associated with mental health disorders (15).

Institutions should pay attention to the impact of patient death on physicians at every stage of their career. Organizational and physician-directed interventions showed effectiveness in protecting the mental health of medical staff. Strategies to help physicians and medical students better cope with a patient's death and improve their skills in caring for a dying patient could include better integration of palliative and end of life care in the undergraduate and postgraduate curriculum, systematic debriefings of the healthcare team after exposure to emotionally traumatic events, setting up of rituals intended to create time and space for pausing and reflecting, and easy access to a mental health provider. Interpersonal support is a valued and common coping strategy, and senior physicians should be educated in being effective mentors, capable of responding to medical students' and junior physicians' emotional and professional needs.

A paradigm shift also seems essential. Physicians are less likely to seek for mental health care than non-physicians. Barriers include fears concerning confidentiality and negative consequences for their career. Away from the image of the hero physician or the long-suffering physician, the medical profession and society should allow more space for the physicians' emotions, not as a sign of weakness, but as an essential component of a life in tune with its deeper parts. It is not just a matter of supporting individuals since

burnout physicians are more likely to report medical errors and an intent to leave their job (16,17). Paying attention to the well-being of physicians is a responsibility of the medical community towards patients and society at large.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Annals of Palliative Medicine*. The article did not undergo external peer review.

Conflicts of Interest: Both authors have completed the ICMJE uniform disclosure form (available at <https://apm.amegroupp.com/article/view/10.21037/apm-23-257/coif>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Wiemann B, Ketteler E, Fahy B. Surgeon and medical student response to patient death. *Ann Palliat Med* 2023;12:70-80.
2. Harvey SB, Epstein RM, Glozier N, et al. Mental illness and suicide among physicians. *Lancet* 2021;398:920-30.
3. Morris RS, Ruck JM, Conca-Cheng AM, et al. Shared Decision-Making in Acute Surgical Illness: The Surgeon's Perspective. *J Am Coll Surg* 2018;226:784-95.
4. Rivard SJ, Vitous CA, De Roo AC, et al. "The captain of the ship." A qualitative investigation of surgeon identity formation. *Am J Surg* 2022;224:284-91.

5. Tesoro R, Suppan M, Dupuis A, et al. Futility of end-of-life and emergency surgery in extreme high-risk patients: anesthesiologists' versus surgeons' perspective. *Braz J Anesthesiol* 2022;72:434-6.
6. Fowler AJ, Wahedally MAH, Abbott TEF, et al. Death after surgery among patients with chronic disease: prospective study of routinely collected data in the English NHS. *Br J Anaesth* 2022;128:333-42.
7. Vascular Events in Noncardiac Surgery Patients Cohort Evaluation (VISION) Study Investigators, Spence J, LeManach Y, et al. Association between complications and death within 30 days after noncardiac surgery. *CMAJ* 2019;191:E830-7.
8. Joliat GR, Demartines N, Uldry E. Systematic review of the impact of patient death on surgeons. *Br J Surg* 2019;106:1429-32.
9. Milch RA. The surgeon-patient relationship in advanced illness. *Surg Oncol Clin N Am* 2001;10:25-30.
10. Schwarze ML, Bradley CT, Brasel KJ. Surgical "buy-in": the contractual relationship between surgeons and patients that influences decisions regarding life-supporting therapy. *Crit Care Med* 2010;38:843-8.
11. Whitehead PR. The lived experience of physicians dealing with patient death. *BMJ Support Palliat Care* 2014;4:271-6.
12. Bashir MU, Nordhues HC, Merry SP, et al. Dealing With Patient Death on International Health Electives: A Qualitative Study of Residents' Experiences. *Acad Med* 2019;94:1170-7.
13. Ho CY, Kow CS, Chia CHJ, et al. The impact of death and dying on the personhood of medical students: a systematic scoping review. *BMC Med Educ* 2020;20:516.
14. Mata DA, Ramos MA, Bansal N, et al. Prevalence of Depression and Depressive Symptoms Among Resident Physicians: A Systematic Review and Meta-analysis. *JAMA* 2015;314:2373-83.
15. Hughes TM, Waljee JF, Fang Y, et al. New-Onset Depression Among Surgical Interns. *JAMA Surg* 2022;157:543-5.
16. Menon NK, Shanafelt TD, Sinsky CA, et al. Association of Physician Burnout With Suicidal Ideation and Medical Errors. *JAMA Netw Open* 2020;3:e2028780.
17. Schwarzkopf D, Rüdell H, Thomas-Rüdell DO, et al. Perceived Nonbeneficial Treatment of Patients, Burnout, and Intention to Leave the Job Among ICU Nurses and Junior and Senior Physicians. *Crit Care Med* 2017;45:e265-73.

Cite this article as: Tesoro R, Escher M. Patients' death: "Everybody hurts... sometimes". *Ann Palliat Med* 2023;12(4):673-675. doi: 10.21037/apm-23-257