



# Effect of levocetirizine hydrochloride on the growth of human dermal papilla cells: a preliminary study

Sijian Wen<sup>#</sup>, Jumei Wei<sup>#</sup>, Jiajuan Bao, Ting Guo, Wenjun Zheng, Xiaosheng Zhuang, Youkun Lin

Department of Dermatology, The First Affiliated Hospital of Guangxi Medical University, Nanning 530021, China

*Contributions:* (I) Conception and design: S Wen, Y Lin, J Wei; (II) Administrative support: X Zhuang; (III) Provision of study materials or patients: T Guo, W Zheng; (IV) Collection and assembly of data: J Wei, J Bao; (V) Data analysis and interpretation: S Wen, J Wei; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

<sup>#</sup>These authors contributed equally to this work.

*Correspondence to:* Youkun Lin. Department of Dermatology, The First Affiliated Hospital of Guangxi Medical University, Nanning 530021, China.

Email: linyoukun7@aliyun.com.

**Background:** To conduct an in vitro investigation into the effect of different concentrations of levocetirizine hydrochloride on the growth of human dermal papilla cells (hDPCs) the underlying mechanisms involved.

**Methods:** hDPCs were cultured in Dulbecco's Modified Eagle Medium (DMEM) containing different concentrations of levocetirizine hydrochloride for 48 h. The growth of hDPCs was observed by immunofluorescence staining, and the cell proliferation was detected by MTT assay. After the hDPCs were cultured in DMEM containing 1, 10, 100, 1,000, and 10,000 ng/mL levocetirizine hydrochloride for 48 h, the mRNA expressions of cyclooxygenase 2 (COX-2), prostaglandin D2 synthase (PTGDS), prostaglandin E2 (PGE2), prostaglandin F2 $\alpha$  (PGF2 $\alpha$ ), G protein-coupled receptor 44 (GPR44), protein kinase B (AKT), and glycogen synthase kinase 3 $\beta$  (GSK3 $\beta$ ) were determined by real-time fluorescence-based quantitative polymerase chain reaction (PCR), and the protein expressions of PTGDS, phosphorylated protein kinase B (pAKT), and phosphorylated glycogen synthase kinase 3 $\beta$  (pGSK3 $\beta$ ) were detected by Western blotting. After the hDPCs were cultured in DMEM containing 1, 10, 100, 1,000, 10,000 ng/mL levocetirizine hydrochloride for 24 h, the secretion levels of prostaglandin D2 (PGD2) and PGD2 receptor (PGD2R) in the culture supernatant were determined by enzyme-linked immunosorbent assay (ELISA). One-way analysis of variance (ANOVA) was performed using SPSS 17.0 software, and the LSD-t test was used for pairwise comparisons.

**Results:** Immunofluorescence staining showed that hDPCs in the 100 ng/mL group grew well, with over 90% confluency. Methyl thiazolyl tetrazolium (MTT) method showed that the proliferation rate of hDPCs significantly differed between different levocetirizine hydrochloride groups and the blank control group ( $F=42.22$ ,  $P<0.05$ ), while the proliferation rate was significantly higher in the 100 ng/mL group ( $115.80\pm 5.10\%$ ) than in the blank control group (100%) ( $t=28.26$ ,  $P<0.05$ ). The relative mRNA expressions of COX-2, PGF2 $\alpha$ , PTGDS, GPR44, and AKT showed significant differences in different levocetirizine hydrochloride groups (the F values were 1.97, 3.66, 2.17, 2.66, and 7.32, respectively; all  $P<0.05$ ), whereas the mRNA expressions of PGE2 and GSK3 $\beta$  showed no significant difference ( $F=0.87$ ,  $F=1.19$ , respectively; both  $P>0.05$ ). The mRNA expressions of COX-2, PTGDS, and GPR44 in the 100 ng/mL group (0.84 $\pm$ 0.08, 0.81 $\pm$ 0.10, and 0.85 $\pm$ 0.09, respectively) were significantly lower than those in the blank control group ( $t=1.97$ ,  $t=2.17$ , and  $t=2.65$ , respectively; all  $P<0.05$ ), whereas the mRNA expressions of PGF2 $\alpha$  and AKT in the 100 ng/mL group (1.96 $\pm$ 0.25 and 1.74 $\pm$ 0.32, respectively) were significantly higher than those in the blank control group ( $t=3.662$  and  $t=7.325$ , respectively; both  $P<0.05$ ). There were significant differences in the levels of PTGDS, pAKT, pGSK3 $\beta$ , PGD2, and PGD2R proteins between the different levocetirizine hydrochloride groups (the F values were 11.84, 3.89, 4.07, 66.15, and 44.33, respectively). The protein expressions of PTGDS, PGD2, and PGD2R in the 100 ng/mL group (0.32 $\pm$ 0.05, 141.62 $\pm$ 5.44, and 215.08 $\pm$ 9.55, respectively) were significantly lower than those in the blank control group (0.73 $\pm$ 0.06, 180.08 $\pm$ 6.15, and 273.24 $\pm$ 3.18,

respectively) (the *t* values were 5.66, 45.07, and 92.05, respectively; all *P*<0.05), whereas the protein expressions of pAKT and pGSK3 $\beta$  in the 100 ng/mL group (0.59 $\pm$ 0.05 and 0.46 $\pm$ 0.03, respectively) were significantly higher than those in the blank control group (0.46 $\pm$ 0.02 and 0.35 $\pm$ 0.042, respectively) (*t*=16.59, *t*=7.73, respectively; both *P*<0.05).

**Conclusions:** Levocetirizine hydrochloride may promote the growth and proliferation of hDPC in vitro by inhibiting the PGD2-GPR44 pathway and activating the AKT signaling pathway.

**Keywords:** Cetirizine; prostaglandin; alopecia; G protein-coupled receptor; dermal papilla cells (DPCs)

Submitted Nov 11, 2019. Accepted for publication Jan 08, 2020.

doi: 10.21037/apm.2020.01.15

View this article at: <http://dx.doi.org/10.21037/apm.2020.01.15>

## Introduction

Androgenetic alopecia (AGA) is an androgen-dependent hereditary disorder influenced by a variety of genetic and environmental factors. AGA is the most common type of alopecia and has a chronic disease course, with its main symptoms being progressive hair loss and hair follicle atrophy, which may be accompanied by greasy scalp, dandruff, and itching. Since hair loss directly affects the patient's appearance, it often has a significant impact on the patient's social activities and mental health and may lead to a decreased quality of life, while worsening social barriers and causing depression in severe cases (1). With recent years of socioeconomic development, many patients have pursued a more positive outcome, and the role of AGA treatment has thus been increasingly recognized.

The etiology and mechanism of AGA are not fully understood. Hair follicle atrophy, miniaturization, shortened anagen phase, prolonged telogen phase, hair follicle involvement, and abnormal growth cycle may be the key links during its pathogenesis. It is well believed that the occurrence and development of AGA are closely related to endocrine disorders, especially the abnormal metabolism of androgen in the hair follicles on the scalp (2). Drugs for AGA include hormone regulators, vasodilators, and immunomodulators; however, few drugs that can effectively promote hair growth have been available. The drugs that are commonly used for AGA are still finasteride and minoxidil; however, finasteride has been reported to cause sexual side effects such as loss of libido and impotence, whereas the use of minoxidil has been associated with irritant dermatitis, allergic contact dermatitis, headache, and hypotension (3). Although many new methods have been developed for AGA treatment including autologous hair transplantation, hair follicle reconstruction, and low-energy laser therapy (4),

these are expensive and invasive and thus cannot be widely applied.

Levocetirizine hydrochloride is a safe, highly selective, and highly affinitive new-generation histamine H1 receptor antagonist that has been widely used in the treatment of skin, respiratory, and ocular allergies. It selectively, competitively, and irreversibly binds to transmembrane G protein-coupled receptors and thus inhibits the release of various inflammatory mediators associated with allergy. As the R-enantiomer of cetirizine, levocetirizine hydrochloride has higher bioavailability and lower hepatic clearance, and its plasma binding rate can reach 95%. It is characterized by low tissue affinity, low cardiotoxicity, and low sedation effect (5). A recent study (6) revealed that topical use of cetirizine 1% significantly increased total hair density in the balding area, whereas the vellus hair density showed an evident decrease. Another study demonstrated that levocetirizine hydrochloride, by binding to COX-2, inhibited its ability to promote prostaglandin expression (7).

More evidence has suggested that the prostaglandin pathways (especially the PGDS/PGD pathway) are involved in the pathogenesis of AGA (8). Prostaglandins (PG) are a group of unsaturated fatty acids that are enzymatically metabolized. They are widely distributed in various tissues and body fluids and can mediate multiple physiological functions and inflammatory reactions such as cell proliferation, differentiation, and apoptosis. Cyclooxygenase (COX) induces PGs to be converted from arachidonic acid to unstable prostaglandin H<sub>2</sub> and then further processed into various biologically active prostaglandins (e.g., prostacyclin, PGD<sub>2</sub>, PGE<sub>2</sub>, PGF<sub>2 $\alpha$</sub> , and PGI<sub>2</sub>) or thromboxane A<sub>2</sub> (TXA<sub>2</sub>) by the corresponding prostaglandin synthetases. Comparisons of the effects of different PG subtypes on hair growth have demonstrated

that PGE2 and PGF2 $\alpha$  can promote hair growth, whereas PGD2 can inhibit hair growth and aggravate hair follicle miniaturization (9). Jeong *et al.* (10) have found that PGD2 leads to the activation of androgen receptors by regulating the DP2 and AKT/GSK3 $\beta$  signaling pathways on hDPCs. Our current study therefore investigated the effects of levocetirizine hydrochloride on the growth, proliferation, and expression of PG-related signaling factors in human hair dermal papilla cells (DPCs) and explored its effects on hair growth along with the related molecular mechanisms.

## Methods

### Reagents and equipment

The main reagents used in this study included fetal bovine serum (FBS) and Dulbecco's Modified Eagle Medium (DMEM) (Gibco, USA), levocetirizine hydrochloride powder (Huaopond Pharm, Chongqing, China), rabbit anti-human  $\alpha$  smooth muscle actin ( $\alpha$ -SMA) antibody (PTGlab, USA), Cy3-labeled goat anti-rabbit IgG (IgG-Cy3) (Beyotime, Shanghai, China), and methyl thiazolyl tetrazolium (MTT) (Sigma, USA). The reagent kits included cellular RNA, small RNA, DNA, and protein extraction kit (Magan, USA), reverse transcription, and fluorescence quantitative PCR kits (TAKARA, Japan), and BCA protein quantification kit (Beyotime, Shanghai, China). Other reagents included anti-actin antibody, anti-GAPDH antibody, prostaglandin D2 synthase (PTGDS) (Abcam, USA) as well as anti-pAKT antibody and anti-pGSK3 $\beta$  antibody (CST, USA). Kits for PGD2 enzyme-linked immunosorbent assay (ELISA) (Wuhan Huamei, China) and for PGD2 receptor ELISA (Jingmei Biotechnology, Jiangsu, China) were also used. The equipment used included microplate reader (Thermo, USA), fluorescence quantitative PCR system, (Roche, Switzerland), and gel imaging analysis system (GE, USA).

### Study methods

#### Cell culture

Human dermal papilla cells (hDPCs) (Promocell, Germany) were thawed and resuscitated and then mixed with high-glucose DMEM containing 15% FBS (15% conditioned DMEM). The mixture was cultured in an incubator (37 °C and 5% CO<sub>2</sub>) for 24 h, and the culture medium was changed every 1–2 days. When the cells were approximately 80% confluent, they were passaged and frozen. The frozen cells

were resuscitated and cultured, and hDPCs were digested at the second passage and used in the following assays.

#### Observation of hDPC growth by using the immunofluorescence technique

hDPCs were seeded in 6-well plates with a density of  $1.5 \times 10^5$ /well. In each well, 2 mL of 15% DMEM conditioned medium was added. After the cells became adherent, the cells were cultured with 15% DMEM-conditioned medium containing levocetirizine hydrochloride 0 (control), 1, 10, 100, 1,000, or 10 000 ng/mL for 48 h. After the cells were mounted and cultured on slides, they were immersed in a phosphate buffer solution with Tween<sup>®</sup> 20 (PBST), fixed in 4% paraformaldehyde, blocked with 1% BSA for 30 min, and incubated with rabbit anti-human  $\alpha$ -SMA antibody in a wet box at 4 °C overnight. After the cells were immersed in PBST again, the mixture was added with IgG-Cy3 and incubated in a dark at room temperature for 1 h. DAPI was added to the wells before observation under a fluorescence microscope. The whole procedure was repeated three times.

#### Measurement of cell proliferation rate by MTT assay

hDPCs were seeded in 96-well plates at  $1 \times 10^4$ /well, and each well was mixed with 200  $\mu$ L of 15% DMEM conditioned medium. After the cells became adherent, the cells were cultured with 15% DMEM-conditioned medium containing levocetirizine hydrochloride 0 (control), 1, 10, 50, 100, 250, 500, 1,000, 10,000, and 100,000 ng/mL for 48 h. Zero wells were set, and five duplicate wells were set for each concentration. After the cells were washed with PBST twice, serum-free DMEM medium containing MTT was added. After the mixture was incubated at 37 °C for 4 h, 150  $\mu$ L of dimethyl sulfoxide was added in each well. The mixture was incubated with agitation in dark on a plate shaker for 10 min. Absorbance (A value) was measured spectrophotometrically at 490 nm. The whole procedure was repeated three times, and the means were calculated. The cell proliferation rate was calculated using the following formula: the proliferation rate (%) = (the average value of all A values for experimental wells – the average value of all A values for zero wells)/(the average value of all A values for control wells – the average value of all A values for zero wells)  $\times$  100%.

#### Determination of the expressions of the mRNA and protein expressions of relevant genes in hDPCs by using real-time fluorescence quantitative PCR and Western blotting

hDPCs were seeded in 6-well plates at  $1.5 \times 10^5$ /well,

**Table 1** Sequences of the target gene primers

Genes	Forward primers (5'-3')	Reverse primers (5'-3')	Length (bp)
<i>COX-2</i>	GAAACCGTCTGAACTATCCTGC	GTCGTTGACCTCGTCTGTTATGT	93
<i>PTGDS</i>	GAGTTAAAGGAGAAATTCACCGC	GGGAGTCCTATTGTTCCGTCAT	109
<i>PGE2</i>	CTCCTTGCCTTTTCACGATTT	ACAACAGAGGACTGAACGCAT	169
<i>PGF2<math>\alpha</math></i>	AGGGAGATGACTTGAGTGGTTG	TTTCCGCTCTGGCAGGTTGT	116
<i>GPR44</i>	GTCATCCTCTTCGTGGTGGG	ACGGCCAAGAAGTAGGTGAAG	125
<i>AKT</i>	GGCAAGGTGATCCTGGTGAA	GTCGTGGGTCTGGAAAGAGTACT	180
<i>GSK-3<math>\beta</math></i>	CCCTCAAATTAAGGCACATCC	CACGGTCTCCAGTATTAGCATCT	239
<i><math>\beta</math> actin</i>	TAGTTGCGTTACACCCTTTCTTG	TCACCTTCACCGTTCAGTTT	194

and each well was mixed with 2 mL of 15% DMEM-conditioned medium. After the cells became adherent, the cells were cultured with 15% DMEM-conditioned medium containing levocetirizine hydrochloride 0 (control), 1, 10, 100, 1,000, or 10 000 ng/mL for 48 h. After the supernatant was discarded, the total RNA and total protein were extracted by using cellular RNA, small RNA, DNA, and protein extraction kits.

- (I) Determination of mRNA expression levels proceeded as follows. Total RNA was synthesized into a cDNA template using a reverse transcription kit, and the target gene primers were synthesized by Shanghai Sangon Biotech using the following sequences (*Table 1*). PCR amplification was performed according to the manufacturer's instructions. The reaction system was 20  $\mu$ L, and the conditions were as follows: Pre-denaturation at 95 °C for 3 min, followed by 45 cycles of denaturation at 95 °C for 3 s and annealing and extension at 60 °C for 30 s. The amplification curve was analyzed after the reaction was ended. With the  $\beta$ -actin gene as an endogenous reference gene, the relative mRNA expression was calculated using the  $2^{-\Delta\Delta C_t}$  method. The experiment was repeated three times. The  $\Delta C_T$  value in the treated group = the average  $C_T$  value of the treated group – the  $C_T$  value of the endogenous control; the  $\Delta C_T$  value in the blank control group = the average  $C_T$  value of the blank control group—the  $C_T$  value of the endogenous control; and the  $\Delta\Delta C_t$  = the  $\Delta C_T$  value in the treated group—the  $\Delta C_T$  value in the blank control group;


- (II) Western blotting for determining the protein expressions of PTGDS, pAKT, and pGSK3 $\beta$  proceeded as follows. After sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE), the extracted total protein was transferred onto a PVDF membrane. After blocking with 1% bovine serum albumin (BSA) for 1.5 h, the proteins were mixed with the corresponding antibodies (1:1,000) and then inoculated overnight on a plate shaker in a cold room (4 °C); subsequently, the secondary antibodies were added before inoculation at room temperature for 1 h. A gel imaging system was used for imaging. The whole procedure was repeated three times.

#### ELISA for measuring the PGD2 and PGD2R levels in the supernatant of the cultured hDPCs


hDPCs were seeded in 6-well plates at  $2 \times 10^5$ /well, and each well had 2 mL of 15% DMEM-conditioned medium added. After the cells became adherent, they were cultured with serum-free DMEM containing levocetirizine hydrochloride 0 (control), 1, 10, 100, 1,000, or 10,000 ng/mL for 24 h. The culture media were collected and centrifuged at 1,000 rpm (radius arm: 15.0 cm) for 5 min. The supernatant was harvested, and the levels of PGD2 and PGD2R were determined according to the kit instructions. The whole procedure was repeated three times.

#### Statistical analysis

The statistical analysis was performed using the SPSS 17.0 software package. Measurement data are presented as  $\bar{x} \pm SD$ . Comparisons among multiple groups were performed



**Figure 1** Immunofluorescence observation of the effects of different concentrations of levocetirizine hydrochloride on the growth of human dermal papilla cells ( $\alpha$ -SMA staining 40 $\times$ 40).



**Figure 2** Effect of levocetirizine hydrochloride on the proliferation of hDPCs *in vitro* (\* $P < 0.05$ , compared with the blank control group). hDPCs, human dermal papilla cells.

using one-way analysis of variance (ANOVA), and the LSD-t test was used for pairwise comparisons. A  $P$  value of  $< 0.05$  was considered statistically significant.

## Results

### *hDPC growth*

The cells in the 100 ng/mL group grew well and arranged closely, with abundant cytoplasm and over 90% confluency. Cell confluency was about 70–80% in the blank control group. The cells in the 10,000 ng/mL group were sparsely distributed and loosely arranged, with a confluency of less than 60% (Figure 1).

### *Effect of levocetirizine hydrochloride on the proliferation of hDPCs*

The MTT results showed that the proliferation rate of hDPCs varied significantly across the different groups ( $F = 42.22$ ,  $P < 0.05$ ). It was significantly higher in the 10, 50, 100, 250, 500 ng/mL groups than in the blank control group (the  $t$  values were 5.31, 12.68, 28.26, 32.43, and 13.01, respectively; all  $P < 0.05$ ), whereas the 1 and 1,000 ng/mL groups showed no such significant difference when compared with the blank control group ( $t = 0.11$ ,  $t = 0.66$ , respectively; both  $P > 0.05$ ). The proliferation rate of hDPCs in the 10,000 and 100,000 ng/mL groups was significantly lower than that in the blank control group ( $t = 143.95$ ,  $t = 161.26$ , respectively; both  $P < 0.05$ ) (Figure 2).


### *Effects of levocetirizine hydrochloride on the mRNA expressions of COX-2, PGE2, PGF2 $\alpha$ , PTGDS, GPR44, AKT, and GSK3 $\beta$*

The mRNA expressions of COX-2, PGF2 $\alpha$ , PTGDS, GPR44, and AKT showed significant differences in different levocetirizine hydrochloride groups (all  $P < 0.05$ ); the mRNA expressions of COX-2, PTGDS, and GPR44 mRNA in the 100 ng/mL group ( $0.84 \pm 0.08$ ,  $0.81 \pm 0.10$ , and  $0.85 \pm 0.09$ , respectively) were significantly lower than those in the blank control group ( $t = 1.97$ ,  $t = 2.17$ , and  $t = 2.66$ , respectively; all  $P < 0.05$ ), whereas the mRNA expressions of PGF2 $\alpha$  and AKT in the 100 ng/mL group ( $1.96 \pm 0.25$  and  $1.74 \pm 0.32$ , respectively) were significantly higher than those in the blank control group

**Table 2** Effect of the different concentrations of levocetirizine hydrochloride on the mRNA expressions of COX-2, PGE2, PGF2 $\alpha$ , PTGDs, GPR44, AKT, and GSK3 $\beta$  in hDPCs (2<sup>- $\Delta\Delta C_t$</sup> )

Group	COX-2	PGE2	PGF2 $\alpha$	PTGDs	GPR44	AKT	GSK3 $\beta$
Blank control group	1	1	1	1	1	1	1
1	0.971 $\pm$ 0.087	1.094 $\pm$ 0.209	1.660 $\pm$ 0.225 <sup>a</sup>	0.965 $\pm$ 0.076	1.069 $\pm$ 0.125	1.348 $\pm$ 0.089 <sup>a</sup>	0.989 $\pm$ 0.082
10	0.902 $\pm$ 0.103	0.884 $\pm$ 0.095	1.646 $\pm$ 0.141 <sup>a</sup>	0.830 $\pm$ 0.089 <sup>a</sup>	0.977 $\pm$ 0.027	1.663 $\pm$ 0.191 <sup>a</sup>	0.928 $\pm$ 0.079
100	0.837 $\pm$ 0.081 <sup>a</sup>	0.956 $\pm$ 0.139	1.962 $\pm$ 0.248 <sup>a</sup>	0.809 $\pm$ 0.097 <sup>a</sup>	0.851 $\pm$ 0.086 <sup>a</sup>	1.738 $\pm$ 0.318 <sup>a</sup>	0.899 $\pm$ 0.060
1000	1.046 $\pm$ 0.107	1.271 $\pm$ 0.437	1.614 $\pm$ 0.158 <sup>a</sup>	0.859 $\pm$ 0.084 <sup>a</sup>	1.018 $\pm$ 0.050	1.600 $\pm$ 0.244 <sup>a</sup>	0.853 $\pm$ 0.094
10000	1.128 $\pm$ 0.092	1.119 $\pm$ 0.198	1.514 $\pm$ 0.592	0.988 $\pm$ 0.108	1.050 $\pm$ 0.081	1.777 $\pm$ 0.110 <sup>a</sup>	0.813 $\pm$ 0.086
F value	1.972	0.872	3.662	2.172	2.658	7.325	1.189
P value	0.028	0.399	0.034	0.045	0.042	0.002	0.371

n=3; <sup>a</sup>, P<0.05 compared with the blank control group. hDPCs, human dermal papilla cells.

**Figure 3** Effect of different concentrations of levocetirizine hydrochloride on the protein expression of PTGDS, pAKT and pGSK3 $\beta$  in hDPCs. hDPCs, human dermal papilla cells.

( $t=3.66$  and  $t=7.33$ , respectively; both  $P<0.05$ ). However, the mRNA expressions of PGE2 and GSK3 $\beta$  showed no significant difference between the different levocetirizine hydrochloride groups (all  $P>0.05$ ) (Table 2).

#### Effects of levocetirizine hydrochloride on the protein expressions of PTGDS, pAKT, and pGSK3 $\beta$ in hDPCs

There were significant differences in the levels of PTGDS, pAKT, and pGSK3 $\beta$  proteins among different levocetirizine hydrochloride groups (the F values were 11.84, 3.89, and 4.07, respectively;  $P<0.05$ ). The level of PTGDS protein in each concentration group was significantly lower than that in the blank control group (the  $t$  values were 11.58, 12.54, 5.67, 8.42, and 8.16, respectively;  $P<0.05$ ). The levels of

pAKT protein in the 100 and 1,000 ng/mL groups were significantly higher than those in the blank control group ( $t=16.60$ ,  $t=12.42$ , respectively); however, the pAKT protein levels in the 1, 10, and 10,000 ng/mL groups were not significantly different from those in the blank control group ( $t=0.28$ ,  $t=0.03$ , and  $t=0.22$ , respectively; all  $P>0.05$ ). The levels of pGSK3 $\beta$  protein in the 100, 1,000, and 10,000 ng/mL groups were significantly higher than those in the blank control group ( $t=7.73$ ,  $t=8.91$ , and  $t=5.94$ , respectively); however, the pGSK3 $\beta$  protein levels in the 1 and 10 ng/mL groups were not significantly different from those in the blank control group ( $t=0.004$ ,  $t=0.03$ , respectively; both  $P>0.05$ ) (Figure 3 and Table 3).

#### Effect of levocetirizine hydrochloride on the secretion of PGD2 and PGD2 receptors in hDPC

As shown by ELISA, the levels of PGD2 and PGD2R receptors in hDPC culture supernatants were significantly different across the different levocetirizine hydrochloride groups (all  $P<0.05$ ). The levels of PGD2 and PGD2R proteins in the 10–10,000 ng/mL groups were significantly lower than those in the blank control group (the  $t$  values were 20.75, 45.07, 87.09, 101.62, 46.72, 92.05, 90.67, and 83.16, respectively; all  $P<0.05$ ), although they were not significantly different between the 1 ng/mL group and the blank control group ( $t=0.26$ ,  $t=1.72$ , respectively; both  $P>0.05$ ) (Table 4).

## Discussion

DPCs, located in the dermal papilla at the base of the hair

**Table 3** PTGDS, pAKT, and pGSK3 $\beta$  protein expression levels after treatment with different concentrations of levocetirizine hydrochloride (based on the ratio of the gray value relative to ACTIN and GAPDH) ( $\bar{x} \pm SD$ )

Group	Blank control group	1	10	100	1,000	10,000
PTGDS protein	0.726 $\pm$ 0.058	0.503 $\pm$ 0.039 <sup>a</sup>	0.537 $\pm$ 0.089 <sup>a</sup>	0.322 $\pm$ 0.048 <sup>a</sup>	0.432 $\pm$ 0.051 <sup>a</sup>	0.437 $\pm$ 0.101 <sup>a</sup>
pAKT protein	0.457 $\pm$ 0.021	0.430 $\pm$ 0.084	0.447 $\pm$ 0.093	0.590 $\pm$ 0.052 <sup>a</sup>	0.585 $\pm$ 0.059 <sup>a</sup>	0.440 $\pm$ 0.059
pGSK3 $\beta$ protein	0.350 $\pm$ 0.042	0.348 $\pm$ 0.045	0.358 $\pm$ 0.068	0.457 $\pm$ 0.028 <sup>a</sup>	0.467 $\pm$ 0.044 <sup>a</sup>	0.436 $\pm$ 0.043 <sup>a</sup>

n=3; <sup>a</sup>, P<0.05 compared with the blank control group.

**Table 4** Effect of different concentrations of levocetirizine hydrochloride on the levels of PGD2 and PGD2R in hDPC culture supernatant ( $\bar{x} \pm SD$ , pg/mL)

Group	PGD2	PGD2R
Blank control group	180.08 $\pm$ 6.15	273.24 $\pm$ 3.18
1	178.15 $\pm$ 2.72	265.74 $\pm$ 7.42
10	158.92 $\pm$ 2.72 <sup>a</sup>	237.58 $\pm$ 7.96 <sup>a</sup>
100	141.62 $\pm$ 5.44 <sup>a</sup>	215.08 $\pm$ 9.55 <sup>a</sup>
1,000	126.23 $\pm$ 4.57 <sup>a</sup>	188.82 $\pm$ 12.20 <sup>a</sup>
10,000	124.31 $\pm$ 4.84 <sup>a</sup>	181.95 $\pm$ 13.79 <sup>a</sup>
F value	66.15	44.33
P value	<0.01	<0.01

n=3; <sup>a</sup>, P<0.05 compared with the blank control group. hDPC, human dermal papilla cells.

follicle, are a group of specially differentiated mesenchymal cells. Studies have confirmed that DPCs can induce the proliferation and differentiation of epithelial cells to form hair follicles, regulate hair growth cycle, and thus play an important role in the pathogenesis of AGA. It is now widely accepted that hair follicle stem cells are activated after receiving signals from the dermal papilla, which activates the hair follicles and triggers the hair to shift from the stationary phase into the growth phase (11). Abnormal androgen metabolic pathway is the primary triggering factor of AGA. Androgen receptors are mainly located in DPCs, further indicating that DPCs are the main target cells of androgen action on hair follicles (2). *In vitro* culture has shown that DPCs induce the proliferation and differentiation of epithelial cells through Wnt/ $\beta$ -catenin signaling pathway and promote the formation of hair follicles (12). In addition, DPCs can promote and maintain the growth and development of hair follicles by secreting various cytokines and growth factors; particularly, the signal

molecules including FGF, TGF- $\beta$ 1, IGF-1, VEGF, BMP, and DKK-1, which are produced by autocrine or paracrine, can interweave to form complex regulatory networks (13). Therefore, many studies have tended to use DPCs as an *in vitro* screening model to explore the mechanisms of hair growth regulators at the cellular and molecular levels.

No experiment has explored the effect of levocetirizine hydrochloride on hDPCs. In 2015, Wichitnithad *et al.* (14) found that, after healthy subjects were orally administered with levocetirizine hydrochloride at a therapeutic dose of 5 mg, the C<sub>max</sub> fluctuated at 196 $\pm$ 36 ng/mL, as determined by fast liquid chromatography-mass spectrometry. In the study performed by Jang *et al.* (15), the peak concentration of cetirizine in the epidermis of the skin reached 1.6–2.4 ng/mL after oral administration of 10 mg or 20 mg of cetirizine in healthy subjects. Petersen *et al.* (16) infected human nasal mucosal epithelial cells with human rhinovirus (HRV) to mimic a model of airway inflammation, which was treated with different concentrations of levocetirizine hydrochloride (molecular weight 388 g/mol); it was found that 19.4 ng/mL levocetirizine hydrochloride could effectively inhibit the expression of inflammatory cytokines including IL-6, IL-8, and NF- $\kappa$ B induced by HRV. We selected drug dilution concentrations using the above intervals, with the highest concentration being 0.01% levocetirizine hydrochloride (100,000 ng/mL). Both  $\alpha$ -SMA immunofluorescence staining and MTT assay showed that levocetirizine hydrochloride significantly promoted the proliferation of hDPCs, with good cell growth and a significantly increased proliferation rate at the concentration of 100 ng/mL. These findings suggest that levocetirizine hydrochloride might promote the growth and proliferation of hDPCs through the regulation of certain cell secreting factors, signaling molecules, and signal transduction pathways.

Many recent articles have described the role of prostaglandins in hair growth and development. Garza *et al.* (17) confirmed that the mRNA and protein expression

levels of both PTGDS and PGD2 increased in the alopecia area of AGA patients, particularly when the hair at the hair follicle was in a transition period between growth and regression. They also demonstrated that the topical use of PGD2 in mice transplanted with human hair follicles led to the miniaturization of hair follicles by inhibiting GPR44, resulting in hair thinning and shedding (18). PGD2 is an important inflammatory regulatory molecule that enhances or inhibits inflammatory responses by interacting with D prostanoid receptor 1 (PTGDR), D prostanoid receptor 2 (PGD2R), and TP receptors. In general, as a pro-inflammatory factor, PGD2 activates the DP2 receptor in human airway epithelial cells, stimulates massive mucus secretion by airway epithelial cells, causes bronchial muscle contraction, and leads to airway hyperresponsiveness (19). Recently, some studies have suggested that PGD2 may inhibit inflammatory response in certain environments. For instance, it can activate DP1 receptor to regulate and induce lung dendritic cells and regulatory T cells, and ultimately reduce asthma attacks. As a receptor for PGD2, GPR44 is expressed not only in the outer root sheath of the hair follicle but also in the dermal papilla of the outer sheath region (20-21). In addition, overexpression of COX-2 in mouse models also causes hair loss (22). Treatment of AGA using PTGDS inhibitors has been reported (23). These results suggest that prostaglandins are involved in hair growth and may be a new therapeutic target for alopecia.

In 1989, Charlesworth *et al.* found that cetirizine could inhibit inflammatory cell infiltration and while also significantly reducing the expression of PGD2 (24). Levocetirizine hydrochloride, by binding to COX-2, inhibits its ability to promote prostaglandin expression (7). Such effects are clearly independent of their antihistaminic activity. Recently, Jeong *et al.* (25) found that PGD2 activates androgen receptors by regulating DP2 and AKT signaling pathways on hDPCs, thereby stimulating the production of hair growth inhibitory factors such as TGF $\beta$ 1, Creb, LEF1, and IGF-1. AKT, also known as protein kinase B, is an important signal transduction molecule that regulates cell proliferation, differentiation, migration, and apoptosis. For one, the AKT signaling pathway promotes the proliferation of hair follicle stem cells and induces epidermal hyperplasia (26). For another, the activation of AKT signaling pathway activates multiple downstream target protein genes including GSK3 $\beta$ , Bcl-2 family members, caspase-9, and NF- $\kappa$ B, and thus participates in the proliferation and differentiation of DPCs and the growth of hair follicles. GSK3 $\beta$ , for example, is

both a downstream gene in the AKT signaling pathway and also a key factor in the classical Wnt/ $\beta$ -catenin signaling pathway (which is the first dermal signaling pathway).

In our current study, we used hDPCs to explore the relationship between levocetirizine hydrochloride and hair growth. Real-time quantitative PCR showed that levocetirizine hydrochloride 100 ng/mL significantly inhibited the mRNA expressions of COX-2, PTGDS, and GPR44 and promoted the mRNA expression of AKT. Western blotting and ELISA also showed that levocetirizine hydrochloride reduced the protein expression levels of PTGDS, PGD2, and PGD2 receptors while up-regulating the protein expression levels of pAKT protein and pGSK3 $\beta$ . In addition, comparisons of the effects of different concentrations of levocetirizine hydrochloride on hDPCs showed that low- and high-concentrations of levocetirizine hydrochloride were not conducive to the expressions of the above signaling molecules and pathway molecules. For example, levocetirizine hydrochloride 1 ng/mL could only down-regulate the protein expression of PTGDS in hDPCs, whereas levocetirizine hydrochloride 10,000 ng/mL significantly inhibited the protein expressions of PTGDS, PGD2, and PGD2R. Therefore, levocetirizine hydrochloride may promote the growth and proliferation of hDPC *in vitro* by inhibiting the PGD2-GPR44 pathway and activating the AKT signaling pathway.

In addition, we found that levocetirizine hydrochloride promoted the expression of PGF2 $\alpha$  but did not affect the expression of PGE2. Although both of them are prostaglandins, their effects on hair growth are exactly the opposite of those of PGD2. In 2009, the US Food and Drug Administration (FDA) approved the clinical use of Latanoprost, a PGF2 $\alpha$  analogue, to promote the growth of human eyelashes and eyebrows (27). It is believed that PGF2 $\alpha$  and its analogues promote the proliferation of hair follicle epithelial cells mainly by relaxing the small arteries and thus increasing local blood supply (28). PGE2 has also been shown to improve hair loss induced by radiation in mouse models (29). At present, no experiment has investigated the downstream signaling pathway of AKT and the regulation mechanism of PGF2 $\alpha$ . Moreover, the mechanism via which levocetirizine hydrochloride regulates PDC2-GPR44 in living bodies also warrants further study.

In conclusion, our experiment demonstrated that levocetirizine hydrochloride could promote the growth and proliferation of hDPCs and significantly inhibit the expressions of relevant factors including COX-2, PTGDS, PGD2, PGD2R, and GPR44 in the PGD2-GPR44



pathway. The role of AKT signaling pathway in this process was also explored. These findings provide an experimental and theoretical basis for the treatment of AGA with levocetirizine hydrochloride.

### Acknowledgments

*Funding:* This study was supported by the National Natural Science Foundation of China (81502744) and the Guangxi Natural Science Foundation (2015GXNSFB139131).

### Footnote

*Conflicts of Interest:* The authors have no conflicts of interest to declare.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

### References

1. Yamazaki M, Miyakura T, Uchiyama M, et al. Oral finasteride improved the quality of life of androgenetic alopecia patients. *J Dermatol* 2011;38:773-7.
2. Urysiak-Czubatka I, Kmiec ML, Broniarczyk-Dyla G. Assessment of the usefulness of dihydrotestosterone in the diagnostics of patients with androgenetic alopecia. *Postepy Dermatol Alergol* 2014;31:207-15.
3. Zhang Y. A study on the difference between normal hair follicle and pathological hair follicle in males with alopecia. Second Military Medical University, 2016.
4. Ghanaat M. Types of hair loss and treatment options, including the novel low-level light therapy and its proposed mechanism. *South Med J* 2010;103:917-21.
5. Kanei A, Asano K, Kanai K, et al. Inhibitory action of levocetirizine on the production of eosinophil chemoattractants RANTES and eotaxin *in vitro* and *in vivo*. *In Vivo* 2014;28:657-66.
6. Rossi A, Campo D, Fortuna MC, et al. A preliminary study on topical cetirizine in the therapeutic management of androgenetic alopecia. *J Dermatolog Treat* 2018;29:149-151.
7. Abo Dena AS, Abdel Gaber SA. *In vitro* drug interaction of levocetirizine and diclofenac: Theoretical and spectroscopic studies. *Spectrochim Acta A Mol Biomol Spectrosc* 2017;181:239-48.
8. Arias-Santiago S, Arrabal-Polo MA, Buendía-Eisman A, et al. Androgenetic alopecia as an early marker of benign prostatic hyperplasia. *J Am Acad Dermatol* 2012;66:401-8.
9. Heilmann S, Nyholt DR, Brockschmidt FF, et al. No genetic support for a contribution of prostaglandins to the aetiology of androgenetic alopecia. *Br J Dermatol* 2013;169:222-4.
10. Jeong KH, Jung JH, Kim JE, et al. Prostaglandin D2-Mediated DP2 and AKT Signal Regulate the Activation of Androgen Receptors in Human Dermal Papilla Cells. *Int J Mol Sci* 2018. doi: 10.3390/ijms19020556.
11. Greco V, Chen T, Rendl M, et al. A two-step mechanism for stem cell activation during hair regeneration. *Cell Stem Cell* 2009;4:155-69.
12. Kwack MH, Kim MK, Kim JC, et al. Wnt5a attenuates Wnt/ $\beta$ -catenin signalling in human dermal papilla cells. *Exp Dermatol* 2013;22:229-31.
13. Madaan A, Verma R, Singh AT, et al. Review of Hair Follicle Dermal Papilla cells as *in vitro* screening model for hair growth. *Int J Cosmet Sci* 2018;40:429-50.
14. Wichitnithad W, Jithavech P, Sanphanya K, et al. Determination of levocetirizine in human plasma by LC-MS-MS: validation and application in a pharmacokinetic study. *J Chromatogr Sci* 2015;53:1663-72.
15. Jang YJ, Wang JH, Kim JS, et al. Levocetirizine inhibits rhinovirus-induced ICAM-1 and cytokine expression and viral replication in airway epithelial cells. *Antiviral Res* 2009;81:226-33.
16. Petersen LJ, Church MK, Rihoux JP, et al. Measurement of interstitial cetirizine concentrations in human skin: correlation of drug levels with inhibition of histamine-induced skin responses. *Allergy* 1999;54:607-11.
17. Garza LA, Liu Y, Yang Z, et al. Prostaglandin D2 inhibits hair growth and is elevated in bald scalp of men with androgenetic alopecia. *Sci Transl Med* 2012;4:126ra34.
18. Nieves A, Garza LA. Does prostaglandin D2 hold the cure to male pattern baldness?. *Exp Dermatol* 2014;23:224-7.
19. Chen LX, Ran DH, Huang HP, et al. Prostaglandin D2

- stimulates human airway mucus hypersecretion mediated by DP2. *Basic & Clinical Medicine* 2016;36:24-9.
20. Colombe L, Vindrios A, Michelet JF, et al. Prostaglandin metabolism in human hair follicle. *Exp Dermatol* 2007;16:762-9.
  21. Colombe L, Michelet JF, Bernard BA. Prostanoid receptors in anagen human hair follicles. *Exp Dermatol* 2008;17:63-72.
  22. Bol DK, Rowley RB, Ho CP, et al. Cyclooxygenase-2 overexpression in the skin of transgenic mice results in suppression of tumor development. *Cancer Res* 2002;62:2516-21.
  23. Fong P, Tong HH, Ng KH, et al. In silico prediction of prostaglandin D2 synthase inhibitors from herbal constituents for the treatment of hair loss. *J Ethnopharmacol* 2015;175:470-80. .
  24. Charlesworth EN, Kagey-Sobotka A, Norman PS, et al. Effect of cetirizine on mast cell-mediator release and cellular traffic during the cutaneous late-phase reaction. *J Allergy Clin Immunol* 1989;83:905-12.
  25. Jeong KH, Jung JH, Kim JE, et al. Prostaglandin D2-Mediated DP2 and AKT Signal Regulate the Activation of Androgen Receptors in Human Dermal Papilla Cells. *Int J Mol Sci* 2018. doi: 10.3390/ijms19020556.
  26. Ohnemus U, Uenal M, Inzunza J, et al. The hair follicle as an estrogen target and source. *Endocr Rev* 2006;27:677-706.
  27. Faghihi G, Andalib F, Asilian A. The efficacy of latanoprost in the treatment of alopecia areata of eyelashes and eyebrows. *Eur J Dermatol* 2009;19:586-7.
  28. Barrón-Hernández YL, Tosti A. Bimatoprost for the treatment of eyelash, eyebrow and scalp alopecia. *Expert Opin Investig Drugs* 2017;26:515-22.
  29. Geng L, Hanson WR, Malkinson FD. Topical or systemic 16, 16 dm prostaglandin E2 or WR-2721 (WR-1065) protects mice from alopecia after fractionated irradiation. *Int J Radiat Biol* 1992;61:533-7.

**Cite this article as:** Wen S, Wei J, Bao J, Guo T, Zheng W, Zhuang X, Lin Y. Effect of levocetirizine hydrochloride on the growth of human dermal papilla cells: a preliminary study. *Ann Palliat Med* 2020;9(2):308-317. doi: 10.21037/apm.2020.01.15