



Utility of intraparenchymal blood patch testing following CT-guided lung biopsy

Khezar Syed¹^, Bryan S. Benn², Jonathan S. Kurman¹

¹Division of Pulmonary & Critical Care, Medical College of Wisconsin, Milwaukee, WI, USA; ²Department of Pulmonary Medicine, Respiratory Institute, Cleveland Clinic, Cleveland, OH, USA

Correspondence to: Jonathan S. Kurman, MD, MBA. Assistant Professor of Medicine, Director of Interventional Pulmonology, Division of Pulmonary & Critical Care, Medical College of Wisconsin, 8701 W Watertown Plank Rd., Milwaukee, WI 53226, USA. Email: jkurman@mcw.edu.

Comment on: Jain N, Zurla Z, Crowley S, *et al.* Tract injection of autologous blood (intraparenchymal blood patching) in percutaneous transthoracic CT-guided lung biopsy and the incidence of pneumothorax: a retrospective analysis. *Precis Cancer Med* 2022;5:34.

Keywords: Intraparenchymal blood patching (IBP); pneumothorax; CT-guided biopsy

Received: 14 November 2022; Accepted: 01 December 2022; Published online: 29 December 2022.

doi: 10.21037/pcm-22-62

View this article at: <https://dx.doi.org/10.21037/pcm-22-62>

Lung cancer remains the leading cause of cancer-related deaths among men and women in the United States (1,2), with lung nodule prevalence ranging from 14–36% (3). With improved access to computed tomography (CT) scanners and the implementation of low-dose CT lung cancer screening in the United States and Europe, lung nodule detection rates will likely increase. Despite technological advances and the development of surveillance guidelines, confidence in our imaging technology's discriminatory capability to differentiate intermediate-risk lesions is still lacking (3,4). As advances in oncological treatments continue to evolve and therapies become increasingly tailored to specific mutations, tissue biopsies will likely continue to play a significant role in allowing precision treatments to be designed in the future (5).

Multiple options exist to obtain tissue for histological and molecular analysis. These include image-guided transthoracic biopsies, bronchoscopy-guided biopsies, and surgical biopsies. The chosen method depends on multiple factors, including location, lesion size, available expertise, and underlying comorbid conditions (6). A United States-based proprietary claims database review reveals that most patients undergo CT-guided needle biopsy (CTNB) for peripheral pulmonary

nodules (7-9). CTNB has excellent specificity and accuracy, with literature reporting pooled diagnostic accuracy of 92.1% (9,567/10,383), a sensitivity of 92.1% (7,343/7,975), and a specificity close to 100% for the diagnosis of malignancy (7,10). However, despite the excellent accuracy of CTNB, risks exist, including, most commonly, pneumothorax. Multiple studies put the incidence of pneumothorax at 15–62% (4,6,7,10,11). Dr. Jain and colleagues explore an intervention to mitigate the incidence risk of pneumothorax in their article (12).

The authors performed a retrospective analysis of all CTNB lung nodule biopsies carried out at two hospitals between January 2013 and January 2020. Of the 848 biopsies performed, 536 met the inclusion criteria. Cases were excluded if a chest tube was already in place, an alternative sealant material was used, or the guiding needle did not traverse the lung parenchyma to reach the sampled lesion. The control group consisted of 227 patients who underwent CTNB between 2013–2015. A total of 309 patients who had a CTNB from 2016 onwards received an intraparenchymal blood patching (IBP), which utilizes 10–15 mL of the patient's blood obtained while securing intravenous access. The blood was then instilled into the biopsy tract as the introducer needle was removed.

[^] ORCID: 0000-0002-1054-5245.

The primary endpoints examined were post-biopsy pneumothorax, the need for chest tube placement, and the requirement of hospital admission. Thirty-seven percent of patients (84/227) who did not receive IBP developed a pneumothorax post-biopsy, while only 14.6% (45/309) of patients who received an IBP developed a pneumothorax. The difference was statistically significant ($P < 0.0001$), with an absolute risk reduction of 22.4% and a number needed to treat (NNT) of 4.4. In addition, there was a significant decrease in the incidence of chest tube placement from 17.2% (39/227) in the control group to 3.9% (12/309) ($P < 0.0001$) among patients who received IBP, with an absolute risk reduction of 13.3%. Similarly, there was a significant decrease in hospital admissions from 16.3% (37/227) to 2.9% (9/309) in patients who received an IBP ($P < 0.0001$).

Caveats exist when interpreting the results of this study. First, there was a statistical difference between the prevalence of patients with underlying chronic obstructive pulmonary disease (COPD) in the two groups (46.7% in the control arm *vs.* 35.4% in the treatment arm, $P = 0.008$). Previous studies have shown COPD to be a risk factor for developing pneumothorax post-biopsy (7,13). Second, the two groups differed in home oxygen usage (7.1% in the control arm *vs.* 1.9% in the treatment arm, $P = 0.003$). Although this finding may reflect the difference between the prevalence of COPD in the two groups, it may also indicate a difference in COPD severity between the groups. Third, one physician performed all the biopsies with IBP, while multiple physicians may have performed the earlier biopsies. Thus, the difference in pneumothorax rate may reflect differences in expertise and technique. Finally, as this study was a retrospective analysis, readers should be cognizant of biases associated with retrospective analyses, namely selection bias and confounding bias. Multivariate analysis could have mitigated potentially confounding bias by controlling for underlying pulmonary disease, severity, and the operator's expertise level. Without that further analysis, however, the observed effect may be due to other variables, such as technique, patient rollover, needle removal during expiration, patient factors, or keeping the biopsied lung dependent (14). Unfortunately, these variables were also not uniform across all cases, and further details are unavailable.

Jain *et al.*'s (12) findings highlight the need for updated guidelines governing CTNB. Image-guided percutaneous lung biopsies have been performed since 1960 (13). Pneumothorax and bleeding were the most frequent complications initially. In 1974, McCartney *et al.*

described their experience of 25 cases where autologous blood patching or gel foam was used to seal the needle track, preventing pneumothoraces (15). Two subsequent studies, however, failed to show any statistical difference in pneumothorax rate with IBP (16,17). The quest to decrease the incidence of post-biopsy pneumothorax continued. After lab models reaffirmed the utility of blood patching (18), Lang *et al.* revisited IBP with a multicenter randomized control trial (RCT), reaffirming its benefits (19). Despite encouraging results, the only formal guideline dedicated to CTNB did not endorse IBP and questioned its utility (20).

Over the years, many other materials have been trialed to help seal the needle track, including normal saline, compressed collagen foam plugs, fibrin glue, and absorbable hemostat gelatin powder (21-24). In addition, many maneuvers, such as rapid rollover (puncture site down) and breath-hold after expiration during needle extraction, have been developed and trialed to decrease the incidence of pneumothorax, chest tube insertion, and hospital admission (14,20). Even without formal endorsement from any specialty society, many physicians have been using IBP and other maneuvers to help decrease pneumothorax rates following CTNB (21,25). Huo *et al.* systematically reviewed post-biopsy maneuvers from 21 articles, totaling more than 7,000 patients (21). The maneuvers included IBP, needle tract plugs, changes in patient position, and needle extraction during exhalation. The pooled odds in six studies that evaluated IBP (including four RCTs) reduced the risk of pneumothorax by half and chest tube insertion by three-fold, indicating that efforts to reduce the incidence of complications appear effective.

Multiple studies have examined the economic burden associated with diagnostic testing of lung nodules and the substantial cost incurred after complications, which can be as much as four times the cost of the actual procedure (8,9,11). For example, Huo *et al.* revealed that the additional cost of minor to intermediate complications, such as pneumothorax without or with a chest tube, ranges from \$5,881 to \$20,457 (9). Thus, efforts to decrease the risk of complications are certainly needed.

As the era of precision medicine continues to evolve, the biopsy of pulmonary nodules will continue to play an integral role in establishing a diagnosis and developing a treatment plan. CTNB will remain a mainstay of lung nodule evaluation for the foreseeable future. Unfortunately, pneumothorax will also likely remain the most common complication of this procedure. IBP appears to be a safe procedure when performed after CTNB to seal the needle

tract and decrease the incidence of pneumothorax. The reproducible decrease in pneumothorax rate with IBP seen in numerous trials lends support for its routine use following the withdrawal of the introducer needle after CTNB. Incorporating IBP into future guidelines for the performance of CTNB should be considered.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Precision Cancer Medicine*. The article did not undergo external peer review.

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://pcm.amegroups.com/article/view/10.21037/pcm-22-62/coif>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- American Cancer Society. Cancer Facts & Figures 2022. Atlanta: American Cancer Society; 2022. Accessed November 5, 2022. Available online: <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/annual-cancer-facts-and-figures/2022/2022-cancer-facts-and-figures.pdf>
- Cronin KA, Scott S, Firth AU, et al. Annual report to the nation on the status of cancer, part 1: National cancer statistics. *Cancer* 2022;128:4251-84.
- Nair A, Devaraj A, Callister MEJ, et al. The Fleischner Society 2017 and British Thoracic Society 2015 guidelines for managing pulmonary nodules: keep calm and carry on. *Thorax* 2018;73:806-12.
- Chiu YW, Kao YH, Simoff MJ, et al. Costs of Biopsy and Complications in Patients with Lung Cancer. *Clinicoecon Outcomes Res* 2021;13:191-200.
- Marshall D, Laberge JM, Firetag B, et al. The changing face of percutaneous image-guided biopsy: molecular profiling and genomic analysis in current practice. *J Vasc Interv Radiol* 2013;24:1094-103.
- Gould MK, Donington J, Lynch WR, et al. Evaluation of individuals with pulmonary nodules: when is it lung cancer? Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2013;143:e93S-e120S.
- Vachani A, Zhou M, Ghosh S, et al. Complications After Transthoracic Needle Biopsy of Pulmonary Nodules: A Population-Level Retrospective Cohort Analysis. *J Am Coll Radiol* 2022;19:1121-9.
- Zheng F, Lavin J, Sprafka JM. Patient out-of-pocket costs for suspicious pulmonary nodule biopsy in lung cancer patients. *J Med Econ* 2021;24:1173-7.
- Huo J, Xu Y, Sheu T, et al. Complication Rates and Downstream Medical Costs Associated With Invasive Diagnostic Procedures for Lung Abnormalities in the Community Setting. *JAMA Intern Med* 2019;179:324-32.
- DiBardino DM, Yarmus LB, Semaan RW. Transthoracic needle biopsy of the lung. *J Thorac Dis* 2015;7:S304-16.
- Zlevor AM, Mauch SC, Knott EA, et al. Percutaneous Lung Biopsy with Pleural and Parenchymal Blood Patching: Results and Complications from 1,112 Core Biopsies. *J Vasc Interv Radiol* 2021;32:1319-27.
- Jain N, Zurla Z, Crowley S, et al. Tract injection of autologous blood (intraparenchymal blood patching) in percutaneous transthoracic CT-guided lung biopsy and the incidence of pneumothorax: a retrospective analysis. *Precis Cancer Med* 2022;5:34.
- Lalli AF, Naylor B, Whitehouse WM. Aspiration biopsy of thoracic lesions. *Thorax* 1967;22:404-7.
- Najafi A, Al Ahmar M, Bonnet B, et al. The PEARL Approach for CT-guided Lung Biopsy: Assessment of Complication Rate. *Radiology* 2022;302:473-80.
- McCartney R, Tait D, Stilson M, et al. A technique for the prevention of pneumothorax in pulmonary aspiration biopsy. *Am J Roentgenol Radium Ther Nucl Med* 1974;120:872-5.
- Bourgouin PM, Shepard JA, McCloud TC, et al.

- Transthoracic needle aspiration biopsy: evaluation of the blood patch technique. *Radiology* 1988;166:93-5.
17. Herman SJ, Weisbrod GL. Usefulness of the blood patch technique after transthoracic needle aspiration biopsy. *Radiology* 1990;176:395-7.
 18. Moore EH, Shelton DK, Wisner ER, et al. Needle aspiration lung biopsy: reevaluation of the blood patch technique in an equine model. *Radiology* 1995;196:183-6.
 19. Lang EK, Ghavami R, Schreiner VC, et al. Autologous blood clot seal to prevent pneumothorax at CT-guided lung biopsy. *Radiology* 2000;216:93-6.
 20. Manhire A, Charig M, Clelland C, et al. Guidelines for radiologically guided lung biopsy. *Thorax* 2003;58:920-36.
 21. Huo YR, Chan MV, Habib AR, et al. Post-Biopsy Manoeuvres to Reduce Pneumothorax Incidence in CT-Guided Transthoracic Lung Biopsies: A Systematic Review and Meta-analysis. *Cardiovasc Intervent Radiol* 2019;42:1062-72.
 22. Billich C, Muche R, Brenner G, et al. CT-guided lung biopsy: incidence of pneumothorax after instillation of NaCl into the biopsy track. *Eur Radiol* 2008;18:1146-52.
 23. Zaetta JM, Licht MO, Fisher JS, et al. A lung biopsy tract plug for reduction of postbiopsy pneumothorax and other complications: results of a prospective, multicenter, randomized, controlled clinical study. *J Vasc Interv Radiol* 2010;21:1235-43.e1-3.
 24. Maybody M, Muallem N, Brown KT, et al. Autologous Blood Patch Injection versus Hydrogel Plug in CT-guided Lung Biopsy: A Prospective Randomized Trial. *Radiology* 2019;290:547-54.
 25. Graffy P, Loomis SB, Pickhardt PJ, et al. Pulmonary Intraparenchymal Blood Patching Decreases the Rate of Pneumothorax-Related Complications following Percutaneous CT-Guided Needle Biopsy. *J Vasc Interv Radiol* 2017;28:608-613.e1.

doi: 10.21037/pcm-22-62

Cite this article as: Syed K, Benn BS, Kurman JS. Utility of intraparenchymal blood patch testing following CT-guided lung biopsy. *Precis Cancer Med* 2023;6:1.