



Potential of ketamine use in sialendoscopy—a narrative review

Nancy Zhu¹, Tim T. Wang¹, Jingping Wang²

¹Department Oral & Maxillofacial Surgery, Massachusetts General Hospital and Clinical Fellow, Harvard School of Dental Medicine, Boston, MA, USA; ²Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital, Boston, MA, USA

Contributions: (I) Conception and design: N Zhu, J Wang; (II) Administrative support: None; (III) Provision of study materials or patients: None; (IV) Collection and assembly of data: All authors; (V) Data analysis and interpretation: All authors; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Jingping Wang, MD, PhD. Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital, 55 Fruit Street, GRB 444, Boston, MA 02114, USA. Email: jwang23@mgh.harvard.edu.

Background and Objective: Sialendoscopy is a procedure that is commonly used nowadays to both diagnose and treat salivary gland stones. Ketamine is a well-studied anesthetic that is known to induce hyper-salivation. We aim to review the current evidence surrounding possible synergistic usage of ketamine in sialendoscopy procedures as a sialagogue, anesthetic and analgesic agent.

Methods: A literature search was completed on PubMed, Embase, and CINAHL online databases using keywords “ketamine” and “sialendoscopy”.

Key Content and Findings: The literature search did not reveal any studies that directly studied the possibility of using ketamine in sialendoscopy procedures. However, many articles were found on the diverse uses and the hyper-salivation side effect of ketamine as well as on the need for agents to increase success of identifying the orifice of salivary ducts during sialendoscopy procedures.

Conclusions: Ketamine is a promising anesthetic agent to use in sialendoscopy procedures over other agents given its analgesic and sialorrhea properties. Additionally, ketamine has a well-documented broad safety profile that makes it a favorable anesthetic for short procedures that would be beneficial in short procedures such as sialendoscopy. Future studies evaluating the safety and efficacy of ketamine in sialendoscopy procedures are needed to provide evidence that ketamine reduces need for opioid analgesics and increases procedural success due to hyper-salivation side effect.

Keywords: Opioid-free analgesic; sialorrhea; ketamine; sialendoscopy

Received: 09 September 2022; Accepted: 08 February 2023; Published online: 14 February 2023.

doi: 10.21037/joma-22-28

View this article at: <https://dx.doi.org/10.21037/joma-22-28>

Introduction

Background

Ketamine was first introduced as a human anesthetic in the 1960s by Corssen and Domino, and is now one of the most widely used anesthetic agent to this day (1). Ketamine is a noncompetitive antagonist at the N-methyl-D-aspartate (NMDA) receptor. The widely available form used today for anesthesia is a racemic mixture of two isomers, R (–) and S (+). The S (+) form binds more strongly to NMDA receptors and is largely the driving isomer for the anesthetic effects of ketamine, while the R (–) form has been shown to

elicit a sustained antidepressant effect (2).

Ketamine is often favored as an anesthetic option because it has a relatively stable effect on systemic vascular resistance. It also does not cause significant respiratory depression and preserves upper airway reflexes during anesthesia (1). Additionally, ketamine has been found to induce sialorrhea, or hypersalivation. While this raised some concerns about potentially increased risk of laryngospasms, these concerns have yet to be validated by scientific evidence (3). It is not entirely clear whether this side effect is dose dependent, as current literature demonstrates conflicting evidence (4,5). Nevertheless, the

Table 1 The search strategy summary

Items	Specification
Date of search	August 1, 2022–August 29, 2022
Databases and other sources searched	PubMed, Embase, CINAHL
Search terms used	Keywords: ketamine, sialendoscopy
Timeframe	No limitations
Inclusion and exclusion criteria	No language or study type restrictions
Selection process	No restrictions

effect of sialorrhea raises the possibility of synergistic use of ketamine in sialendoscopy.

Sialendoscopy, a minimally invasive technique for salivary gland surgery, was first proposed by Katz in 1991 (6). With subsequent improvement in equipment and techniques, sialendoscopy is increasingly used to diagnose and treat salivary gland stones, known as sialoliths. Sialoliths can form in major (i.e., parotid, submandibular or sublingual) or minor salivary glands. The cause of salivary gland stones is not fully understood, though saliva composition, duct anatomy, foreign bodies are thought to contribute to their formation (7). Most commonly, these stones are formed in the submandibular gland because the saliva there has a higher concentration of calcium and has a more alkaline pH than other salivary glands (8,9). Additionally, the route of the submandibular gland ducts is the most tortuous—running from the angle of the mandible to the base of the tongue—which predisposes this gland to relative stasis and therefore increased risk of stone formation.

Rationale and knowledge gap

Oral and maxillofacial surgeons (OMSs) frequently employ sialendoscopy to treat salivary gland pathologies in patients who fail conservative therapies (i.e., hydration, sialagogue usage, gland massage). These procedures can be done either in the clinic under IV sedation or in the operating room under general anesthesia (10). In the following, we examine the current evidence regarding the usage of ketamine in sialendoscopy. To our knowledge, there are no currently published studies to this end.

Objective

We aim to evaluate the evidence around the hyper-salivatory and perioperative analgesic effects of ketamine, and how

they might be synergistically applied for sialendoscopy. We present this article in accordance with the Narrative Review reporting checklist (available at <https://joma.amegroups.com/article/view/10.21037/joma-22-28/rc>).

Methods

A literature search was completed on PubMed, Embase, and CINAHL online databases using keywords “ketamine” and “sialendoscopy” in August 2022 (*Table 1*). Our inclusion criteria were studies of any type that provided evidence for the use of ketamine over other anesthetic agents in sialendoscopy procedures. No language or date limitations were set. While no articles that directly studied the use of ketamine in sialendoscopy were found, we did identify a number of studies regarding the diverse uses of ketamine in various clinical settings, as well as a number of studies that leveraged sialagogues in sialendoscopy. These findings further promote the synergistic potential of ketamine use in sialendoscopy.

Discussion

Sialorrhea in sialendoscopy

Ketamine has long been known to cause sialorrhea (1). Often, atropine is given intra-operatively in conjunction with ketamine to mitigate this side effect (5). The mechanism of action of sialorrhea is unclear. M1 and M3 are the known major acetylcholine receptors in the salivary gland. A study demonstrated decreased salivation in mice with both M1 and M3 knockouts (11). Clinically, patients with dry mouth syndrome are treated with cholinomimetics such as pilocarpine and carbachol. Ketamine, on the other hand, has been shown to inhibit muscarinic acetylcholine receptors (12). As such, one might expect ketamine to

reduce rather than increase salivary flow. As such, this is an area that is poorly understood and an area that requires further research.

However, the hyper-salivatory effect of ketamine may be helpful in sialendoscopy, particularly for the first part of the procedure. Sialendoscopy involves two distinct phases: salivary duct access and intraductal exploration (13). Access is thought to be the most technique-sensitive and time-consuming portion of the procedure due to the delicate structure and variable location of the salivary gland papilla (14,15). Current techniques include use of guidewire to locate the duct followed by serial dilation that slide over the guidewire, addition of viscous lidocaine infiltration in the duct for hydro-dilatory effects, or even angiocath-guided insertion of the endoscope (16,17). Still, identifying the orifice remains a challenge. The ductal orifice is the narrowest aspect of the duct, measuring as small as 0.1–0.2 mm, for both the submandibular and parotid gland (14,18). In this setting, ketamine could be helpful as a sialagogue. One study demonstrated increased visibility of the submandibular duct from 63% to 95% with the application of oral ascorbic acid (vitamin C) and another with lemon juice (19,20). Even if the duct in question is completely obstructed, there may be some benefit in using sialagogues to better identify the contralateral duct as a reference. Once the duct is accessed, it is explored for pathology. At any point, if hypersalivation needs to be reduced, glycopyrrolate can be given to the patient to minimize secretions.

Perioperative analgesic effects of ketamine

Ketamine has a noticeable analgesic action at small doses (defined as less than 1 mg/kg when administered intravenously) (21,22). Ketamine does so through both analgesic and anti-hyperalgesia properties at the NMDA receptor level (23). Additionally, ketamine has been shown to be more effective if given before noxious stimulus (24). This effect is referred to as “preemptive analgesia”, meaning that the memory of pain is reduced, thereby also decreasing the need for post-operative pain medications. As such, ketamine given prior to sialendoscopy procedures may be helpful in supplementing intraoperative local anesthesia and reducing post-operative pain (25-27).

Currently, there are no widely used guidelines within OMS regarding post-sialendoscopy pain control. Studies suggest that this procedure is generally well tolerated by patients and that patients can typically resume daily life

immediately post-operatively (28,29). Pain control regimens may consist of mostly over-the-counter analgesics, such as non-steroidal anti-inflammatory drugs and acetaminophen. Nevertheless, opioids may also be prescribed by some OMSs for patients who are more sensitive or on a “just-in-case” basis (30). After all, OMSs commonly prescribe short courses of opioids after other office-based procedures, such as third molar extraction and orthognathic surgery (31,32). To this end, ketamine may be particularly helpful in helping mitigate the post-operative pain level and possibly reduce the need for opioids.

Safety of ketamine for OMS

Ketamine is a well-studied drug that has been shown to be able to provide adequate sedation without compromising the airway (1,33). It is a common agent used in procedural sedation in OMS offices and also the emergency department for this reason (34-36). Regarding sialendoscopy specifically, completing the procedure in the office setting and the operating room had no significant differences in clinical outcomes (10,37,38). Furthermore, a recent study found that the mean total time burden of in-office sialendoscopy was 39 versus 278 minutes for the OR, corresponding to an average total procedure charge of \$720 versus \$13,956 (37). As such, when clinically feasible, in-office IV sedation for sialendoscopy should be considered.

Limitations and future study

While the usage of ketamine in sialendoscopy is promising, there remain limitations that should be considered. Ketamine has other known side effects, such as psychiatric disturbances in mood state and body image, floating sensations, vivid dreams or illusions, and occasional frank delirium during emergence. These side effects can be minimized by the administration of a benzodiazepine (39). There are also patients for whom ketamine use is contraindicated, such as those with renal or hepatic dysfunction (40).

Nevertheless, ketamine remains a promising choice that warrants investigation for safety and efficacy of usage in sialendoscopy. Studies can evaluate the effect of ketamine use on OMSs perceptions of procedural difficulty that are then corroborated with procedural duration. On the patient side, studies can evaluate the effect of ketamine use on post-operative pain levels in sialendoscopy. Further, these studies may elucidate other potential complications specific to sialendoscopy, persistent sialorrhea after

identifying the duct of interest or poor patient experience due to psychiatric side effects of ketamine. Ultimately, it will be helpful to build standardized guidelines across OMS regarding the benefits and drawbacks of employing ketamine in sialendoscopy.

Conclusions

Ketamine's effects of hypersalivation and analgesia, along with its broad safety profile, make it a potentially synergistic anesthetic to use in sialendoscopy. Possible benefits range from shortening procedure times to improving patient pain levels. As such, future studies evaluating the safety and efficacy of ketamine use in sialendoscopy are strongly indicated and can be influential in shaping future standards of practice.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Journal of Oral and Maxillofacial Anesthesia*, for the series "Opioid-free Anesthesia and Opioid-sparing Anesthesia in OMF Surgery". The article has undergone external peer review.

Reporting Checklist: The authors have completed the Narrative Review reporting checklist. Available at <https://joma.amegroups.com/article/view/10.21037/joma-22-28/rc>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://joma.amegroups.com/article/view/10.21037/joma-22-28/coif>). The series "Opioid-free Anesthesia and Opioid-sparing Anesthesia in OMF Surgery" was commissioned by the editorial office without any funding or sponsorship. JW serves as the unpaid editorial board member of *Journal of Oral and Maxillofacial Anesthesia* from August 2021 to July 2023 and served as unpaid Guest Editor of the series. The authors have no other conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Corssen G, Domino EF. Dissociative anesthesia: further pharmacologic studies and first clinical experience with the phencyclidine derivative CI-581. *Anesth Analg* 1966;45:29-40.
2. Yang C, Shirayama Y, Zhang JC, et al. R-ketamine: a rapid-onset and sustained antidepressant without psychotomimetic side effects. *Transl Psychiatry* 2015;5:e632.
3. Green SM, Roback MG, Krauss B, et al. Laryngospasm during emergency department ketamine sedation: a case-control study. *Pediatr Emerg Care* 2010;26:798-802.
4. McGlone RG, Howes MC, Joshi M. The Lancaster experience of 2.0 to 2.5 mg/kg intramuscular ketamine for paediatric sedation: 501 cases and analysis. *Emerg Med J* 2004;21:290-5.
5. Heinz P, Geelhoed GC, Wee C, et al. Is atropine needed with ketamine sedation? A prospective, randomised, double blind study. *Emerg Med J* 2006;23:206-9.
6. Katz P. Endoscopy of the salivary glands. *Ann Radiol (Paris)* 1991;34:110-3.
7. Hammett JT, Walker C. Sialolithiasis. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 [cited 2022 Aug 31]. Available online: <http://www.ncbi.nlm.nih.gov/books/NBK549845/>
8. Lustmann J, Regev E, Melamed Y. Sialolithiasis. A survey on 245 patients and a review of the literature. *Int J Oral Maxillofac Surg* 1990;19:135-8.
9. Miloro M, Ghali GE, Larsen PE, et al. editors. *Peterson's Principles of Oral and Maxillofacial Surgery*. 4th ed. 2022 edition. Cham: Springer, 2022:2350.
10. Bawazeer N, Carvalho J, Djennaoui I, et al. Sialendoscopy under conscious sedation versus general anesthesia. A comparative study. *Am J Otolaryngol* 2018;39:754-8.
11. Nakamura T, Matsui M, Uchida K, et al. M(3) muscarinic acetylcholine receptor plays a critical role in parasympathetic control of salivation in mice. *J Physiol*

- 2004;558:561-75.
12. Durieux ME. Inhibition by ketamine of muscarinic acetylcholine receptor function. *Anesth Analg* 1995;81:57-62.
 13. Gallo A, Benazzo M, Capaccio P, et al. Sialoendoscopy: state of the art, challenges and further perspectives. Round Table, 101(st) SIO National Congress, Catania 2014. *Acta Otorhinolaryngol Ital* 2015;35:217-33.
 14. Wang R, Su YX. Reliable ductal access in sialoendoscopy. *Int J Oral Maxillofac Surg* 2020;49:1592-4.
 15. Chossegros C, Guyot L, Richard O, et al. A technical improvement in sialendoscopy to enter the salivary ducts. *Laryngoscope* 2006;116:842-4.
 16. Squires L, King J, Hsieh TY, et al. Rapid, Effective Cannulation for Salivary Duct Access: Use of Hydro-Dilation With Viscous Lidocaine. *Laryngoscope* 2021;131:E2432-5.
 17. Su CH, Hung SH. Easy insertion into the duct: The use of an angiocatheter as a sialendoscopy applicator. *Laryngoscope* 2018;128:1392-4.
 18. Zenk J, Hosemann WG, Iro H. Diameters of the main excretory ducts of the adult human submandibular and parotid gland: a histologic study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1998;85:576-80.
 19. Bozzato A, Hertel V, Koch M, et al. Vitamin C as contrast agent in diagnosis of salivary duct obstruction. *Laryngorhinootologie* 2009;88:290-2.
 20. McCain JP, Montero J. Surgical Retrieval of Submandibular Stones. *Atlas Oral Maxillofac Surg Clin North Am* 2018;26:111-7.
 21. Schmid RL, Sandler AN, Katz J. Use and efficacy of low-dose ketamine in the management of acute postoperative pain: a review of current techniques and outcomes. *Pain* 1999;82:111-25.
 22. Vadivelu N, Schermer E, Kodumudi V, et al. Role of ketamine for analgesia in adults and children. *J Anaesthesiol Clin Pharmacol* 2016;32:298-306.
 23. Laulin JP, Maurette P, Corcuff JB, et al. The role of ketamine in preventing fentanyl-induced hyperalgesia and subsequent acute morphine tolerance. *Anesth Analg* 2002;94:1263-9, table of contents.
 24. Kohrs R, Durieux ME. Ketamine: teaching an old drug new tricks. *Anesth Analg* 1998;87:1186-93.
 25. Brinck EC, Tiippana E, Heesen M, et al. Perioperative intravenous ketamine for acute postoperative pain in adults. *Cochrane Database Syst Rev* 2018;12:CD012033.
 26. Blankstein KC. Low-dose intravenous ketamine: an effective adjunct to conventional deep conscious sedation. *J Oral Maxillofac Surg* 2006;64:691-2.
 27. Gao M, Rejaei D, Liu H. Ketamine use in current clinical practice. *Acta Pharmacol Sin* 2016;37:865-72.
 28. Nahlieli O, Baruchin AM. Sialoendoscopy: three years' experience as a diagnostic and treatment modality. *J Oral Maxillofac Surg* 1997;55:912-20.
 29. Papadaki ME, McCain JP, Kim K, et al. Interventional sialoendoscopy: early clinical results. *J Oral Maxillofac Surg* 2008;66:954-62.
 30. Moore PA, Hersh EV. Just-in-case opioid prescribing. *J Dent Educ* 2020;84:1327-8.
 31. Resnick CM, Calabrese CE, Afshar S, et al. Do Oral and Maxillofacial Surgeons Over-Prescribe Opioids After Extraction of Asymptomatic Third Molars? *J Oral Maxillofac Surg* 2019;77:1332-6.
 32. Bousquet B, Green MA, Caillouette CN, et al. How Much Opioid Medication Do Patients Need After Orthognathic Surgery? *J Oral Maxillofac Surg* 2022;80:1174-82.
 33. Lanning CF, Harmel MH. Ketamine anesthesia. *Annu Rev Med* 1975;26:137-41.
 34. Newton A, Fitton L. Intravenous ketamine for adult procedural sedation in the emergency department: a prospective cohort study. *Emerg Med J* 2008;25:498-501.
 35. Tucker MR, Hann JR, Phillips CL. Subanesthetic doses of ketamine, diazepam, and nitrous oxide for adult outpatient sedation. *J Oral Maxillofac Surg* 1984;42:668-72.
 36. Wiemer SJ, Nathan JM, Heggstad BT, et al. Safety of Outpatient Procedural Sedation Administered by Oral and Maxillofacial Surgeons: The Mayo Clinic Experience in 17,634 Sedations (2004 to 2019). *J Oral Maxillofac Surg* 2021;79:990-9.
 37. Coniglio AJ, Deal AM, Bhate O, et al. In-office versus Operating Room Sialendoscopy: Comparison of Outcomes, Patient Time Burden, and Charge Analysis. *Otolaryngol Head Neck Surg* 2019;160:255-60.
 38. Trujillo O, Drusin MA, Pagano PP, et al. Evaluation of Monitored Anesthesia Care in Sialendoscopy. *JAMA Otolaryngol Head Neck Surg* 2017;143:769-74.
 39. Pai A, Heining M. Ketamine. *Continuing Education in Anaesthesia Critical Care & Pain* 2007;7:59-63.
 40. Gorlin AW, Rosenfeld DM, Ramakrishna H. Intravenous sub-anesthetic ketamine for perioperative analgesia. *J Anaesthesiol Clin Pharmacol* 2016;32:160-7.

doi: 10.21037/joma-22-28

Cite this article as: Zhu N, Wang TT, Wang J. Potential of ketamine use in sialendoscopy—a narrative review. *J Oral Maxillofac Anesth* 2023;2:9.