

# Antireflux surgery to prevent the progression from Barrett's esophagus to esophageal adenocarcinoma: yes or no?

Marco G. Patti<sup>1</sup>, Fernando A. Herbella<sup>2</sup>, Francisco Schlottmann<sup>3,4</sup>

<sup>1</sup>Department of Surgery, University of Virginia, Charlottesville, VA, USA; <sup>2</sup>Department of Surgery, Escola Paulista de Medicina, Sao Paulo, Brazil;

<sup>3</sup>Department of Surgery, Hospital Alemán of Buenos Aires, Buenos Aires, Argentina; <sup>4</sup>Department of Surgery, University of Illinois at Chicago, Chicago, IL, USA

Correspondence to: Marco G. Patti, MD. Department of Surgery, University of Virginia, Charlottesville, VA, USA. Email: Marco.patti@gmail.com.

**Keywords:** Barrett's esophagus (BE); esophageal adenocarcinoma; gastroesophageal reflux disease (GERD); antireflux surgery

Submitted Jul 27, 2022. Accepted for publication Oct 25, 2022.

doi: 10.21037/jgo-22-719

View this article at: <https://dx.doi.org/10.21037/jgo-22-719>

Gastroesophageal reflux disease (GERD) is due to the reflux of gastric contents from the stomach into the esophagus. Therefore, it is not “acid reflux” only, but a combination of acid and other substances such as bile (1,2). The pathogenesis of GERD is multifactorial, and the most important factors are the transient relaxation and/or low pressure of the lower esophageal sphincter (LES) and weak peristalsis, which causes a slower clearance of the refluxate with increased contact time with the esophageal mucosa. In addition, in obese and morbidly obese patients there is another factor that promotes reflux, the increased thoraco-abdominal gradient between the stomach and the chest, secondary to an increase in the positive intra-abdominal pressure and a more negative intra-thoracic pressure (3).

GERD causes symptoms and/or mucosal damage. Symptoms are defined as esophageal (heartburn, regurgitation, dysphagia) and extraesophageal (aspiration, cough, hoarseness). Endoscopy defines the mucosal injury. The Los Angeles (LA) classification of esophagitis is by far the most widely used system to describe the endoscopic appearance of the esophagitis and grade its severity (4). The LA classification classifies the severity of the esophagitis as follows: grade A, mucosal breaks  $\leq 5$  mm, without continuity across mucosal folds; grade B, mucosal breaks  $>5$  mm, without continuity across mucosal folds; grade C, mucosal breaks continuous between  $>2$  mucosal folds, involving  $<75\%$  of the esophageal circumference; and grade D, mucosal breaks involving  $>75\%$  of the esophageal circumference. About 10 to 15% of patients with GERD

develop an intestinal metaplasia of the epithelium from squamous to columnar with goblet cells, known as Barrett's esophagus (BE) (5). The importance of this finding is due to the possibility that BE can eventually progress from metaplasia to low- and high-grade dysplasia, and cancer (6). While the risk for malignant transformation is low in patients with metaplasia, around 0.33% (7), it is higher in patients with low grade dysplasia, around 0.5% (8). In patients with high grade dysplasia the annual incidence of adenocarcinoma is remarkably high, around 7% (9).

It is very important to keep in mind the composition of the refluxate because medications available today such as proton pump inhibitors (PPI), can only decrease the production of acid by the parietal cells but they do not affect reflux through the LES. On the other hand, a fundoplication can stop any type of reflux, independent of composition and pH, by restoring the competence of the gastroesophageal junction because of an increase in LES pressure and a decrease in transient LES relaxations (10). It has been shown that a fundoplication is more effective than medications in terms of relief of symptoms, particularly regurgitation (11). But by stopping reflux and a further insult to the mucosa, can a fundoplication stop the progression from metaplasia to dysplasia and cancer? This is indeed a very controversial issue, and while some studies have shown a preventive effect, others have failed to show any impact of surgery on the natural history of the disease.

In 2003, Oelschlager and colleagues of the University of Washington in Seattle, studied the effect of laparoscopic

antireflux surgery (LARS) in 90 patients with BE (12). Heartburn improved in 96% of patients and regurgitation in 84%. At a median of 30 months, postoperative endoscopy showed complete regression of the metaplasia in 30 of 54 (55%) patients with short segment BE (< 3 cm), but in none of the 36 patients with a long segment BE ( $\geq 3$  cm). The authors concluded that LARS should be recommended in patients with BE to control symptoms and prevent the progression to cancer. Unfortunately, there was no further follow-up in these patients after the initial study.

Oberg studied the development of dysplasia and adenocarcinoma in 140 patients with Barrett's metaplasia, 94 patients treated with acid suppressing therapy (H2 blocking agents or PPI) and 46 with antireflux surgery (13). At a median follow-up of 5.8 years, endoscopy showed that dysplasia development was significantly less common after antireflux surgery compared with medical therapy. Chang compared patients with BE treated medically with patients after antireflux surgery and determined that the probability of regression was much higher after surgery (1.9% versus 15.4%) (14).

Other studies, however, have shown opposite results. Parrilla *et al.* reported on the long-term results of a randomized and prospective study comparing medical and surgical treatment of BE (15). An excellent and similar control of symptoms was achieved in both groups (91%). Progression to high grade dysplasia occurred in 5% of 43 patients treated medically and in 3% of 58 patients after antireflux surgery. Corey and colleagues performed a meta-analysis to determine if an antireflux operation decreases the incidence of esophageal carcinoma in BE compared to medical therapy (16). They included 4,678 patient-years of follow-up in the surgical group and 4,906 patient-years in the medical group. There was no significant difference in the cancer rates between the two groups: 3.8 cancers/1,000 patient-years compared to 5.3 in the medical group ( $P=0.29$ ). Finally, Maret-Ouda studied the incidence of adenocarcinoma after antireflux surgery in a cohort study from the 5 Nordic countries (17). They compared 48,863 patients who had surgery to 893,208 that did not, and found that the surgical treatment of GERD did not decrease the risk of developing an esophageal adenocarcinoma.

As shown, there is no consensus on the effect of LARS in terms of blocking the progression from metaplasia to dysplasia and cancer. In addition, a conclusion is made even more difficult by the existing interobserver variability in the interpretation of dysplasia, which in part explains the

heterogeneity in terms of progression and treatment.

Based on the existing data, and until a properly conducted multicenter prospective and randomized trial is performed, we do not recommend LARS in patients with BE metaplasia to avoid progression to dysplasia and cancer. As stressed by the most recent guidelines of the American Gastroenterological Association, antireflux surgery should not be considered as an antineoplastic measure in patients with BE (18). The indications for surgery should be the same as for the treatment of GERD without BE, specifically for patients who have symptoms refractory to medical treatment or have complications secondary to the use of acid suppressing therapy.

## Acknowledgments

*Funding:* None.

## Footnote

*Provenance and Peer Review:* This article was commissioned by the editorial office, *Journal of Gastrointestinal Oncology* for the series "Current Management of Upper Gastrointestinal Malignancies". The article did not undergo external peer review.

*Conflicts of Interest:* All authors have completed the ICMJE uniform disclosure form (available at <https://fomm.amegroups.com/article/view/10.21037/jgo-22-719/coif>). The series "Current Management of Upper Gastrointestinal Malignancies" was commissioned by the editorial office without any funding or sponsorship. FS and MGP served as the unpaid Guest Editors of the series. The authors have no other conflicts of interest to declare.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

- Oberg S, Ritter MP, Crookes PF, et al. Gastroesophageal reflux disease and mucosal injury with emphasis on short-segment Barrett's esophagus and duodenogastroesophageal reflux. *J Gastrointest Surg* 1998;2:547-54.
- Fein M, Ireland AP, Ritter MP, et al. Duodenogastric reflux potentiates the injurious effects of gastroesophageal reflux. *J Gastrointest Surg* 1997;1:27-32; discussion 33.
- Del Grande LM, Herbella FAM, Patti MG. The Role of the Transdiaphragmatic Pressure Gradient (TGP) in the Pathophysiology of Proximal Reflux. *J Gastrointest Surg* 2021;25:351-6.
- Lundell LR, Dent J, Bennett JR, et al. Endoscopic assessment of oesophagitis: clinical and functional correlates and further validation of the Los Angeles classification. *Gut* 1999;45:172-80.
- Johansson J, Hakanasson HO, Mellblom L, et al. Prevalence of pre-cancerous and other metaplasia in the distal esophagus and gastro-esophageal junction. *Scand J Gastroenterol* 2005;40:893-902.
- Schlottmann F, Patti MG, Shaheen NJ. From Heartburn to Barrett's Esophagus, and Beyond. *World J Surg* 2017;41:1698-1704.
- Desai TK, Krishnan K, Samala N, et al. The incidence of oesophageal adenocarcinoma in non-dysplastic Barrett's esophagus: a meta-analysis. *Gut* 2012;61:970-6.
- Singh S, Manickam P, Amin AV, et al. Incidence of esophageal adenocarcinoma in Barrett's esophagus with low-grade dysplasia: a systematic review and meta-analysis. *Gastrointest Endosc* 2014;79:897-983.e3.
- Rastogi A, Puli S, El-Serag HB, et al. Incidence of esophageal adenocarcinoma in patients with Barrett's esophagus and high-grade dysplasia: a meta-analysis. *Gastrointest Endosc* 2008;67:394-8.
- Herbella FA, Tedesco P, Nipomnick I, et al. Effect of partial and total laparoscopic fundoplication on esophageal body motility. *Surg Endosc* 2007;21:285-8.
- Rickenbacher N, Kötter T, Kochen MM, et al. Fundoplication versus medical management of gastroesophageal reflux disease: systematic review and meta-analysis. *Surg Endosc* 2014;28:143-55.
- Oelschlager BK, Barreca M, Chang L, et al. Clinical and pathologic response of Barrett's esophagus to laparoscopic antireflux surgery. *Ann Surg* 2003;238:458-66.
- Oberg S, Wenner J, Johansson J, et al. Barrett esophagus: risk factors for progression to dysplasia and adenocarcinoma. *Ann Surg* 2005;242:49-54.
- Chang EY, Morris CD, Seltman AK, et al. The effect of antireflux surgery on esophageal carcinogenesis in patients with barrett esophagus: a systematic review. *Ann Surg* 2007;246:11-21.
- Parrilla P, Martínez de Haro LF, Ortiz A, et al. Long-term results of a randomized prospective study comparing medical and surgical treatment of Barrett's esophagus. *Ann Surg* 2003;237:291-8.
- Corey KE, Schmitz SM, Shaheen NJ. Does a surgical antireflux procedure decrease the incidence of esophageal adenocarcinoma in Barrett's esophagus? A meta-analysis. *Am J Gastroenterol* 2003;98:2390-4.
- Maret-Ouda J, Santoni G, Wahlin K, et al. Esophageal Adenocarcinoma After Antireflux Surgery in a Cohort Study From the 5 Nordic Countries. *Ann Surg* 2021;274:e535-40.
- Shaheen NJ, Falk GW, Iyer PG, et al. Diagnosis and Management of Barrett's Esophagus: An Updated ACG Guideline. *Am J Gastroenterol* 2022;117:559-87.

**Cite this article as:** Patti MG, Herbella FA, Schlottmann F. Antireflux surgery to prevent the progression from Barrett's esophagus to esophageal adenocarcinoma: yes or no? *J Gastrointest Oncol* 2022. doi: 10.21037/jgo-22-719