



Robotic surgery for esophageal cancer: what is the evidence?

Francisco Schlottmann^{1,2}, Fernando A. M. Herbella³, Marco G. Patti⁴

¹Department of Surgery, Hospital Alemán of Buenos Aires, Buenos Aires, Argentina; ²Department of Surgery, University of Illinois at Chicago, Chicago, IL, USA; ³Department of Surgery, Escola Paulista de Medicina, Federal University of Sao Paulo, Sao Paulo, Brazil; ⁴Department of Surgery, University of Virginia, Charlottesville, VA, USA

Correspondence to: Francisco Schlottmann, MD, MPH. Department of Surgery, Hospital Aleman of Buenos Aires, Av. Pueyrredon1640, Buenos Aires, Argentina. Email: fschlottmann@hotmail.com.

Keywords: Esophageal cancer; robotic surgery; esophagectomy

Submitted Jul 27, 2022. Accepted for publication Oct 25, 2022. Published online Oct 29, 2022.

doi: 10.21037/jgo-22-718

View this article at: <https://dx.doi.org/10.21037/jgo-22-718>

Esophageal cancer is a major public health burden, ranking 7th in terms of global cancer incidence and 6th in overall cancer mortality (1). Although esophageal squamous cell carcinoma (ESCC) is the most common histological type worldwide, Western countries have witnessed a rising incidence of esophageal adenocarcinoma (EAC) in the last decades (partly due to the increased prevalence of obesity) (2,3).

Multimodal therapies (i.e., chemotherapy, radiotherapy, immunotherapy, surgery) are often needed to treat esophageal cancer patients. Nevertheless, surgical resection remains the cornerstone in the management of early and locally-advanced esophageal cancer (4). An esophagectomy is a challenging and demanding operation, which requires refined surgical maneuvers and thorough anatomical knowledge of diverse surgical fields (i.e., abdomen, thorax and neck). Since its introduction in the early 1990s, minimally invasive esophagectomy (MIE) has been adopted by several centers achieving lower rates of postoperative complications and better quality of life, as compared to the conventional approach (5-7). MIE, however, is still technically complex (two-dimensional vision, straight non-articulated instruments, limited mediastinal surgical space, mirrored intracorporeal movements) and is associated with high morbidity rates (8).

Robotic surgery has well known advantages over open and conventional MIE (i.e., laparoscopy and thoracoscopy) that facilitate a more precise surgical dissection: enhanced visualization through high-definition and magnified three-dimensional (3D) imaging, stable surgical field, tremor filtration, fingertip control of EndoWrist instruments,

and better ergonomics/reduced fatigue among others. The potential benefits of these technical advantages in esophageal cancer surgery have been explored in previous studies. A randomized trial comparing robot-assisted minimally invasive esophagectomy (RAMIE) with open transthoracic esophagectomy found that RAMIE had lower rates of overall surgery-related complications (59% *vs.* 80%, $P=0.02$), reduced pulmonary complications (32% *vs.* 58%, $P=0.005$), reduced cardiac complications (22% *vs.* 47%, $P=0.006$), lower postoperative pain, and better short-term quality of life. Oncological outcomes were comparable between both approaches (9). Another study also showed RAMIE was associated with decreased intraoperative blood loss, less pulmonary complications and reduced overall morbidity (10).

Opposite to the comparison between RAMIE and open esophagectomy, differences in outcomes between RAMIE and video-assisted esophagectomy are less notorious. Weksler *et al.* compared RAMIE with conventional MIE and found no significant differences in postoperative complications, length of intensive care unit stay or length of hospital stay, arguing that RAMIE did not offer clear advantages over thoracoscopic MIE (11). To note, only few patients were included in the analysis (11 *vs.* 26) and patients who had intrathoracic anastomosis were excluded. A larger propensity scored analysis compared RAMIE ($n=66$) with conventional thoracoscopic esophagectomy ($n=66$) and also reported similar outcomes in terms of blood loss, rates of overall complications, and length of stay (12). In line with these findings, a randomized trial including

only patients with resectable ESCC showed that RAMIE and conventional MIE had comparable rates of overall complications, major complications, anastomotic leakage, and 90-day mortality. RAMIE, however, showed better thoracic lymph node dissection in patients who received neoadjuvant therapy as well as better lymph node dissection along the left recurrent laryngeal nerve (13). A recent meta-analysis including only patients with intrathoracic anastomosis did find benefits of RAMIE (n=974) over conventional MIE (n=5,275): less intraoperative blood loss, lower rates of postoperative pneumonia, reduced overall morbidity, and higher rates of R0 resections (14). Other studies also found superior left recurrent laryngeal lymph node dissection and lower incidence of recurrent laryngeal nerve paralysis with the robotic approach (15,16).

Medical expenditures related to the use of the robot are one of the Achilles' heel of robotic surgery. Both the acquisition and the periodic maintenance of a robotic unit represent an elevated economic burden. For that reason, many large academic institutions all over the world still do not have a robot (17). Cost-effectiveness analyses for robotic esophagectomy, however, are lacking. For instance, savings related to reduced postoperative complications and shorter length of hospital stay could counterbalance the higher costs of RAMIE, as compared to open esophagectomy. Better ergonomics associated with less work-related musculoskeletal complaints (common during conventional MIE) should also be considered (18,19). In addition, the development of new robotic platforms by other manufacturers will create competition in the market likely reducing the required expenses for robotic surgery.

Further advantages of robotic surgery might appear in the near future. Current platforms are capable of capturing kinematic data (e.g., instruments position, distance traveled, grip force) that could be used to design deep-learning models, evaluate surgical performance, and standardize procedures (20). Computerized platforms will also eventually allow navigational (i.e., image guided) surgery and artificial intelligence algorithms that could enhance esophageal cancer surgery.

Overall, current evidence shows that both RAMIE and conventional MIE are associated with better postoperative outcomes than open esophagectomy. In patients undergoing Ivor Lewis esophagectomy (i.e., intrathoracic anastomosis) the robotic approach might offer greater benefits than the classic thoracoscopic technique. In addition, lymph node dissection appears to be improved with the robotic

platform. Potential oncological benefits associated with the use of robotic surgery, if any, remain elusive.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Journal of Gastrointestinal Oncology* for the series "Current Management of Upper Gastrointestinal Malignancies". The article did not undergo external peer review.

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://jgo.amegroups.com/article/view/10.21037/jgo-22-718/coif>). The series "Current Management of Upper Gastrointestinal Malignancies" was commissioned by the editorial office without any funding or sponsorship. FS and MGP served as the unpaid Guest Editors of the series. The authors have no other conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Bray F, Ferlay J, Soerjomataram I, et al. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394-424.
2. Li J, Xu J, Zheng Y, et al. Esophageal cancer: Epidemiology, risk factors and screening. *Chin J Cancer*

- Res 2021;33:535-47.
3. Schlottmann F, Dreifuss NH, Patti MG. Obesity and esophageal cancer: GERD, Barrett's esophagus, and molecular carcinogenic pathways. *Expert Rev Gastroenterol Hepatol* 2020;14:425-33.
 4. Demarest CT, Chang AC. The Landmark Series: Multimodal Therapy for Esophageal Cancer. *Ann Surg Oncol* 2021;28:3375-82.
 5. Cuschieri A, Shimi S, Banting S. Endoscopic oesophagectomy through a right thoracoscopic approach. *J R Coll Surg Edinb* 1992;37:7-11.
 6. Biere SS, van Berge Henegouwen MI, Maas KW, et al. Minimally invasive versus open oesophagectomy for patients with oesophageal cancer: a multicentre, open-label, randomised controlled trial. *Lancet* 2012;379:1887-92.
 7. Maas KW, Cuesta MA, van Berge Henegouwen MI, et al. Quality of Life and Late Complications After Minimally Invasive Compared to Open Esophagectomy: Results of a Randomized Trial. *World J Surg* 2015;39:1986-93.
 8. Casas MA, Angeramo CA, Bras Harriott C, et al. Surgical outcomes after totally minimally invasive Ivor Lewis esophagectomy. A systematic review and meta-analysis. *Eur J Surg Oncol* 2022;48:473-81.
 9. van der Sluis PC, van der Horst S, May AM, et al. Robot-assisted Minimally Invasive Thoracoscopic Esophagectomy Versus Open Transthoracic Esophagectomy for Resectable Esophageal Cancer: A Randomized Controlled Trial. *Ann Surg* 2019;269:621-30.
 10. Mederos MA, de Virgilio MJ, Shenoy R, et al. Comparison of Clinical Outcomes of Robot-Assisted, Video-Assisted, and Open Esophagectomy for Esophageal Cancer: A Systematic Review and Meta-analysis. *JAMA Netw Open* 2021;4:e2129228.
 11. Weksler B, Sharma P, Moudgill N, et al. Robot-assisted minimally invasive esophagectomy is equivalent to thoracoscopic minimally invasive esophagectomy. *Dis Esophagus* 2012;25:403-9.
 12. Zhang Y, Han Y, Gan Q, et al. Early Outcomes of Robot-Assisted Versus Thoracoscopic-Assisted Ivor Lewis Esophagectomy for Esophageal Cancer: A Propensity Score-Matched Study. *Ann Surg Oncol* 2019;26:1284-91.
 13. Yang Y, Li B, Yi J, et al. Robot-assisted Versus Conventional Minimally Invasive Esophagectomy for Resectable Esophageal Squamous Cell Carcinoma: Early Results of a Multicenter Randomized Controlled Trial: the RAMIE Trial. *Ann Surg* 2022;275:646-53.
 14. Angeramo CA, Bras Harriott C, Casas MA, et al. Minimally invasive Ivor Lewis esophagectomy: Robot-assisted versus laparoscopic-thoracoscopic technique. Systematic review and meta-analysis. *Surgery* 2021;170:1692-701.
 15. Tong C, Xu Y, Cao H, et al. Short-term outcomes of robot-assisted versus thoracoscopic-assisted minimally invasive esophagectomy. *Ann Thorac Surg* 2021. [Epub ahead of print]. doi: 10.1016/j.athoracsur.2021.06.073.
 16. Fujita T, Sato K, Ozaki A, et al. Propensity-Matched Analysis of the Short-Term Outcome of Robot-Assisted Minimally Invasive Esophagectomy Versus Conventional Thoracoscopic Esophagectomy in Thoracic Esophageal Cancer. *World J Surg* 2022;46:1926-33.
 17. Childers CP, Maggard-Gibbons M. Estimation of the Acquisition and Operating Costs for Robotic Surgery. *JAMA* 2018;320:835-6.
 18. Alleblas CCJ, de Man AM, van den Haak L, et al. Prevalence of Musculoskeletal Disorders Among Surgeons Performing Minimally Invasive Surgery: A Systematic Review. *Ann Surg* 2017;266:905-20.
 19. Kingma BF, de Maat MFG, van der Horst S, et al. Robot-assisted minimally invasive esophagectomy (RAMIE) improves perioperative outcomes: a review. *J Thorac Dis* 2019;11:S735-42.
 20. Chang TC, Seufert C, Eminaga O, et al. Current Trends in Artificial Intelligence Application for Endourology and Robotic Surgery. *Urol Clin North Am* 2021;48:151-60.

Cite this article as: Schlottmann F, Herbella FAM, Patti MG. Robotic surgery for esophageal cancer: what is the evidence? *J Gastrointest Oncol* 2023;14(4):1913-1915. doi: 10.21037/jgo-22-718