

Aberrant epithelial remodeling with impairment of cilia architecture in non-cystic fibrosis bronchiectasis

Zhuang-Gui Chen^{1,2,3*}, Ying-Ying Li^{2*}, Zhao-Ni Wang^{1*}, Ming Li⁴, Hui-Fang Lim⁵, Yu-Qi Zhou³, Liang-Ming Cai¹, Ya-Ting Li¹, Li-Fen Yang¹, Tian-Tuo Zhang³, De-Yun Wang²

¹Department of Pediatrics, The Third Affiliated Hospital of Sun Yat-sen University, Guangzhou 510630, China; ²Department of Otolaryngology, Yong Loo Lin School of Medicine, National University Health System, National University of Singapore, Singapore; ³Department of Pulmonary Diseases, The Third Affiliated Hospital of Sun Yat-sen University, Institute of Respiratory Diseases of Sun Yat-sen University, Guangzhou 510630, China; ⁴Department of Pulmonary Diseases, The First Affiliated Hospital of Sun Yat-sen University, Guangzhou 510700, China; ⁵Division of Respiratory & Critical Care Medicine, Yong Loo Lin School of Medicine, National University Health System, National University of Singapore, Singapore

Contributions: (I) Conception and design: ZG Chen, TT Zhang, DY Wang; (II) Administrative support: ZG Chen, M Li, HF Lim, TT Zhang, DY Wang; (III) Provision of study materials or patients: ZG Chen, YY Li, ZN Wang; (IV) Collection and assembly of data: YY Li, ZN Wang, M Li, HF Lim, YQ Zhou; (V) Data analysis and interpretation: LM Cai, YT Li, LF Yang; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

*These authors contributed equally to this work.

Correspondence to: Dr. De-Yun Wang. Department of Otolaryngology, Yong Loo Lin School of Medicine, National University Health System, National University of Singapore, 1E Kent Ridge Road, Singapore 119228. Email: entwdy@nus.edu.sg; Dr. Tian-Tuo Zhang. Department of Pulmonary Diseases, The Third Affiliated Hospital of Sun Yat-sen University, Institute of Respiratory Diseases of Sun Yat-sen University, No. 600, Tianhe Road, Tianhe District, Guangzhou 510630, China. Email: zhtituli@163.com.

Background: Aberrant epithelial remodeling and/or abnormalities in mucociliary apparatus in airway epithelium contribute to infection and inflammation. It is uncertain if these changes occur in both large and small airways in non-cystic fibrosis bronchiectasis (non-CF bronchiectasis). In this study, we aim to investigate the histopathology and inflammatory profile in the epithelium of bronchi and bronchioles in bronchiectasis.

Methods: Excised lung tissue sections from 52 patients with non-CF bronchiectasis were stained with specific cellular markers and analyzed by immunohistochemistry and immunofluorescence to assess the epithelial structures, including ciliated cells and goblet cells morphology. Inflammatory cell counts and ciliary proteins expression levels of centrosomal protein 110 (CP110) and dynein heavy chain 5, axonemal (DNAH5) were assessed.

Results: Epithelial hyperplasia is found in both bronchi and bronchioles in all specimens, including hyperplasia and/or hypertrophy of goblet cells. The median cilia length is longer in hyperplastic epithelium [bronchi: 8.16 (7.03–9.14) μm , $P < 0.0001$; bronchioles: 7.46 (6.41–8.48) μm , $P < 0.0001$] as compared to non-hyperplastic epithelium (bronchi: 5.60 μm ; bronchioles: 4.89 μm). Hyperplastic epithelium is associated with overexpression of CP110 and decreased intensity of DNAH5 expression in both bronchial and bronchiolar epithelium. Though infiltration of neutrophils is predominant (63.0% in bronchi and 76.7% in bronchioles), eosinophilic infiltration is also present in the mucosa of bronchi (30.8%) and bronchioles (54.8%).

Conclusions: Aberrant epithelial remodeling with impaired mucociliary architecture is present in both large and small airways in patients with refractory non-CF bronchiectasis. Future studies should evaluate the interplay between these individual components in driving chronic inflammation and lung damage in patients.

Keywords: Epithelial remodeling; mucociliary; non-cystic fibrosis; bronchiectasis; pathology

Submitted Oct 25, 2017. Accepted for publication Jan 29, 2018.

doi: 10.21037/jtd.2018.02.13

View this article at: <http://dx.doi.org/10.21037/jtd.2018.02.13>

Introduction

Non-cystic fibrosis bronchiectasis (non-CF bronchiectasis) is a suppurative lung disease characterized by dilated and distorted airway architecture and abnormal mucus production; patients present with chronic cough with purulent sputum production and recurrent infective exacerbations and progressive functional decline (1). Non-CF bronchiectasis imposes a significant socioeconomic burden on patients (2). They need longer hospital stays, more frequent outpatient visits, and more extensive medical therapy than control subjects, with a cost of approximately \$630 million annually in America (3). Mortality rate ranged 10–16% over an approximate 4-year observation period (3).

The pathogenesis of non-CF bronchiectasis is still under investigation. Currently, the most widely accepted concept for its development is the “vicious cycle hypothesis” (4). It suggests that characteristic airway distortion and abnormal mucus impedes mucociliary clearance (MCC), which in turn promotes bacterial colonization, persistent neutrophilic inflammation and proteases production that perpetuate the process of airway destruction (5–7).

The airway epithelium forms the first line of defense to injurious external stimuli and regulates the immune functions that bridges both innate and adaptive immunity and might be an important regulator of this destructive process. In non-CF bronchiectasis, the epithelium is damaged and has increased susceptibility to injury as compared to normal airway epithelium. Failure of appropriate growth and differentiation of epithelial cells causes persistent mucosal injury (8). Ciliary dyskinesia and mucus over-production weakens clearance of respiratory pathogens from the airway (7). These dysregulated immune responses and repairment contribute to airway structural remodeling, characterized by epithelial hyperplasia, loss of cartilage and fibro-muscular tissues and fibrosis in the bronchi (3,5,9). These pathologic changes might lead to small airways obstruction, chronic inflammation, and progressive lung function impairment (1,9,10).

The histopathological changes of the airway epithelium in the bronchi and bronchioles in non-CF bronchiectasis have not been described. This is an important area to investigate as different lung compartments play different roles and interact to cause lung damage. In recent years, the small airway (<2 mm in diameter) is recognized to be the major site of airflow limitation in chronic inflammatory airway diseases such as asthma and chronic obstructive pulmonary disease (COPD) (11,12). However, small airways

haven't been well studied in non-CF bronchiectasis as thoracic high-resolution computed tomography (HRCT) and spirometric tests cannot accurately assess small airway dysfunction. Thus, we aim to investigate if there is any change in (I) epithelial remodeling (e.g., epithelial hyperplasia and metaplasia); and (II) morphology of goblet and ciliated cells. The results will contribute to a better understanding in pathogenesis of non-CF bronchiectasis.

Methods

Study patients and study design

The study conformed to guidelines of the amended Declaration of Helsinki and has obtained the approval from the Institutional Review Boards of The Third Affiliated Hospitals of Sun Yat-sen University [No. (2017) 2-22]. All patients have given their written consents. We evaluated lung tissues from 52 non-CF bronchiectasis patients aged from 17 to 50 years, who underwent a lobectomy between January 2010 and March 2014 at the First and Third Affiliated Hospitals of Sun Yat-sen University, China. The indications for surgery include: recurrent infection refractory to antibiotics and/or massive hemoptysis that were resistant to medical treatment. The diagnosis of non-CF bronchiectasis was based on thoracic HRCT. Smokers were defined as current cigarette smokers with a smoking history of more than 10 pack years. Lung specimens and clinical information of included patients were collected for histological evaluation and analysis, respectively.

Histological evaluation

Immunohistochemical (IHC) staining

Intraoperative lung biopsies of bronchiectasis were embedded in paraffin and sectioned at 4 µm with a microtome (Leica, Wetzlar, Germany). IHC staining was performed with a modified horseradish peroxidase (HRP) technique with the Dako Cytomation EnVision+System-HRP (Dako A/S). Sections were processed with Target Retrieval Buffer (Dako A/S). Endogenous peroxidase activity was blocked with 0.3% H₂O₂. Sections were then stained and incubated at 4 °C overnight. Eosinophils and lymphocytes were stained by hematoxylin and eosin (H&E). Neutrophils were stained with mouse anti-human neutrophil elastase monoclonal antibody (mAb) (Clone NP57; Dako A/S, Glostrup, Denmark). Cilia were stained with rabbit anti-human dynein heavy chain axonemal

(DNAH5) polyclonal antibody (pAb) (Abcam, Cambridge, MA). The slides were then incubated with Dako Envision +System-HRP (Dako A/S) at room temperature for 30 minutes followed by applying HRP substrate (diaminobenzidine) for color development. All slides were counterstained with hematoxylin.

Immunofluorescence (IF)

Ciliated cells were also assessed by IF staining with mouse anti-human type IV- β tubulin mAb (Clone, ONS.1A6; Abcam) and rabbit anti-human centrosomal protein 110 (CP110) (Proteintech group, Chicago, USA). Tissue sections were processed with Target Retrieval Buffer (Dako A/S). The slides were then incubated with primary antibodies at 4°C overnight, then incubated with Alexa Fluor 488 or 594 conjugated secondary antibodies (goat-anti mouse or rabbit immunoglobulin G [IgG], H+L; Molecular Probes, Carlsbad, CA) at 1:400 in the dark at room temperature for 1 hour, and mounted with Antifade reagent with 4', 6-diamidino-2-phenylindole (DAPI; Molecular Probes).

Goblet cells were analyzed by staining with Alcian Blue periodic acid-Schiff (PAS). Sections were immersed in Alcian Blue pH 2.5 staining solution for 30 minutes, followed by periodic acid solution for 5 minutes, Schiff reagent for 15 minutes, and counterstained with hematoxylin.

Normal adjacent lung tissue as healthy controls

As it's not feasible to obtain open lung biopsy samples from healthy subjects, and the heterogeneity of pathological changes throughout the bronchial tree in patients with bronchiectasis, comparisons were made with the normal lung tissue areas without inflammatory cell infiltration and/or epithelial hyperplasia/remodeling, from the same tissue sections (hereafter referred to as 'healthy controls').

Bronchi and bronchioles

Based on the histological criteria for classifying intrapulmonary structures, airways with wider and non-collapsed lumen, lined with pseudostratified columnar epithelium and the presence of hyaline cartilage plates, goblet cells and glands were defined as bronchi. Airways with narrow collapsible lumen, simple columnar ciliated or cuboidal non-ciliated epithelium without cartilage plates, glands or goblet cells (except larger respiratory bronchioles)

were defined as bronchioles (13,14).

Evaluation of epithelium

In bronchi, epithelium with >4 layers was considered hyperplastic and severe epithelial hyperplasia as >6 layers (15). In bronchioles, epithelium with ≥ 2 layers was considered hyperplastic and severe epithelial hyperplasia as >4 layers (14). Squamous metaplasia was determined by the appearance of the stratified squamous epithelium.

Inflammatory cell infiltration evaluation

Five individual fields with infiltration of inflammatory cells were selected for total and differential cells counts. Total cells counts were derived from counting 200 leukocytes (under $\times 400$ magnification). Differential cell counts of the inflammatory cells were expressed as the percentage—number of positive staining cells/200 leukocytes $\times 100\%$. Inflammatory cell infiltration is categorized by more than 10%.

Goblet cell evaluation

In bronchi, goblet cell hyperplasia was defined if they were in ≥ 2 layers, and goblet cell hypertrophy was defined when doubling in the volume (14). Generally, goblet cells disappear in subsequent divisions in bronchioles. Therefore, goblet cell hyperplasia was defined when it appeared in bronchiolar epithelium.

Cilia length evaluation

Cilia length was evaluated on hyperplastic and non-hyperplastic epithelium in bronchi and bronchioles by assessing type IV- β tubulin staining at $\times 400$ magnifications, as previously described (15). Five areas of the cilia staining from the tissue sections or cytospin cells were randomly selected. Cilia length was measured using Image J software. The mean value of cilia length was calculated from 20 measurements for each paraffin section.

Evaluation of CP110 and DNAH5

Staining images were captured at $\times 400$ magnifications with a fluorescence microscope. Protein expression of these cilia related genes were analyzed by Image J, through calculating the value of the positively stained area and the

Table 1 Clinical characteristics and microbiological results

Characteristics	All patients (n=52)	Male (n=24)	Female (n=28)	P [§]
Age, year	49.06±12.49	52.25±12.7	46.32±11.85	0.088
Smoker	14 (26.93)	11 (48.53)	3 (10.71)	0.017
Positive sputum cultures	6 (11.54)	3 (12.50)	3 (10.71)	NA
<i>Pseudomonas aeruginosa</i>	3 (5.77)	1 (4.17)	2 (7.14)	NA
<i>Haemophilus influenzae</i>	2 (3.85)	2 (8.33)	0	NA
<i>Haemophilus parainfluenzae</i>	1 (1.92)	0	1 (3.57)	NA
Infection*	16 (30.77)	6 (25.00)	10 (35.71)	NA
Aspergillus species	8 (15.38)	3 (12.50)	5 (17.86)	NA
Mycobacterium tuberculosis	6 (11.53)	2 (8.33)	4 (14.28)	NA
Bacterial infection	2 (3.85)	1 (4.16)	1 (3.57)	NA
Inflammatory cell infiltration**				
Eosinophilic	32 (61.53)	15 (62.50)	17 (60.71)	NA
Neutrophilic	41 (78.84)	22 (91.67)	19 (67.85)	0.142
Lymphocytic	39 (75.00)	15 (62.50)	24 (85.71)	0.046

Data are presented as number (%) and mean ± standard deviation. [§], the statistical analysis was performed using chi-square tests; *, infection is evidenced by histopathological examination with specific stain in the excised lung tissue; as for TB detection, patients were diagnosed through the specific pathologic change of tuberculosis in patients' biopsy samples, such as caseous necrosis; **, inflammatory cell infiltration is categorized by more than 10%.

mean fluorescence intensity (MFI) for each marker. Total fluorescence intensity (TFI) measurements were performed through multiplying the positive area by MFI and corrected by subtracting the background autofluorescence (15). The low intensity of DNAH5 positive staining was defined as DNAH5 abnormal expression (16).

Microbiological examination

Patients were routinely received microbiological testing using sputum smear microscopy and sputum culture before operations. After surgery, microorganisms such as fungi and tubercle bacillus will be also detected in the lung specimen. However, as most patients have received long course of antibiotic treatment, microbiological examinations cannot completely explain their etiologies.

Statistical analysis

All data were analyzed using the SPSS statistical software V18.0 (SPSS Inc., Chicago, USA). Continuous variables

were expressed using mean ± standard deviation if normally distributed or median (interquartile range) if not. Categorical variables were expressed by percentage. Differences in continuous variables were compared by using Student's *t*-test or non-parametric test such as Wilcoxon signed rank test, and in categorical variables were compared by using the Chi-squared test or Fisher exact test. Correlation analysis was performed using Spearman *r*. In all analyses, differences were considered significant if *P*<0.05.

Results

Patient characteristics

The demographics and clinical characteristics of patients were summarized in *Table 1*. Their mean age was 49.06 years old. There were significantly more male smokers (21.15%) than female smokers (5.77%, *P*=0.017). *Pseudomonas aeruginosa* (5.77%), *Haemophilus influenzae* (3.85%) and *Haemophilus parainfluenzae* (1.92%) showed positive in bronchial wash culture in 11.54% patients. Aspergillus species (15.38%) and *mycobacterium tuberculosis*

Table 2 Differences in histology in bronchi and bronchioles

Histological change	Bronchi (n=27)*	Bronchioles (n=44)*	P
Epithelial hyperplasia	25 (92.6)	43 (97.7)	0.141
Mild-moderate [#]	13 (48.2)	29 (65.9)	0.139
Severe ^{##}	12 (44.5)	14 (31.8)	0.284
Goblet cell hyperplasia	20 (74.1)	35 (79.6)	0.592
Goblet cell hypertrophy	17 (63.0)	10 (22.7)	0.001
Goblet cell hyperplasia and hypertrophy	15 (55.6)	9 (20.5)	0.004
Squamous metaplasia	2 (7.4)	1 (2.3)	0.141
Cilia length, μm	8.16 (7.03–9.14)	7.46 (6.41–8.48)	<0.0001**

Data are presented as number (%), and median (interquartile range). [#], mild-moderate epithelial hyperplasia is defined as the presence of 5–6 cell layers in bronchi and 3–4 cell layers in bronchioles; ^{##}, Severe epithelial hyperplasia is defined as >6 cell layers in bronchi and > 4 cell layers in bronchioles; *, Bronchi only, bronchioles only and both bronchi and bronchiole structures were seen in 7, 24 and 20 specimens respectively; **, Compared to median lengths of bronchi and bronchioles in non-diseased control regions i.e., 5.60 (4.99–6.61) μm and 4.89 (4.50–5.99) μm respectively.

(MTB, 11.53%) were the main pathogens identified in pathologic analysis in lung biopsy specimens. Lymphocytic inflammation was more prevalent among females as compared to males (62.5% vs. 85.7%, $P=0.046$).

Histopathological examinations

We were able to identify airway epithelium in 51 out of 52 patient specimens – bronchi only, bronchioles only and both bronchi and bronchiole structures were seen in 7, 24 and 20 specimens respectively. *Table 2* summarizes the histological differences between the bronchi and bronchioles.

Epithelium hyperplasia/squamous metaplasia

In bronchiectatic airways (*Figure 1A*) compared to the non-diseased areas, there was an increase in epithelial hyperplasia in both the bronchi (*Figure 1B*) and in bronchioles (*Figure 1C*). There was no difference in the prevalence or severity of epithelial hyperplasia between the bronchi and bronchioles (92.6% vs. 97.7%, $P=0.141$, *Figure 1D*). Squamous metaplasia was noted in 2 specimens (1 bronchi and 1 bronchiole specimen, *Figure 1B,C,D*).

Goblet cell hyperplasia and hypertrophy

The prevalence of goblet cell hyperplasia (*Figure 2*) was similar in both bronchi and bronchioles (74.1% vs. 79.6%, $P=0.592$). The prevalence of goblet cell hypertrophy

(*Figure 2*) was significantly higher in bronchi compared to bronchioles (63.0% vs. 22.7%, $P=0.001$). The prevalence of both goblet cell hyperplasia and hypertrophy was higher in bronchi than in bronchioles (55.6% vs. 20.5%, $P=0.004$). Two specimens did not contain goblet cells (identified by PAS staining).

Ciliary impairment

The median cilia length was significantly longer in both bronchi [8.16 (7.03–9.14) μm , $P<0.0001$] and bronchioles [7.46 (6.41–8.48) μm , $P<0.0001$] in bronchiectasis airways with epithelial hyperplasia, compared to healthy controls [5.60 (4.99–6.61) and 4.89 (4.50–5.99) μm respectively] (*Figure 3A,B*). There was a higher expression of CP110 in both bronchi (2.57-fold, $P=0.0007$) and bronchioles (2.18-fold, $P=0.0005$) with airway hyperplasia, compared to healthy controls (*Figure 3C*). There was a positive correlation between cilia length and CP110 expression in both bronchi ($r=0.410$, $P=0.007$) and in bronchioles ($r=0.406$, $P=0.004$) (*Figure 3D,E*). Decreased intensity of DNAH5 (by IHC staining) was found in 12 (54.4%) bronchial and 19 (57.6%) bronchiolar specimens (*Figure 4*).

Inflammatory cell infiltration

In 51 samples, we found three different patterns of inflammatory cell infiltration (*Figure 5*). The patterns of inflammatory cell infiltration were summarized in *Table 3*.




Figure 1 Epithelium of bronchi and bronchioles. Bronchi has large lumen, pseudostratified columnar epithelium and hyaline cartilage plates with glands in the wall, which were showed by “↑”. Bronchioles simple cuboidal epithelium without cartilage and glands in the wall, and goblet cell disappear. “↑” showed an enlarged bronchioles (as compared to the adjacent pulmonary vein) (A, ×40 magnification; scale bar =200 μm). The common epithelium, epithelial hyperplasia, severe hyperplasia and squamous metaplasia in bronchi (B, ×400 magnification; scale bar =20 μm) and in bronchioles (C, ×400 magnification; scale bar =20 μm). Comparison of epithelial hyperplasia and squamous metaplasia (D, Fisher exact test) in bronchi and bronchioles.




Figure 2 Goblet cell hyperplasia and hypertrophy. The goblet cell hyperplasia and hypertrophy in bronchi and bronchioles (A, $\times 400$ magnification; scale bar = $20\ \mu\text{m}$). Comparison (B, Fisher exact test) of goblet cell hyperplasia and hypertrophy in bronchi and bronchioles.

There was more severe eosinophilic infiltration in the bronchioles (31.0%) compared to the bronchi (3.8%, $P=0.008$). Neutrophils were the predominant cell type in both bronchi (62.9%) and bronchioles (76.8%). Besides, we noted lymphocytic infiltration in over half of the bronchi and bronchiole specimens (70.4% and 59.1%, respectively). Furthermore, mixed neutrophilic and eosinophilic inflammation was seen in 6 bronchial (22.2%) and 18 bronchiolar (42.9%) specimens. A small group (<20%) exhibited mixed neutrophilic/eosinophilic/lymphocytic airway infiltration.

Discussion

In the present study, we found a complex histopathological condition with infection in bronchi and bronchioles of patients with non-CF bronchiectasis. We demonstrated that impaired ciliated cells with lengthened cilia structures on the hyperplastic epithelium in both bronchi and bronchioles

of the patients. These histological findings can be of important in understanding the pathogenesis causing acute and chronic airway infection and inflammation, clinical symptoms of coughing and expectoration, and give more evidences to the current “vicious cycle hypothesis”.

MCC is characterized by the upward movement of mucus by ciliary motion that requires a balance between the volume and composition of the mucus, and normal ciliary beat frequency (17,18). In our study, we showed hyperplasia and/or hypertrophy of the goblet cells in both the large and small airways epithelium, which contributed directly to the over-secretion of the mucus and impairment of MCC in the patients. Mucus production, secretion and clearance play a critical role in maintenance of airway health because it maintains hydration in the airway and traps particulates, bacteria, and viruses (18,19). Furthermore, airway goblet cells are proinflammatory effector cells which are able to secrete pro-inflammatory cytokines, chemokines, and growth factors. These inflammatory mediators released




Figure 3 Cilia impairment. CP110 (red) were double stained with cilia (βIV-tubulin, green) in non-hyperplastic and hyperplastic epithelium in bronchi and bronchioles respectively (A, ×400 amplifications; scale bar =20 μm). Statistical analysis was performed in non-hyperplastic and hyperplastic epithelium in bronchi and bronchioles in cilia length in paraffin section (B, Mann-Whitney two-tailed U test; Median values with 25th and 75th percentiles are indicated by horizontal red lines). CP110 were evaluated by TFI (C, Mann-Whitney two-tailed U test; Median values with 25th and 75th percentiles). The correlations between TFI of CP110 and cilia length (one-to-one) were analyzed respectively (D and E).




Figure 4 Abnormal DNAH5 expression pattern. DNAH5 was evaluated by IHC staining (A, upper: $\times 40$ amplifications; scale bar =200 μm ; down: $\times 400$ amplifications; scale bar =20 μm). Statistical analysis was performed in non-hyperplastic and hyperplastic epithelium in bronchi and bronchioles in DNAH5 expression (B, Fisher exact test).

from goblet cells may act in an autocrine and paracrine manner to enhance inflammation in diseases such as asthma and leading to airway goblet cell hyperplasia (20,21). Our findings therefore present pathological evidences to the current “vicious cycle hypothesis” in the development of bronchiectasis.

Although different types of epithelial cells contribute to the MCC function, ciliated cells play a central role in maintaining MCC function and preventing chronic airway diseases (18,22). Among the tissue specimens, significantly lengthened cilia are present on the hyperplastic epithelium both in bronchi and bronchioles. As we described previously (15), these lengthened cilia are accompanied with impairment of MCC function, which is a potential cause of chronic mucosal inflammation or infection in chronic airway diseases (e.g., rhinosinusitis). In patients with non-CF bronchiectasis, this abnormal-cilia structure will

contribute to impaired MCC, making airways susceptible to chronic infection and microbial colonization, favoring a continuous inflammatory response that persists even after infection has been controlled (3,5,7).

The structural damage to the airways is responsible for progressive small airways obstruction and dilation. Gilley and colleagues reported that conditional deletion of motile cilia in the lung of the adult mouse results in slower cilia beating frequency and the development of bronchiectasis (23). They found that airway dilation in bronchiectasis was due to morphological changes of airway epithelial cells with cellular hyperplasia and proliferation of club cells, and led to loss of cilia and MCC impairment. In bronchiectasis, the mucus itself is often abnormal and more complex. Tracheobronchial clearance in bronchiectasis has been shown to be slower than in normal control subjects independent of the presence of infection (3,5,7). Over




Figure 5 Inflammatory cell infiltration. Inflammatory cell infiltration pattern (A, lower powder view: $\times 100$ amplifications; scale bar = $100\ \mu\text{m}$; B, $\times 400$ amplifications; scale bar = $20\ \mu\text{m}$).

time, retained sputum can cause mucous plugs and airway obstruction, obliteration, and damage resulting in more advanced bronchiectasis (6,7). These structural abnormalities in small airways allow for mucus stasis, which favors continued chronic infection and the persisted vicious cycle in patient with bronchiectasis (8,24).

CP110 is a key marker in ciliogenesis process (25,26). Our previous study demonstrated that the increased expression of CP110 in mucosa from patients with chronic rhino sinusitis (CRS) was associated to the poor ciliation (15). DNAH5 encode for dynein proteins of ciliated airway, and DNAH5 mutations is a common cause of primary ciliary dyskinesia with outer dynein arm defects (16). In this study, we assessed the length of cilia in bronchi and bronchioles and found significantly longer cilia with abnormal expression of ciliogenesis markers (e.g., CP110 and DNAH5) in hyperplastic epithelium as compared to normal lung tissue. The longer and messy cilia with outer dynein mutations may cause MCC dysfunction, and decrease the efficacy of pathogens elimination (22). As such, the continual local microbial invasion and inflammatory damage were contribution to permanent lung tissue alteration and airway structural remodeling (6,7). To our knowledge, this is the first study reporting a positive correlation between cilia length and CP110 in bronchiectasis.

An ongoing cycle of airway tracts infection and

inflammation may be established following a combination of deflection in host defense and bacterial infection (4,7,10). In consistent with previous studies (27-29), inflammatory cells infiltration around airways of non-CF bronchiectasis, including eosinophils, neutrophils and lymphocytes, could be found in both bronchi and bronchioles. As the MCC function is compromised, repeating pathogens infection or colonization led to inflammation, which caused persistent infiltration of different inflammatory cells. Interestingly, the high percentage of eosinophils in respiratory epithelium were observed in patients with bronchiectasis in our study. In fact, most of these patients didn't have asthma according to their clinical records. Thus, the fungal infection might be a potential cause to recruit eosinophils infiltration into airways (30), because *Aspergillus* species infection was found to be the most common pathogen in these excised lung tissue sections from patients with non-CF bronchiectasis.

There are several limitations in this study. This is a retrospective study with a relatively small sample size. As a retrospective study, differences in demographic and clinical characteristics should be matched and compared with the healthy controls, which is not available due to the ethics. There is no standard reference of the qualitative and quantitative measures on goblet and ciliary cells in both bronchi and bronchioles, we could therefore only use the control measures obtained from the same specimens in a

Table 3 Patterns of inflammatory cell infiltration

Inflammatory infiltration	Bronchi (n=27), n (%)	Bronchioles (n=44), n (%)	P
Eosinophilic^a			
None	18 (69.2)	19 (45.2)	NA
10–30%	7 (26.9)	10 (23.8)	0.770
>30%	1 (3.8)	13 (31.0)	0.008
Neutrophilic^b			
None	10 (37.0)	10 (23.3)	NA
10–30%	11 (40.7)	23 (53.5)	0.253
>30%	6 (22.2)	10 (23.3)	0.515
Lymphocytic			
None	8 (29.6)	18 (40.9)	NA
10–30%	13 (48.2)	21 (47.7)	0.595
>30%	6 (22.2)	5 (11.4)	0.268
Eosinophilic & neutrophilic			
None	21 (77.8)	24 (57.1)	NA
10–30%	5 (18.5)	13 (31.0)	0.266
>30%	1 (3.7)	5 (11.9)	0.382
All three	5 (18.5)	5 (11.4)	0.489

^a, two samples don't have HE staining (but one of them had bronchi stained); ^b, one sample doesn't have neutrophil staining for bronchioles.

relative non-diseased area. In this study, the mean age of patients was 49.06 years old, which was younger than the average age in other large-sample reports. However, some studies on surgical treatment of bronchiectasis reported that the median age of these patients was less than 50 years old (31,32). This may be explained by that older patients are more likely accompanied by comorbidities and therefore be less eligible for surgery.

In conclusion, our data demonstrates the different pathologic patterns in bronchi and bronchioles and the impairments of cilia architecture in patients with refractory non-CF bronchiectasis. These findings are of significant importance to our understanding of the “viscous cycle hypothesis” in the development of non-CF bronchiectasis.

Acknowledgements

We thank Dr. Yan Yan and Dr. Li Chunwei for technologic

support. We thank all patients for providing their clinical information and specimens.

Funding: This study was supported by the National Natural Science Foundation of China (Grant no. 81470219).

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

Ethical Statement: The study conformed to guidelines of the amended Declaration of Helsinki and has obtained the approval from the Institutional Review Boards of The Third Affiliated Hospitals of Sun Yat-sen University [No. (2017) 2-22]. All patients have given their written consents.

References

1. Pasteur MC, Bilton D, Hill AT. British Thoracic Society guideline for non-CF bronchiectasis. *Thorax* 2010;65:i1-58.
2. Seitz AE, Olivier KN, Steiner CA, et al. Trends and burden of bronchiectasis-associated hospitalizations in the United States, 1993-2006. *Chest* 2010;138:944-9.
3. McShane PJ, Naureckas ET, Tino G, et al. Non-cystic fibrosis bronchiectasis. *Am J Respir Crit Care Med* 2013;188:647-56.
4. Boyton RJ, Altmann DM. Bronchiectasis: current concepts in pathogenesis, immunology, and microbiology. *Annu Rev Pathol* 2016;11:523-54.
5. Whitters D, Stockley R. Immunity and bacterial colonisation in bronchiectasis. *Thorax* 2012;67:1006-13.
6. Cole PJ. Inflammation: a two-edged sword--the model of bronchiectasis. *Eur J Respir Dis Suppl* 1986;147:6-15.
7. Chalmers JD, Hill AT. Mechanisms of immune dysfunction and bacterial persistence in non-cystic fibrosis bronchiectasis. *Mol Immunol* 2013;55:27-34.
8. Gohy ST, Hupin C, Pilette C, et al. Chronic inflammatory airway diseases: the central role of the epithelium revisited. *Clin Exp Allergy* 2016;46:529-42.
9. King P. Pathogenesis of bronchiectasis. *Paediatr Respir Rev* 2011;12:104-10.
10. Wilson R, Dowling RB, Jackson AD. The biology of bacterial colonization and invasion of the respiratory mucosa. *Eur Respir J* 1996;9:1523-30.
11. Doberer D, Trejo BH, Wenzel SE. Should lung biopsies be performed in patients with severe asthma? *Eur Respir Rev* 2015;24:525-39.

12. Contoli M, Bousquet J, Fabbri LM, et al. The small airways and distal lung compartment in asthma and COPD: a time for reappraisal. *Allergy* 2010;65:141-51.
13. Park S, and Anderson P. Pathology Education Instructional Resource (PEIR). Available online: http://peir.path.uab.edu/wiki/Main_Page
14. Shimizu T, Takahashi Y, Kawaguchi S, et al. Hypertrophic and metaplastic changes of goblet cells in rat nasal epithelium induced by endotoxin. *Am J Respir Crit Care Med* 1996;153:1412-8.
15. Li YY, Li CW, Chao SS, et al. Impairment of cilia architecture and ciliogenesis in hyperplastic nasal epithelium from nasal polyps. *J Allergy Clin Immunol* 2014;134:1282-92.
16. Olbrich H, Horváth J, Fekete A, et al. Axonemal localization of the dynein component DNAH5 is not altered in secondary ciliary dyskinesia. *Pediatr Res* 2006;59:418-22.
17. Liu Y, Di YP. Effects of second hand smoke on airway secretion and mucociliary clearance. *Front Physiol* 2012;3:342.
18. Sears PR, Yin WN, Ostrowski LE. Continuous mucociliary transport by primary human airway epithelial cells in vitro. *Am J Physiol Lung Cell Mol Physiol* 2015;309:L99-108.
19. Sibila O, Suarez-Cuartin G, Rodrigo-Troyano A, et al. Secreted mucins and airway bacterial colonization in non-CF bronchiectasis. *Respirology* 2015;20:1082-8.
20. Tanabe T, Shimokawaji T, Kanoh S, et al. Secretory phospholipases A2 are secreted from ciliated cells and increase mucin and eicosanoid secretion from goblet cells. *Chest* 2015;147:1599-609.
21. Tanabe T, Rubin BK. Airway goblet cells secrete pro-inflammatory cytokines, chemokines, and growth factors. *Chest* 2016;149:714-20.
22. Boon M, Wallmeier J, Ma L, et al. MCIDAS mutations result in a mucociliary clearance disorder with reduced generation of multiple motile cilia. *Nat Commun* 2014;5:4418.
23. Gilley SK, Stenbit AE, Pasek RC, et al. Deletion of airway cilia results in non-inflammatory bronchiectasis and hyperreactive airways. *Am J Physiol Lung Cell Mol Physiol* 2014;306:L162-9.
24. Tilley AE, Walters MS, Shaykhiev R, et al. Cilia dysfunction in lung disease. *Annu Rev Physiol* 2015;77:379-406.
25. Walentek P, Quigley IK, Sun DI, et al. Ciliary transcription factors and miRNAs precisely regulate Cp110 levels required for ciliary adhesions and ciliogenesis. *Elife* 2016;5:e17557.
26. Yadav SP, Sharma NK, Liu C, et al. Centrosomal protein CP110 controls maturation of the mother centriole during cilia biogenesis. *Development* 2016;143:1491-501.
27. Regamey N, Tsartsali L, Hilliard TN, et al. Distinct patterns of inflammation in the airway lumen and bronchial mucosa of children with cystic fibrosis. *Thorax* 2012;67:164-70.
28. Amalakuhan B, Maselli DJ, Martinez-Garcia MA. Update in Bronchiectasis 2014. *Am J Respir Crit Care Med* 2015;192:1155-61.
29. Gaga M, Bentley AM, Humbert M, et al. Increases in CD4+ T lymphocytes, macrophages, neutrophils and interleukin 8 positive cells in the airways of patients with bronchiectasis. *Thorax* 1998;53: 685-91.
30. O'Dea EM, Amarsaikhan N, Li H, et al. Eosinophils are recruited in response to chitin exposure and enhance Th2-mediated immune pathology in *Aspergillus fumigatus* infection. *Infect Immun* 2014;82:3199-205.
31. Jin YX, Zhang Y, Duan L, et al. Surgical treatment of bronchiectasis: a retrospective observational study of 260 patients. *Int J Surg* 2014;12:1050-4.
32. Rademacher J, Ringshausen FC, Suhling H, et al. Lung transplantation for non-cystic fibrosis bronchiectasis. *Respir Med* 2016;115:60-5.

Cite this article as: Chen ZG, Li YY, Wang ZN, Li M, Lim HF, Zhou YQ, Cai LM, Li YT, Yang LF, Zhang TT, Wang DY. Aberrant epithelial remodeling with impairment of cilia architecture in non-cystic fibrosis bronchiectasis. *J Thorac Dis* 2018;10(3):1753-1764. doi: 10.21037/jtd.2018.02.13