

The Society for Translational Medicine: indications and methods of percutaneous transthoracic needle biopsy for diagnosis of lung cancer

Qinghua Zhou¹, Jingsi Dong¹, Jie He², Deruo Liu³, David H. Tian^{4,5}, Shugeng Gao², Shanqing Li⁶, Lunxu Liu⁷, Jianxing He^{8,9}, Yunchao Huang¹⁰, Shidong Xu¹¹, Weimin Mao¹², Qunyou Tan¹³, Chun Chen¹⁴, Xiaofei Li¹⁵, Zhu Zhang¹⁶, Gening Jiang¹⁷, Lin Xu¹⁸, Lanjun Zhang¹⁹, Jianhua Fu¹⁹, Hui Li²⁰, Qun Wang²¹, Lijie Tan²¹, Danqing Li⁶, Qinghua Zhou¹, Xiangning Fu²², Zhongmin Jiang²³, Haiquan Chen^{24,25}, Wentao Fang²⁵, Xun Zhang²⁶, Yin Li²⁷, Ti Tong²⁸, Zhentao Yu²⁹, Yongyu Liu³⁰, Xiuyi Zhi³¹, Tiansheng Yan³², Xingyi Zhang²⁸, Roberto F. Casal³³, Eugenio Pompeo³⁴, Angelo Carretta³⁵, Marc Riquet³⁶, Ottavio Rena³⁷, Pierre-Emmanuel Falcoz³⁸, Hisashi Saji³⁹, Ali Zamir Khan⁴⁰, Jose Luis Danguilan^{41,42}, Diego Gonzalez-Rivas⁴³, Nicolas Guibert⁴⁴, Chengchu Zhu⁴⁵, Jianfei Shen⁴⁵

¹Lung Cancer Center, West China Hospital, Sichuan University, Chengdu 610041, China; ²Department of Thoracic Surgical Oncology, Cancer Institute & Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, National Cancer Center, Beijing 100021, China; ³Department of Thoracic Surgery, China-Japan Friendship Hospital, Beijing 100029, China; ⁴The Collaborative Research (CORE) Group, Macquarie University, Sydney, Australia; ⁵Department of Cardiothoracic Surgery, Royal North Shore Hospital, Sydney, Australia; ⁶Department of Thoracic Surgery, Peking Union Medical College Hospital, Chinese Academy of Medicine, Beijing 100006, China; ⁷Department of Cardiovascular and Thoracic Surgery, West China Hospital, Sichuan University, Chengdu 610041, China; ⁸Department of Thoracic Surgery, The First Affiliated Hospital of Guangzhou Medical University, Guangzhou 510120, China; ⁹State Key Laboratory of Respiratory Disease, National Clinical Research Center for Respiratory Disease, Guangzhou Institute of Respiratory Disease, The First Affiliated Hospital of Guangzhou Medical University, Guangzhou Medical University, Guangzhou 510120, China; ¹⁰Department of Thoracic and Cardiovascular Surgery, The Third Affiliated Hospital of Kunming Medical University (Yunnan Tumor Hospital), Kunming 650100, China; ¹¹Department of Thoracic surgery, Harbin Medical University Cancer Hospital, Harbin 150086, China; ¹²Department of Thoracic Surgery, Zhejiang Cancer Hospital, Hangzhou 310022, China; ¹³Department of Thoracic Surgery, Institute of Surgery Research, Daping Hospital, Third Military Medical University, Chongqing 400042, China; ¹⁴Department of Thoracic Surgery, Fujian Medical University Union Hospital, Fuzhou 350001, China; ¹⁵Department of Thoracic Surgery, Tangdu Hospital, The Fourth Military Medical University, Xi'an 710032, China; ¹⁶Department of Thoracic Surgery, First Affiliated Hospital of Xinjiang Medical University, Urumqi 830054, China; ¹⁷Department of Thoracic Surgery, Shanghai Pulmonary Hospital of Tongji University, Shanghai 200433, China; ¹⁸Department of Thoracic Surgery, Nanjing Medical University Affiliated Cancer Hospital, Jiangsu Key Laboratory of Molecular and Translational Cancer Research, Cancer Institute of Jiangsu Province, Nanjing 210009, China; ¹⁹Department of Thoracic Surgery, Sun Yat-sen University Cancer Center, Guangzhou 510060, China; ²⁰Department of Thoracic Surgery, Beijing Chao-Yang Hospital, Beijing 100043, China; ²¹Department of Thoracic Surgery, Shanghai Zhongshan Hospital of Fudan University, Shanghai 200032, China; ²²Department of Thoracic Surgery, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, China; ²³Department of Thoracic Surgery, Shandong Provincial Qianfoshan Hospital, Shandong University, Jinan 250014, China; ²⁴Department of Thoracic Surgery, Fudan University Shanghai Cancer Center, Shanghai 200032, China; ²⁵Department of Thoracic Surgery, Shanghai Chest Hospital, Jiao Tong University, Shanghai 200000, China; ²⁶Department of Thoracic Surgery, Tianjin Chest Hospital, Tianjin 300051, China; ²⁷Department of Thoracic Surgery, Henan Cancer Hospital, Zhengzhou 450000, China; ²⁸Department of Thoracic Surgery, Second Hospital of Jilin University, Changchun 130041, China; ²⁹Department of Esophageal Cancer, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center for Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin 300060, China; ³⁰Department of Thoracic Surgery, Liaoning Cancer Hospital and Institute, Shenyang 110042, China; ³¹Department of Thoracic Surgery, Xuanwu Hospital of Capital Medical University, Beijing 100053, China; ³²Department of Thoracic Surgery, Peking University Third Hospital, Beijing 100083, China; ³³Department of Pulmonary Medicine, The University of Texas MD Anderson Cancer Center, Texas, USA; ³⁴Department of Thoracic Surgery, Policlinico Tor Vergata, Department of Biomedicine and Prevention, Tor Vergata University, Rome, Italy; ³⁵Department of Thoracic Surgery, San Raffaele Scientific Institute, Milan, Italy; ³⁶Georges Pompidou European Hospital, General Thoracic Surgery Department, Georges Pompidou European Hospital, Paris, France; ³⁷Thoracic Surgery Unit, University of Eastern Piedmont, AOU Maggiore della Carità, Vercelli, Italy; ³⁸Department of Thoracic Surgery, Nouvel Hôpital Civil, Hôpitaux Universitaires de Strasbourg, Strasbourg, France; ³⁹Department of Chest Surgery, St. Marianna University School of Medicine, Kawasaki, Japan; ⁴⁰Department of Minimally Invasive Thoracic Surgery, Medanta The Medicity, Gurgaon, India; ⁴¹Lung Center of the Philippines, Quezon City, Philippines, USA; ⁴²University of the Philippines College of Medicine, Manila, Philippines, USA; ⁴³Department of Thoracic Surgery, Coruña University Hospital, 15006 Coruña, Spain; ⁴⁴Pulmonology Department, Larrey University Hospital, Toulouse, France; ⁴⁵Department of Cardiothoracic Surgery, Taizhou Hospital of Zhejiang Province, Wenzhou Medical University, Taizhou 317000, China

Correspondence to: Jie He, MD, PhD; Shugeng Gao, MD, PhD. Department of Thoracic Surgical Oncology, Cancer Institute & Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, National Cancer Center, Beijing 100021, China. Email: hejie@cicams.ac.cn; gaoshugeng@vip.sina.com.

Submitted May 18, 2018. Accepted for publication Aug 24, 2018.

doi: 10.21037/jtd.2018.09.28

View this article at: <http://dx.doi.org/10.21037/jtd.2018.09.28>

Introduction

In 1883, Leyden successfully performed percutaneous transthoracic needle biopsy (PTNB) of the lung in three patients with pneumonia according the records (1); three years later, Menetrier was the first recorded to employ this technique for the diagnosis of lung cancer (2). So far, PTNB has a history of more than 100 years (3-5). In recent years, PTNB has increasingly been applied in clinical settings along with innovations in imaging devices (6,7), improvement of puncture needles, and advances in cellular pathology (8,9). Currently, PTNB is often performed under the guidance of ultrasound or computed tomography (CT) (10,11).

Indications

The indications for PTNB of unidentified lung lesions with high suspicion of lung cancer include:

- (I) Unidentified peripheral pulmonary nodules or masses especially if localized close to the chest wall, whose pathological nature cannot be identified even after repeated sputum cytology and bronchoscopy (12-22);
- (II) Radiologic imaging suggestive of neoplastic lung lesions with no surgical indication (e.g., lymphomas) or currently non-resectable lung cancer amenable to neoadjuvant chemo-radiotherapy (15);
- (III) Unidentified mediastinal mass or pleural lesion with high suspicion of cancer (10,12-14);
- (IV) Patients who cannot tolerate bronchoscopy but require pathological or cytological characterization of an unidentified lung lesion (10);
- (V) Staging of histologically proven malignancies that have spread to the pulmonary hilum, mediastinum, chest wall, or to more distant sites (14).

Contraindications

Contraindications for PTNB of lung lesions include:

- (I) Isolated pulmonary lesions that are highly suspected

to be malignant and are potentially surgically resectable in order to minimize potential risks of PNB-related cancer seeding and dissemination (23);

- (II) High risk of bleeding (e.g., patients undergoing anticoagulant therapy or with conditions such as von Willebrand disease or severe thrombocytopenia) (24);
- (III) Radiologic imaging suggestive of vascular lesions such as arterial or venous malformations, arteriovenous fistula, and aneurysms (22,25);
- (IV) Severe diffuse or bullous emphysema, severe pulmonary fibrosis;
- (V) Severe pulmonary hypertension; cor pulmonale and/or myocardial infarction; severe cardiopulmonary insufficiency (21);
- (VI) Radiologic imaging suggestive of pulmonary Hydatid disease;
- (VII) Poor general clinical status making PTNB intolerable (24).

Pre-PTNB work-up

Before performing PTNB a rapid work-up must be conducted, including:

- (I) Baseline blood tests, including coagulation profiles, hepatitis/HIV/syphilis status, and urine testing (26);
- (II) ECG and complete pulmonary function tests including transfer factor;
- (III) Chest computed tomography (CT). Enhanced CT is preferred if the lesion is close to the mediastinum and/or major vessel (27-29).

Types of PTNB needles

Percutaneous puncture needles are divided into three classes: aspiration needle, cutting or core needle, and bone-drilling needle.

- (I) Aspiration needles (30-31). These needles are used to obtain cytological specimens. Model sizes range between 16–24 G with an external diameter of

0.6–1.6 mm. They are characterised by a small external diameter and thin walls and thus will cause lesser injury to the tissues with fewer complications. The types of aspiration needles commercially available include Chiba needles, Turner needle, and Greene needles;

- (II) Cutting needles (32-34). These needles are used to obtain tissues samples for histological examinations. They have a relatively large diameter, and are associated with more severe injuries to the tissues and more frequent complications. The types of cutting needles commercially available include Vin-Silverman needles, Trucut needles, and Rotex needles. In recent years, the spring-loaded automatic biopsy needle (also known as biopsy gun) has replaced the conventional cutting needle in many centers. This new device reduces operative time, has higher high success rate, is more convenient in sampling, and reduces the risk of needle track implantation;
- (III) Bone-drilling needles (35). Bone-drilling needles are employed for the biopsy of bone lesions. Their tips have sharp cutting teeth, which enable them to pass through hard bone/cartilage tissues to harvest histological specimens. If appropriate, the Ackermann needles can be used.

PTNB-guiding devices

Percutaneous lung puncture biopsy can be performed under the guidance of fluoroscopy, ultrasound, CT, or magnetic resonance imaging (MRI). Currently, CT-guided PTNB is the most commonly used approach (10,36-37). The addition of spiral CT-based real-time fluoroscopy can make the procedure safer, quicker, and more accurate. For lesions close to the chest wall and if sufficiently large, ultrasound-guided PTNB may be more appropriate (38). MRI-guided biopsy can be performed in multiple axial planes and at any angle, which helps to increase the success rate (39); however, specially designed puncture instruments that can be used in a magnetic field are required.

Technical details of PTNB

Technical details of PTNB can be summarized in a few main steps:

- (I) The patient will be asked to lie in supine, prone, or lateral position based on the location of the lesion.

The chest wall side close to the lesion should be placed upward to facilitate the procedure (13);

- (II) A catheter position-ruler is attached to the site to be punctured, which is then scanned at slice thickness of 5 mm and slice interval of 5 mm (the slice thickness and slice interval can be 2–3 mm for small lesions);
- (III) The optimal puncture level and site are decided, and the angle and depth of the needle insertion are measured (4,40-43);
- (IV) The lesion center is selected as the target puncture level. At this step it is important to avoid skeletal structures, heart/great vessels or pulmonary emphysematous bullae if present (40);
- (V) After the body surface puncture site is marked, disinfection and draping should be routinely performed with the puncture site at the center. Subsequently, the chest wall is anesthetized layer by layer using 2% lidocaine (4,44,45);
- (VI) The patient is asked to take a relaxed breath, and then the needle is inserted via the pre-set needling angle. After the needle approaches the pleura, the patient is asked to hold breath before the needle is quickly advanced to the target site; CT scan is performed immediately at the target level to ensure that the needle tip has reached the target site, after which aspiration biopsy or sampling using an automatic biopsy gun can then be performed (10);
- (VII) The content of the aspiration biopsy needle is gently expelled onto a glass slide, evenly smeared, and then fixed using 95% ethanol. After the tissue needle is withdrawn, the tissue specimen strip is fixed in 10% formalin. Multiple slides are prepared for histological analysis (44,45);
- (VIII) Routine CT scan is performed after puncture to promptly identify any possible complications such as pneumothorax or significant bleeding visible as pleural effusion (40);
- (IX) At the completion of the procedure, the patient is observed for 2–4 hours. Chest X-ray in posterior-anterior and lateral views are performed after 24 hours.

Complications

Percutaneous needle biopsy of lung lesions can cause a variety of complications (31,46-66), which can become even life-threatening complications requiring prompt diagnosis

and adequate treatment. While no special treatment is needed for mild complications, active management and surgical consultation/rescue may be required for severe conditions. After the procedure, the patient has bed rest and avoid vigorous physical activity for the first 24 hours.

The most frequently reported complications include:

- (I) Pneumothorax (5,46-59): pneumothorax is the most common complication. A small pneumothorax may spontaneously resolve without any treatment. Conversely, patients with large pneumothoraces greater than 30% of the lung volume on chest X-ray or becomes symptomatic should have a chest tube inserted;
- (II) Hemorrhage (46-62): localized hematoma in lungs may resolve without treatment. In a similar manner no special treatment is required if small amount of hemoptysis (e.g., phlegm with blood) occurs. Hemostatic drugs may be administered in presence of major or massive hemoptysis. Massive hemothorax is often caused by the injury of intercostal arteries or the pulmonary artery, for which an open or video-assisted thoracoscopic surgery may be required to achieve hemostasis;
- (III) Air embolism (43,55,64,65): air embolism is an extremely rare complication but may cause cerebral or spinal cord infarction, myocardial infarction, arrhythmia, heart failure, or even death. Once air embolism occurs, the patient should be immediately placed left lateral decubitus or Trendelenburg's position to prevent residual air in left atrium to pass into the systemic circulation; concurrently, 100% oxygen must be administered to the patient to promote the absorption of the air embolus. The patient can also be transferred for hyperbaric oxygen treatment;
- (IV) Needle-tract seeding of tumor cells (17,64,65): needle-tract seeding is a relatively common complication of percutaneous lung puncture biopsy, particularly in biopsies performed for sub-pleural lung lesions. Despite its frequency, its risk is often underestimated in the clinical setting. To minimize this risk, negative pressure should be maintained during needle withdrawal while the needle core be protected with a cannula. Repeated use of puncture needle or cutting needle should be avoided;
- (V) Vasovagal pleural irritation (12,66): pleural irritation, characterized by constant cough, dizziness, sweating, paleness, palpitations, weak pulse, cold extremities,

decreased blood pressure, a feeling of chest oppression, prostration, and even consciousness disorders, occurs more frequently in young adults and in women. Once pleural irritation symptoms do occur, chest puncture should be stopped immediately; the patient is then made supine and the hemodynamic parameters be closely monitored. Mild symptoms can resolve spontaneously after rest or psychological counseling. In patients with excessive sweating and/or hypotension, administration of oxygen through a Venturi mask and of intravenous supplementation of fluid boluses of 500 mL of 10% glucose can be given to provide energy while supplementing body fluids. If necessary, 0.3–0.5 mL of adrenaline (1:1,000) should be injected subcutaneously to prevent shock according to the usual clinical experience;

- (VI) Reactive cardiopulmonary arrest (64): reactive cardiopulmonary arrest is an extremely rare and life-threatening complication. Once it occurs, the patient must be immediately rescued according to the standard cardiopulmonary resuscitation care.

Acknowledgements

None.

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

References

1. Leyden H. Uber infectiose Pneumonie. *Deutsch Med Wochenschr*, 9.
2. Menetrier, P. Cancer primitif du poumon. *Bull Soc Anat* 1886;61:643.
3. Haaga JR, Alfydi RJ. Precise biopsy localization by computer tomography. *Radiology* 1976;118:603-7.
4. Westcott JL. Percutaneous transthoracic needle biopsy. *Radiology* 1988;169:593-601.
5. Kazerooni EA, Lim FT, Mikhail A, et al. Risk of pneumothorax in CT-guided transthoracic needle aspiration biopsy of the lung. *Radiology* 1996;198:371-5.
6. Sone S, Takashima S, Li F, et al. Mass screening for lung cancer with mobile spiral computed tomography scanner. *Lancet* 1998;351:1242-5.

7. Marshall D, Simpson KN, Earle CC, et al. Potential cost-effectiveness of one-time screening for lung cancer (LC) in a high risk cohort. *Lung Cancer* 2001;32:227-36.
8. Stanley JH, Fish GD, Andriole JG, et al. Lung lesions: cytologic diagnosis by fine-needle biopsy. *Radiology* 1987;162:389-91.
9. Travis WD, Brambilla E, Burke AP, et al. Introduction to The 2015 World Health Organization Classification of Tumors of the Lung, Pleura, Thymus, and Heart. *J Thorac Oncol* 2015;10:1240-2.
10. Manhire A, Charig M, Clelland C, et al. Guidelines for radiologically guided lung biopsy. *Thorax* 2003;58:920-36.
11. Nakajima T, Yasufuku K, Fujiwara T, et al. Recent advances in endobronchial ultrasound-guided transbronchial needle aspiration. *Respir Investig* 2016;54:230-6.
12. Shaham D. SEMI-INVASIVE AND INVASIVE PROCEDURES FOR THE DIAGNOSIS AND STAGING OF LUNG CANCER I: Percutaneous Transthoracic Needle Biopsy. *Radiol Clin N Am* 2000;38:525-34.
13. Yankelevitz DF, Vazquez M, Henschke CI. Special techniques in transthoracic needle biopsy of pulmonary nodules. *Radiol Clin N Am* 2000;38:267-79.
14. Conces DJ Jr, Schwenk GR Jr, Doering PR, et al. Thoracic needle biopsy. Improved results utilizing a team approach. *Chest* 1987;91:813-6.
15. Penketh AR, Robinson AA, Barker V, et al. Use of percutaneous needle biopsy in the investigation of solitary pulmonary nodules. *Thorax* 1987;42:967.
16. Poe RH, Tobin RE. Sensitivity and specificity of needle biopsy in lung malignancy. *Am Rev Respir Dis* 1980;122:725.
17. Komiya T, Kusunoki Y, Kobayashi M, et al. Transcutaneous needle biopsy of the lung. *Acta Radiol* 1997;38:821.
18. Westcott JL, Rao N, Colley DP. Transthoracic needle biopsy of small pulmonary nodules. *Radiology* 1997;202:97-103.
19. Greif J, Marmor S, Schwarz Y, et al. Percutaneous Core Needle Biopsy vs. Fine Needle Aspiration in Diagnosing Benign Lung Lesions. *Acta Cytol* 1999;43:756-60.
20. The diagnosis, assessment and treatment of diffuse parenchymal lung disease in adults. Introduction. *Thorax* 1999;54 Suppl 1:S1-14.
21. Harrison BD, Thorpe RS, Kitchener PG, et al. Percutaneous Trucut lung biopsy in the diagnosis of localised pulmonary lesions. *Thorax* 1984;39:493.
22. Klein JS, Zarka MA. Transthoracic needle biopsy: an overview. *J Thorac Imaging* 1997;12:232-49.
23. Haramati LB, Austin JH. Complications after CT-guided needle biopsy through aerated versus nonaerated lung. *Radiology* 1991;181:778.
24. Sharma A, Shepard JO. Lung Cancer Biopsies. *Radiol Clin North Am* 2018;56:377-90.
25. Kocijancic I, Kocijancic K. CT-guided percutaneous transthoracic needle biopsy of lung lesions – 2-year experience at the Institute of Radiology in Ljubljana. *Radiol Oncol* 2007;41:99-106.
26. British Thoracic Society Bronchoscopy Guidelines Committee, a Subcommittee of Standards of Care Committee of British Thoracic Society. British Thoracic Society guidelines on diagnostic flexible bronchoscopy. *Thorax* 2001;56 Suppl 1:i1-21.
27. Laroche C, Fairbairn I, Moss H, et al. Role of computed tomographic scanning of the thorax prior to bronchoscopy in the investigation of suspected lung cancer. *Thorax* 2000;55:359-63.
28. Bungay HK, Pal CR, Davies CW, et al. An evaluation of computed tomography as an aid to diagnosis in patients undergoing bronchoscopy for suspected bronchial carcinoma. *Clin Radiol* 2000;55:554-60.
29. Henschke CI, Davis SD, Auh Y, et al. Detection of bronchial abnormalities: comparison of CT and bronchoscopy. *J Comput Assist Tomogr* 1987;11:432-5.
30. Burt ME, Flye MW, Webber BL, et al. Prospective evaluation of aspiration needle, cutting needle, transbronchial, and open lung biopsy in patients with pulmonary infiltrates. *Ann Thorac Surg* 1981;32:146-53.
31. Yap J, Tan KP. A comparative study of the Chiba and Turner needles in percutaneous lung biopsy. *Singapore Med J* 1988;29:14-16.
32. Klein JS, Salomon G, Stewart EA. Transthoracic needle biopsy with a coaxially placed 20-gauge automated cutting needle: results in 122 patients. *Radiology* 1996;198:715-20.
33. Greif J, Marmor S, Schwarz Y, et al. Percutaneous core cutting needle biopsy compared with fine-needle aspiration in the diagnosis of peripheral lung malignant lesions: results in 156 patients. *Cancer* 1998;84:144-7.
34. Lee KL, Griffith JF, Ng WH, et al. CT-guided bone biopsy using battery-powered needle drilling biopsy system: comparison with manual system. *Essr Annual Scientific Meeting* 2013. doi: 10.1594/essr2013/P-0028
35. Milman N. Percutaneous lung biopsy with a fine bore cutting needle (Vacu-Cut): improved results using drill technique. *Thorax* 1995;50:560.

36. Cardella JF, Bakal CW, Bertino RE, et al. Quality Improvement Guidelines for Image-guided Percutaneous Biopsy in Adults. *J Vasc Interv Radiol* 2003;14:S227-30.
37. Yang W, Sun W, Li Q, et al. Diagnostic Accuracy of CT-Guided Transthoracic Needle Biopsy for Solitary Pulmonary Nodules. *Plos One* 2015;10:e0131373.
38. Mao F, Dong Y, Ji Z, et al. Comparison of contrast-enhanced ultrasound and conventional ultrasound for guiding peripheral pulmonary biopsies. *Int J Clin Exp Med* 2017;10:3677-84.
39. Liu S, Li C, Yu X, et al. Diagnostic accuracy of MRI-guided percutaneous transthoracic needle biopsy of solitary pulmonary nodules. *Cardiovasc Intervent Radiol* 2015;38:416-21.
40. Romanes GJ. *Cunningham's Textbook of Anatomy*. Oxford: Oxford University Press, 1981.
41. Glassberg RM, Sussman SK, Glickstein MF. CT anatomy of the internal mammary vessels: importance in planning percutaneous transthoracic procedures. *Ajr Am J Roentgenol* 1990;155:397.
42. Kreula J. A new method for investigating the sampling technique of fine needle aspiration biopsy. *Invest Radiol* 1990;25:245.
43. Aberle DR, Gamsu G, Golden JA. Fatal systemic arterial air embolism following lung needle aspiration. *Radiology* 1987;165:351.
44. Weisbrod GL. Percutaneous fine-needle aspiration biopsy of the mediastinum. *Clin Chest Med* 1987;8:27.
45. Wang KP, Kelly SJ, Britt JE. Percutaneous needle aspiration biopsy of chest lesions. New instrument and new technique. *Chest* 1988;93:993-7.
46. Hwang HS, Chung MJ, Lee JW, et al. C-arm cone-beam CT-guided percutaneous transthoracic lung biopsy: usefulness in evaluation of small pulmonary nodules. *AJR Am J Roentgenol* 2010;195:W400-7.
47. Braak SJ. Pulmonary Masses: Initial Results of Cone-beam CT Guidance with Needle Planning Software for Percutaneous Lung Biopsy. *Cardiovasc Intervent Radiol* 2012;35:1414-21.
48. Lee WJ, Chong S, Seo JS, et al. Transthoracic fine-needle aspiration biopsy of the lungs using a C-arm cone-beam CT system: diagnostic accuracy and post-procedural complications. *Br J Radiol*. 2012;85:e217-22.
49. Choi MJ, Kim Y, Hong YS, et al. Transthoracic needle biopsy using a C-arm cone-beam CT system: diagnostic accuracy and safety. *Br J Radiol* 2012;85:e182-7.
50. Lee SM, Park CM, Lee KH, et al. C-arm cone-beam CT-guided percutaneous transthoracic needle biopsy of lung nodules: clinical experience in 1108 patients. *Radiology* 2014;271:291-300.
51. Cheng YC, Tsai SH, Cheng Y, et al. Percutaneous Transthoracic Lung Biopsy: Comparison Between C-Arm Cone-Beam CT and Conventional CT Guidance. *Transl Oncol* 2015;8:258-64.
52. Jiao D, Yuan H, Zhang Q, et al. Flat detector C-arm CT-guided transthoracic needle biopsy of small (≤ 2.0 cm) pulmonary nodules: diagnostic accuracy and complication in 100 patients. *Radiol Med* 2016;121:268-78.
53. Rotolo N, Floridi C, Imperatori A, et al. Comparison of cone-beam CT-guided and CT fluoroscopy-guided transthoracic needle biopsy of lung nodules. *Eur Radiol* 2016;26:381-9.
54. Jin KN, Park CM, Goo JM, et al. Initial experience of percutaneous transthoracic needle biopsy of lung nodules using C-arm cone-beam CT systems. *Eur Radiol* 2010;20:2108.
55. Choi JW, Park CM, Goo JM, et al. C-arm cone-beam CT-guided percutaneous transthoracic needle biopsy of small (≤ 20 mm) lung nodules: diagnostic accuracy and complications in 161 patients. *AJR Am J Roentgenol* 2012;199:W322-30.
56. Choo JY, Park CM, Lee NK, et al. Percutaneous transthoracic needle biopsy of small (≤ 1 cm) lung nodules under C-arm cone-beam CT virtual navigation guidance. *Eur Radiol* 2013;23:712-9.
57. Cheung JY, Kim Y, Shim SS, et al. Combined fluoroscopy- and CT-guided transthoracic needle biopsy using a C-arm cone-beam CT system: comparison with fluoroscopy-guided biopsy. *Korean J Radiol* 2011;12:89-96.
58. Floridi C, Muollo A, Fontana F, et al. C-arm cone-beam computed tomography needle path overlay for percutaneous biopsy of pulmonary nodules. *Radiol Med* 2014;119:820-7.
59. Jiao de C, Li TF, Han XW, et al. Clinical applications of the C-arm cone-beam CT-based 3D needle guidance system in performing percutaneous transthoracic needle biopsy of pulmonary lesions. *Diagn Interv Radiol* 2014;20:470-4.
60. Jiao DC, Li ZM, Yuan HF, et al. Flat detector C-arm CT-guidance system in performing percutaneous transthoracic needle biopsy of small (≤ 3 cm) pulmonary lesions. *Acta Radiol* 2016;57:677.
61. Busser WM, Braak SJ, Fütterer JJ, et al. Cone beam CT guidance provides superior accuracy for complex needle paths compared with CT guidance. *Br J Radiol* 2013;86:20130310.

62. Hohenforst-Schmidt W, Zarogoulidis P, Vogl T, et al. Cone Beam Computertomography (CBCT) in Interventional Chest Medicine - High Feasibility for Endobronchial Realtime Navigation. *J Cancer* 2014;5:231-41.
63. Hiraki T, Fujiwara H, Sakurai J, et al. Nonfatal Systemic Air Embolism Complicating Percutaneous CT-Guided Transthoracic Needle Biopsy: Four Cases From a Single Institution. *Chest* 2007;132:684-90.
64. Yan GW, Bhetuwal A, Yan GW, et al. A Systematic Review and Meta-Analysis of C-Arm Cone-Beam CT-Guided Percutaneous Transthoracic Needle Biopsy of Lung Nodules. *Pol J Radiol* 2017;82:152-60.
65. Požek, I, Rozman, A. Lung cancer seeding along needle track after CT guided transthoracic fine-needle aspiration biopsy - case report. *Zdravniški Vestnik* 2010;79:659-62.
66. Lazguet Y, Maarouf R, Karrou M, et al. CT guided percutaneous needle biopsy of the chest: initial experience. *Pan Afr Med J* 2016 ;23:211.

Cite this article as: Zhou Q, Dong J, He J, Liu D, Tian DH, Gao S, Li S, Liu L, He J, Huang Y, Xu S, Mao W, Tan Q, Chen C, Li X, Zhang Z, Jiang G, Xu L, Zhang L, Fu J, Li H, Wang Q, Tan L, Li D, Zhou Q, Fu X, Jiang Z, Chen H, Fang W, Zhang X, Li Y, Tong T, Yu Z, Liu Y, Zhi X, Yan T, Zhang X, Casal RF, Pompeo E, Carretta A, Riquet M, Rena O, Falcoz PE, Saji H, Khan AZ, Danguilan JL, Gonzalez-Rivas D, Guibert N, Zhu C, Shen J. The Society for Translational Medicine: indications and methods of percutaneous transthoracic needle biopsy for diagnosis of lung cancer. *J Thorac Dis* 2018;10(9):5538-5544. doi: 10.21037/jtd.2018.09.28