# Prognostic factors for resection of isolated pulmonary metastases in breast cancer patients: a systematic review and meta-analysis

# Jun Fan, Dali Chen, Heng Du, Cheng Shen, Guowei Che

Department of Thoracic Surgery, West China Hospital, Sichuan University, Chengdu 610041, China

*Contributions:* (I) Study Design: J Fan, D Chen, H Du, C Shen and G Che; (II) Search and eligibility assessment: J Fan, D Chen and H Du; (III) Methodological Assessments Data Extraction: J Fan, D Chen and C Shen; (IV) Data Calculation: J Fan and H Du; (V) Manuscript Preparation: J Fan, D Chen, H Du, C Shen and G Che; (VI) Guarantor of the paper: G Che; (VII) Final approval of manuscript: All authors.

Correspondence to: Guowei Che. Department of Thoracic Surgery, West China Hospital, Sichuan University, Chengdu 610041, China. Email: guowei\_che@yahoo.com.

**Background:** Lung is a common organ of metastases in patients with primary breast cancer. Pulmonary metastasis of primary breast cancer is usually considered as a systemic disease, however, the systemic approaches have achieved little progress in terms of prolonging survival time. In contrast, some studies revealed a probable survival benefit of pulmonary metastasectomy for such patients. However, the prognostic factor for pulmonary metastasectomy in breast cancer patients is still a controversial issue. The aim of this study was to conduct a systematic review and meta-analysis of cohort studies to assess the pooled 5-year overall survival (OS) rate and the prognostic factors for pulmonary metastasectomy from breast cancer.

**Methods:** An electronic search in MEDLINE (via PubMed), EMBASE (via OVID), CENTRAL (via Cochrane Library), and Chinese BioMedical Literature Database (CBM) complemented by manual searches in article references were conducted to identify eligible studies. All cohort studies in which survival and/ or prognostic factors for pulmonary metastasectomy from breast cancer were reported were included in the analysis. We calculated the pooled 5-year survival rates, identified the prognostic factors for OS and combined the hazard ratios (HRs) of the identified prognostic factors.

**Results:** Sixteen studies with a total of 1937 patients were included in this meta-analysis. The pooled 5-year survival rates after pulmonary metastasectomy was 46% [95% confidence interval (95% CI): 43-49%]. The poor prognostic factors were disease-free interval (DFI) (<3 years) with HR =1.70 (95% CI: 1.37-2.10), resection of metastases (incomplete) with HR =2.06 (95% CI: 1.63-2.62), No. of pulmonary metastases (>1) with HR =1.31 (95% CI: 1.13-1.50) and the hormone receptor status of metastases (negative) with HR =2.30 (95% CI: 1.43-3.70).

**Conclusions:** Surgery with a relatively high 5-year OS rate after pulmonary metastasectomy (46%), may be a promising treatment for pulmonary metastases in the breast cancer patients with a good performance status and limited disease. The main poor prognostic factors were DFI (<3 years), resection of metastases (incomplete), No. of pulmonary metastasis (>1) and hormone receptor status of metastases (negative). And prospective randomized trials will be needed to address these issues in the future.

Keywords: Breast cancer; pulmonary metastases; prognostic factors; systematic review; meta-analysis

Submitted Mar 21, 2015. Accepted for publication Jul 15, 2015. doi: 10.3978/j.issn.2072-1439.2015.08.10 View this article at: http://dx.doi.org/10.3978/j.issn.2072-1439.2015.08.10

# Introduction

Breast cancer is the most frequently diagnosed cancer and the leading cause of cancer death among females, accounting for 23% of the total cancer cases and 14% of the cancer deaths (1). In Asia, one in every three women faces the risk of breast cancer in their lifetime as per report of WHO 2012. Distant metastases appear in 20% of women who have primary breast cancer (2). And lung is a common organ of metastases in patients with primary breast cancer, with approximately 12% of the breast cancer patients found to have metastatic lesions in the lung (3).

Pulmonary metastasis of primary breast cancer is usually considered as a systemic disease, so most medical oncologists disapprove of surgical procedures for metastatic breast cancer or consider them just as palliative strategies (4). Only systemic treatments are routinely considered for such patients. Yet recently, the systemic approaches even with taxanes and/or anthracycline have achieved little progress in terms of prolonging survival time (5,6). Highdose chemotherapy with stem-cell transplantation has also failed to prolong life expectancy (7). The median survival after chemotherapy remains at about 24 months, and the cure rate is very low (8). Thus, metastatic breast cancer is still regarded as an incurable disease, and the treatment is usually just palliative (9). However, several studies revealed a probable survival benefit of pulmonary metastasectomy for such patients. In the study by Staren et al., which compared the medical and surgical management of pulmonary metastatic disease, the mean 5-year survival of the medically treated group was only 11% as comparing to the surgical group, whose mean 5-year survival was 36% (10). A significantly increased survival for the patients operated on comparing with the patients treated conservatively was also demonstrated by Meimarakis et al. (11) and Yhim et al. (12) in their respective studies. At the same time, with the technological advances in cardiothoracic surgery and anesthesia, the pulmonary metastasectomy is associated with a low perioperative morbidity and lethality now (11,13). Consequently, pulmonary metastasectomy may be a promising treatment for lung metastases of breast cancer. However, the prognostic factor of pulmonary metastasectomy in breast cancer patients is still a controversial issue.

In the current study, we conducted a systematic review and meta-analysis of cohort studies to assess the pooled 5-year overall survival (OS) rate and the prognostic factors for survival in pulmonary metastasectomy from breast Fan et al. Prognostic factors for resection of PM from BC

cancer, which may help the clinicians to evaluate the potential surgical benefits of the patients with different prognostic factors and make clinical decisions.

# **Materials and methods**

This systematic review and meta-analysis was performed according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Statement protocol (14).

# Search strategy

We searched MEDLINE (via PubMed), EMBASE (via OVID), CENTRAL (via Cochrane Library), and Chinese BioMedical Literature Database (CBM) to October 2014 to identify studies relevant to this review. Our search strategy included the following subject headings and/or key words variably combined by "breast neoplasm", "lung metastasis", and "surgery". The detailed search strategy of PubMed is listed here: (((("breast neoplasms" [MeSH Terms]) OR breast cancer\*) OR breast tumor\*) OR breast carcinoma\*) AND (("lung neoplasms/secondary"[MeSH Terms]) OR lung metastas\*) AND (("surgical procedures, operative" [MeSH Terms]) OR resect\*). In addition, reference lists of the articles initially detected were searched by hand to identify additional relevant reports. The eligibility of references retrieved by the search was assessed independently by two of the authors (Jun Fan and Dali Chen), and the review authors resolved differences of opinion by discussion or by appeal to a third review author (Heng Du) when necessary. The full text of the remaining articles, including the references, was examined to determine whether the articles contained relevant information.

# Inclusion and exclusion criteria

Studies were considered eligible if they met all of the following inclusion criteria: (I) studies on surgery for pulmonary metastases from breast cancer; (II) enrolled patients whose primary breast tumors had been resected completely proven by histopathology and metastatic disease was limited to the lung proven by imaging techniques; (III) 5-year survival rate and prognostic factors for patients after pulmonary metastasectomy were reported in the individual studies. Studies were excluded based on any of the following criteria: (I) the following article types were excluded: reviews, letters, laboratory researches, animal experiments;

(II) the language is not English or Chinese; (III) the articles studied on the lung nodules(not only the lung metastases) in breast cancer patients.

# Quality assessment

Quality assessment of individual studies was performed independently by two of the authors (Jun Fan and Dali Chen), using the Newcastle-Ottawa Scale for cohort studies. The scale allocates stars (maximum of 9) for quality of selection, comparability and outcome of study participants (15). Any discrepancies were addressed by joint reevaluation of the Original Article.

# Data extraction

Data were extracted from the selected studies independently by two of the authors (Jun Fan and Dali Chen), using a predefined standardized form and resolved disagreements by discussion between two review authors or by appeal to a third review author (Cheng Shen). We extracted data of the eligible articles' basic characteristics, including first author (year), country, study period, type of study, No. of patients, No. of patients evaluated survival and prognostic factors after pulmonary resection, median follow-up, median/mean age at pulmonary metastasectomy, median disease-free interval (DFI) which is the interval between surgery for the primary breast cancer and the appearance of pulmonary metastases, median OS time after pulmonary metastasectomy, 5-year survival rate and prognostic factors. Moreover, the original data included the Kaplan-Meier (K-M) survival curves or hazard ratio (HR) and 95% confidence interval (95% CI) of survival outcomes. The K-M survival curves' data was extracted by Engauge Digitizer 4.1 (http:// sourceforge.net) and the HR, which couldn't be gotten directly from the individual studies, was estimated by the methods of Tierney et al. (16).

#### **Outcome** assessment

The analysis focused on assessing the pooled 5-year survival and the prognostic factors for OS in the patients who underwent pulmonary metastasectomy from breast cancer.

# Data analysis and statistical analysis

Prognostic factors associated with outcome were extracted

from all cohorts. A prognostic association was considered significant if the reported P value was less than 0.05, if the article reported that an association was significant, or if the 95% CI around a rate ratio or similar statistic did not cross 1. Only prognostic factors that were assessed via univariate analyses (log-rank test) are presented, this decision was taken due to the different statistical techniques and choice of covariates used in the individual multivariate models. We decided that the collective interpretation of factors drawn from different multivariate models may be potentially misleading.

To assess the pooled 5-year survival rate, proportions were Log transformed to satisfy the normality. And then we calculated the pooled 5-year survival rate with 95% CI by the DerSimonian & Laird method. When the 5-year survival rate and/or 95% CI was not available, these values were estimated according to data obtained from the K-M curve, using an actuarial method. We also carried out metaanalysis on the prognostic factors for OS using HR as a statistic. Heterogeneity was quantified using a chi-square heterogeneity statistic and by means of an I<sup>2</sup> statistic for each analysis. Heterogeneity was defined as P<0.10 or  $I^2$ >70%. When homogeneity was fine (P $\ge$ 0.10; I<sup>2</sup> $\le$ 70%), a fixed effect model was used to combine effective sizes or else, a random effect model was used. Meta-regression was used to assess the impact of publish year and the prognostic factors on the 5-year survival rate. The potential publication bias was evaluated by the symmetry of funnel and Egger's test, with P>0.05 indicating no potential publication bias (17). All analyses were performed using R software and the metafor package (18). P values less than 0.05 were considered significant.

# **Results**

# Reference retrieval

After primary retrieval, a total of 2,207 potentially relevant studies were incorporated into our initial study, including 746 in Medline, 1,369 in Embase, 47 in CBM, 40 in CENTRAL and 5 by reference list. A total of 403 were excluded for duplicates and 1,756 were excluded by title/abstract screening. Full texts were retrieved for the remaining 48 studies, and 16 of them met all the criteria for inclusion in the analysis, which included 1,937 patients (*Figure 1, Table 1*). All of the patients were female except two, which were excluded from this analysis.

1444



Figure 1 PRISMA Flow Chart of Reference Retrieval.

# Characteristics and qualities of the included studies

Basic characteristics of included studies were summarized in Table 1. All of the 16 studies were cohort studies, in which only one study (23) was prospective, and the other 15 studies were retrospective. Eleven studies were published after 2000, and five studies were published in the 1990s. Nine studies analyzed survival and prognostic factors after pulmonary resection in only part of the enrolled patients, for the reasons of loss to follow-up or the patients with non-metastatic pulmonary nodules (benign pulmonary nodules or primary lung cancer). The median/mean age at pulmonary metastasectomy of included studies ranged from 49 to 63 years old. Reported median survival in these series was 31-96.9 months. Most of the median follow-up of the involved studies were short, and only in three studies, the median follow-up was more than 5 years. And there were seven studies which were conducted with a sample size less than 50 patients.

Quality assessments of individual studies were shown in *Table 2*. We used the Newcastle-Ottawa Scale (NOS) for cohort studies to assess included studies, which included three aspects (selection, comparability and outcome) and eight items. Fifteen studies scored not less than 6.

#### Fan et al. Prognostic factors for resection of PM from BC

# Prognostic factors for OS

There were fifteen prognostic factors which were summarized in Table 3. The most frequently reported significant prognostic factors were DFI (nine studies), resection of metastases (four studies), No. of pulmonary metastases (three studies) and the hormone receptor status of metastases (three studies). And the number and proportion of the patients with the four main prognostic factors in respective studies were shown in Table 1. None of the involved studies reported there was a significant association between the OS and the following prognostic factors, which included additional adjuvant therapy, age at metastasectomy, type of surgical procedures, nodal status of primary tumor, other recurrence site before lung surgery, histologic grade of breast cancer. Among all of the fifteen prognostic factors, three prognostic factors (Stage of breast cancer at breast surgery, Nodal status of primary tumor, Histologic grade of breast cancer) were related to the primary breast cancer. However, there is only one study which reported that there was a significant association between the stage of breast cancer at breast surgery and survival.

The four main prognostic factors for OS and the pooled HRs were shown in *Table 4* and *Figure 2*. All of the four main prognostic factors had significant associations with the OS and the pooled HR values ranged from 1.31 to 2.30. And there were no significant heterogeneities found in all of the four meta-analyses, so we choose the fixed effect model to do the meta-analyses.

# 5-year OS rate

The pooled 5-year survival rate was 46% (95% CI: 43-49%) (*Figure 3*). And the test for heterogeneity revealed significant heterogeneity ( $I^2$ =78.5%, P<0.001). Therefore, we assessed whether the study year and the four main prognostic factors could explain the heterogeneity by metaregression. We found that there was a significant association between the study year and the 5-year survival rate shown by univariable meta-regression analysis (P=0.0323). And no significant associations were found between the four main prognostic factors and 5-year survival rate (*Table 5*).

# Assessment of publication bias

We assessed the publication bias of the meta-analysis in 5-year survival rate. The funnel plot (*Figure 4*) was drawn with log of proportions along the horizontal axis and

lable I Basic cha	actenstics of 1	Incluaeu	studies									Prognostic fac	tors, No.	(%)
Author (reference)	Country	Study period	Type of study	z	c	Age [year]	Follow-up [months]	DFI [months]	Median OS (months)	o-year OS rate (%)	DFI (<3 y)	Resection of meta* (incomplete)	No. of meta (>1)	HR of meta (negative)
Chen <i>et al.</i> , 2009 (19)	Japan	1991- 2007	Retrospective	41	41	55 [35-81]	25 [2-142]	55 [0-192]	NR	51	15 (36.6)	(0) 0	21 (51.2)	RN
Ludwig <i>et al.</i> , 2003 (20)	Germany	1989- 1998	Retrospective	25	21	54 [35.3-77.3]	RN	99.2 [13.4-165.8]	96.9	53	NR	0) 0	9 (42.9)	NR
Yoshimoto <i>et al.</i> , 2008 (21)	Japan	1960- 2000	Retrospective	06	06	55.1 [32-77]	79.2	67.2 [0-244.8]	75.6	54	33 (36.6)	10 (11.1)	12 (13.3)	NR
Lanza <i>et al.</i> , 1992 (22)	NSA	1981- 1990	Retrospective	44	37	55 [32-79]	NR	NR	47	49.50	NR	0) 0	10 (27.0)	15 (51.7)
Yhim <i>et al.</i> , 2010 (12)	Korea	1997- 2007	Retrospective	45	15	49 [36-65]	50 [5.0-136]	NR	NR	82.10	NR	(0) 0	4 (26.7)	10 (66.7)
Meimarakis <i>et al.</i> , 2013 (23)	Germany	1982- 2007	Prospective	81	81	58.2 [28.3-76.3]	NR	RN	NR	59.60	NR	15 (18.5)	30 (37.0)	14 (31.1)
Welter <i>et al.</i> , 2008 (24)	Germany	1998- 2007	Retrospective	47	47	55.0 [34.7-80]	20.6 [3.2-110]	44.4 [0-309.6]	32	36	18 (38.3)	14 (29.8)	18 (38.3)	25 (56.8)
Staren <i>et al.</i> , 1992 (10)	NSA	1972- 1990	Retrospective	63	33	50	NR	48	55	36	NR	RN	6 (18.2)	NR
McDonald <i>et al.</i> , 1994 (25)	NSA	1982- 1992	Retrospective	60	59	58 [21-81]	42 [1-128.4]	26.4 [1-247.2]	42	37.80	NR	20 (33.9)	28 (47.5)	NR
Simpson <i>et al.</i> , 1997 (26)	Australia	1984- 1996	Retrospective	17	17	59 [40-74]	NR	61.2 [8-218.4]	NR	62	8 (47.1)	RN	3 (18.8)	NR
Friedel <i>et al.</i> , 2002 (27)	International	1960- 1994	Retrospective	467	467	53 [21-87]	34 [0-240]	43	35	35	181 (39.0)	75 (16.1)	148 (32.5)	NR
Tanaka <i>et al.</i> , 2005 (28)	Japan	1992- 2001	Retrospective	52	39	53.7 [35-75]	NR	66.8 [11.5-104.8]	32	30.80	NR	NR	24 (61.5)	NR
Kycler and Laski, 2012 (29)	Poland	1994- 2002	Retrospective	33	33	53.4 [33-75]	71.3 [0-232]	51.9	73.2	54.50	16 (48.5)	18(54.5)	19 (57.6)	20 (60.6)
Rena <i>et al.</i> , 2007 (30)	Italy	1990- 2003	Retrospective	79	25	63 [42-82]	43 [3-190]	NR	NR	38	10 (40.0)	RN	NR	NR
Friedel <i>et al.</i> , 1994 (31)	NSA	1979- 1992	Retrospective	91	89	53 [23-78]	NR [2-144]	NR	31	27	NR	21(23.6)	28 (41.2)	19 (57.6)
Planchard <i>et al.</i> , 2004 (32)	France	1972- 1998	Retrospective	162	125	53 [30-82]	102 [1-264]	36 [2-252]	50.4	45	65 (52.0)	29(23.2)	35 (28.0)	25 (52.1)
N, No. of patients; 5-year OS rate, 5 metastases; HR, h	n, No. of anal -year overall : ormone recep	yzing sur survival r ytor, whic	vival and prognos rate; DFI, disease ih includes estrogi of as no tumor cel	en re-	inter ceptc	after pulmo val, betwee vr (ER) and p	inary resection in surgery for in of the rese	on; median OS or the primary receptor (PR),	, median o tumor and and only v	verall sur d the app when both	vival time bearance h of them	after pulmona of pulmonary are negative,	metasi metasta HR is ne	:asectomy; ses; meta, gative; NR,

Table 2 Quality assessment of individual studies using the Newcastle-Ottawa Scale (NOS) for cohort studies

				2	Sele	ction					Compara	tbility				Outcom	e			
Study		в			q		U			q	e			f		D		ح	й 	core
	a1* a	2* a3	a4	b1*	b2 t	3 c1'	c2*	0 03	4 d1	* d2	e1*	e2* f	*	2* f3	f4	g1* g	2 h1*	h2* h3	h4	
Chen <i>et al.</i> , 2009 (19)	*			*		*					*	*		*			*			2
Ludwig <i>et al.</i> , 2003 (20)	*			*		*					*			*			*			9
Yoshimoto <i>et al.</i> , 2008 (21)	*			*		*					*			*		*	*			7
Lanza <i>et al.</i> , 1992 (22)	*			*		*					*			*			*			9
Yhim <i>et al.</i> , 2010 (12)	*			*		*					*	*		*			*			7
Meimarakis <i>et al.</i> , 2013 (23)	*			*		*			*		*	*		*			*			8
Welter <i>et al.</i> , 2008 (24)	*			*		*					*			*			*			9
Staren <i>et al.</i> , 1992 (10)		*		*		*					*			*			*			9
McDonald <i>et al.</i> , 1994 (25)	*			*		*					*			*			*			9
Simpson <i>et al.</i> , 1997 (26)		*		*		*					*			*		*	*			7
Friedel <i>et al.</i> , 2002 (27)	*			*		*					*	*		*			*			7
Tanaka <i>et al.</i> , 2005 (28)	*			*		*					*			*						5
Kycler and Laski, 2012 (29)	*			*		*					*			*		*	*			7
Rena <i>et al.</i> , 2007 (30)	*			*		*					*			*				*		9
Friedel <i>et al.</i> , 1994 (31)	*			*		*					*	*		*				*		7
Planchard <i>et al.</i> , 2004 (32)	*			*		*					*			*		*		*		7
a, representativeness of the	expo	sed co	short:	a1, tr	uly re	preser	itative;	a2, s	omew	hat repr	esentati	ve; a3,	selec	ted gr	o dno	f users;	a4, no	o descriptic	on of 1	the
derivation of the cohort. b, s	electio	n of th	le non-	-expo	sed c	short:	o1, drav	wn fro	m the	same co	ommunil	ty as th∈	exp	osed ci	ohort;	b2, drav	wn fror	m a differer	it sour	;ee;
b3, no description of the de	rivatior	ז of th∈	∋ non-(	sodxe	ed co	hort. c	, asceri	tainme	ent of	exposure	e: c1, se	cure rec	cord;	c2, stn	ucture	d intervi	iew; c3	3, written se	elf-repo	ort;
c4, no description. d, demo	nstratic	on that	outco	ime oi	f inter	est wa	s not p	resen	t at sta	art of stu	udy: d1,	yes; d2	, no.	e, com	ıparab	ility of o	cohorts	s on the ba	sis of .	the
design or analysis: e1, study	/ contre	ols for	the m	ost im	porta	nt fact	or; e2, s	study	contro	ols for an	y additio	onal fac	tor. f,	assess	sment	of outc	ome: f	1, indepenc	ent bl	lind
assessment; f2, record linka	ge; f3,	self-re	port; f	4, no	descr	iption.	g, was	follov	/-up lo	none enou	igh for o	utcome	s to c	scour: ç	g1, yes	s; g2, no	o. h, ac	lequacy of	follow	dn
of cohorts: h1, complete foll	dn wol	-all s	ubject	s acct	ounted	d for; h	2, subj	jects I	ost to	follow u	ip unlike	ly to inti	roduc	se bias	-sma	dmun lli	er lost	>80% follo	w-nb,	, or

description provided of those lost; h3, follow-up rate <80% and no description of those lost; h4, no statement.

 Table 3 Prognostic factors for which univariate analyses were reported in at least two cohorts

Prognostic factors	Number/significant*
DEI	15/9 (12 19 21 22 26 27 29 30 32)
<5 years	1/0
	8/7 (10 21 26 27 20 20 22)
	6/7 (19,21,20,27,29,50,52) E/1 (10)
<2 years	5/1 (12)
<1 year	1/1 (22)
Resection of metastases	8/4 (23,27,29,31)
(incomplete vs. complete)	
No. of pulmonary	14/3 (19,23,25)
metastases (multiple vs.	
solitary)	
The HR status of	9/3 (12,23,24)
metastases (negative vs.	
positive)	
Size of the largest	6/2 (23,32)
metastasis (≥2 <i>vs.</i> <2 cm)	
Site of metastases	6/2 (19,29)
(bilateral vs. unilateral)	
Stage of breast cancer at	5/1 (21)
breast surgery	
No. of metastasectomies	3/1 (29)
(redo vs. primary)	
Pleural invasion (yes vs. no)	2/1 (23)
Additional adjuvant	6/0
therapy	
Age at metastasectomy	6/0
Type of surgical	4/0
procedures	
Nodal status of primary	4/0
tumor	
Other recurrence site	2/0
before lung surgery	
Histologic grade of breast	2/0
cancer	

\*, number of cohorts in which the factor was reported/ number of cohorts in which a significant association with poor outcome was reported (log-rank test,  $\alpha$ <0.05). DFI, disease-free interval.

standard error along the vertical axis. We didn't find a significant asymmetry in the funnel plot by Egger's test (P=0.7879). So there is no publication bias for meta-analysis in 5-year survival rate.

# Discussion

To our knowledge, it is the first time that a comprehensive and detailed systematic review and meta-analysis is performed to evaluate the survival and prognostic factors for resection of isolated pulmonary metastases in breast cancer patients. We discussed the prognostic factors and their associations with the OS. At the same time, we analyzed the pooled 5-year survival rate and the association between the prognostic factors and 5-year survival.

Pulmonary metastasectomy has become a standard therapy for various metastatic malignancies of the lungs. It has been reported that surgery of lung metastases of almost all solid tumors provided good long-term results (33). And the International Registry of Lung Metastases reports on a 5-year survival of 36% in 4,572 patients having resection of lung metastases (34). However, it is a controversial issue for lung resection in breast cancer. In the current analysis, the pooled 5-year OS rate after pulmonary metastasectomy in the breast cancer patients is 46%. In contrast, the 5-year survival was 16% in a case series of patients with breast cancer metastases limited to the lungs and treated with chemotherapy only (35). This result implies that surgery is a promising treatment for the pulmonary metastases in breast cancer patients. The potential survival benefit may be gotten from the reduction of the tumor burden and tumor heterogeneity by resection of pulmonary metastases. However, it should be emphasized that surgical candidates represented a highly selected population with a good performance status and limited disease. Furthermore, in addition to the primary surgical treatment, almost all of these patients in respective studies more or less received some chemotherapy or/and endocrine therapies. So the interpretation of the result should be prudent. In spite of that, at least this result could imply a trend that some selected breast cancer patients may get some survival benefits from the resection of pulmonary metastases.

Previous researches provided inconsistent findings as to the prognostic factors in the patients undergoing pulmonary metastasectomy from the breast cancer. In the present analysis, we extracted all of the prognostic factors from the sixteen involved studies, which univariate analyses were reported in at least two studies, and counted the number of times that a significant association was reported (*Table 3*). We found that DFI >3 years, complete resection of metastases, solitary pulmonary metastasis and positive hormone receptor status of metastases were most frequently reported as the significant good prognostic

#### Fan et al. Prognostic factors for resection of PM from BC

F B						
Prognostic factors	Ν	Patients	Heterogeneity (I-squared, P)	Model	Hazard ratio (95% CI)	P value
DFI (<3 years)	5	683	34.0%, 0.195	Fixed	1.70 (1.37-2.10)	0.000
Resection of metastases (incomplete)	4	670	25.1%, 0.261	Fixed	2.06 (1.63-2.62)	0.000
No. of pulmonary metastases (>1)	3	569	15.2%, 0.308	Fixed	1.31 (1.13-1.50)	0.000
The hormone receptor status of metastases (–)	6	302	0.0%, 0.623	Fixed	2.30 (1.43-3.70)	0.001

Table 4 Main	prognostic facto	ors for su	irvival and	the pooled	hazard ratios
--------------	------------------	------------	-------------	------------	---------------

N, reference count; DFI, disease-free interval.



Figure 2 Evaluated hazard ratios summary for: (A) DFI (<3 years) for overall survival; (B) resection of metastases (incomplete) for overall survival; (C) No. of pulmonary metastases (>1) for overall survival; (D) hormone receptor status of metastases (negative) for overall survival. DFI, disease-free interval.

factors. So patients with these good prognostic factors may survive longer after pulmonary metastasectomy. And then we analyzed the magnitude of the association between the four main prognostic factors and survival by the pooled HRs (*Table 4*). We found that the hormone receptor status of metastases with the pooled HR 2.30 is the strongest prognostic factors among the four main factors. It may be because that the positive hormone receptor represents a slow-growing tumor biology with less aggressive clinical feature and these patients had a better response

# 1448



**Figure 3** Forest plots showing 5-year survival in each study. Each square represents an individual survival, with the size of the square being proportional to the weight given to the study. The dotted and dashed vertical lines represent combined survival for the whole population.

**Table 5** Relationship between the four main prognostic factors

 and 5-year survival

Characteristics	5-year su	irvival
Characteristics	Coefficient <sup>a</sup>	P value <sup>b</sup>
Study year	0.0204	0.0323
DFI (<3 years)	0.8933	0.5399
Resection of metastases	-0.6099	0.2474
(incomplete)		
No. of pulmonary metastases (>1)	-0.6493	0.2471
The hormone receptor status of	-0.0381	0.9772
metastases (–)		

<sup>a</sup>, coefficient represents a linear relationship between each prognostic factor and the survival outcome by a univariable meta-regression model. The sign of the coefficient indicates the positivity or negativity of the relationship. <sup>b</sup>, the P value (<0.1) of each coefficient indicates whether the linear correlation was significant or not.

to the endocrine treatment. It is interesting that the DFI (<3 years), which was reported as a significant prognostic factor in 7 of 8 studies, didn't have a very strong association with survival (the pooled HR 1.70). Consequently, although there were so many cohorts reporting DFI (<3 years) as a significant factor, the cut-off of DFI, 3 years, maybe was not adequate, for there were some studies choosing 1 year, 2 years or 5 years as their cut-offs. A further study would be needed to address this issue. Another interesting thing we found was that none of the four main prognostic factors



**Figure 4** Funnel plot and Egger's test for evaluation of the publication bias for 5-year survival rate.

would significantly affect the 5-year survival rate. However, it should be noted that there were some missing values in this analysis, which maybe also led to this result. So we should interpret this result cautiously.

With the development of imaging techniques, such as multidetector-row computed tomography (MDCT) and positron emission tomography (PET), we can detect earlier and smaller lung nodules in the breast cancer patients. So more and more breast cancer patients were detected with a lung nodules after mastectomy. However, the histological types of the lung nodules may be pulmonary metastases, primary lung cancers or benign lesions (28,30). The diagnosis of the histological types of lung nodules is crucial to the following therapies. Imaging examinations may be sometimes only indicative but not diagnostic. Fine-needle aspiration biopsy using CT-guidance or trans-bronchial needle aspiration biopsy is available to diagnose the lung nodules. However, when the lung nodule is small or its location is not favourable for diagnostic sampling, the biopsy may fail to diagnose. In that case, surgery, especially video-assisted thoracic surgery, is a useful modality to confirm pathologic diagnosis of pulmonary nodules in breast cancer patients and guide the clinicians to choose the following therapies.

Our study has several limitations. The findings of a meta-analysis depend on the qualities of the individual studies, as their potential problems and biases may affect the pooled effects. According to the quality assessment by NOS, fourteen of the sixteen involved studies scored 6 or 7, which indicated the qualities of most of the studies were only moderate. And there were seven individual studies conducted with a sample less than 50 patients and fifteen of sixteen cohorts were retrospective studies. At the same time, most studies confirmed the diagnosis of lung metastasis only according to the inpatient medical records or the records in clinical cases database. Meanwhile, some prognostic factors were assessed in only a few studies, which may also weaken the role of meta-analysis. And we didn't search unpublished and grey literature database, which may lead to a potential publication bias. Maybe there is also a language bias, for we only screened the literature in English and Chinese.

In conclusion, surgery with a relatively high 5-year OS rate after pulmonary metastasectomy (46%), may be a promising treatment for the breast cancer patients with a good performance status and limited disease, who were diagnosed as isolated pulmonary metastases. For the breast cancer patients with pulmonary nodules, who failed to be diagnosed by biopsy, surgery is an available method to confirm the pathologic diagnosis. At the same time, the main poor prognostic factors for resection of isolated pulmonary metastases in breast cancer patients were DFI (<3 years), resection of metastases (incomplete), No. of pulmonary metastasis (>1) and hormone receptor status of metastases (negative). And prospective randomized trials will be needed to address these issues in the future.

# **Acknowledgements**

We acknowledge Jing Bai, of the Department of Cardiovascular Medicine of the First Affiliated Hospital of Zhengzhou University for her careful revision of the manuscript.

# Footnote

*Conflicts of Interest:* The authors have no conflicts of interest to declare.

# References

- 1. Jemal A, Bray F, Center MM, et al. Global cancer statistics. CA Cancer J Clin 2011;61:69-90.
- Chlebowski RT, Kuller LH, Prentice RL, et al. Breast cancer after use of estrogen plus progestin in postmenopausal women. N Engl J Med 2009;360:573-87.
- Kreisman H, Wolkove N, Finkelstein HS, et al. Breast cancer and thoracic metastases: review of 119 patients. Thorax 1983;38:175-9.
- Cardoso F, Harbeck N, Fallowfield L, et al. Locally recurrent or metastatic breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and followup. Ann Oncol 2012;23 Suppl 7:vii11-9.
- Sledge GW Jr. Should we dream the impossible dream? The meaning of long-term survival in metastatic breast cancer. J Clin Oncol 1996;14:2191-3.
- 6. Giordano SH, Buzdar AU, Smith TL, et al. Is breast cancer survival improving? Cancer 2004;100:44-52.
- Gerrero RM, Stein S, Stadtmauer EA. High-dose chemotherapy and stem cell support for breast cancer: where are we now? Drugs Aging 2002;19:475-85.
- Greenberg PA, Hortobagyi GN, Smith TL, et al. Longterm follow-up of patients with complete remission following combination chemotherapy for metastatic breast cancer. J Clin Oncol 1996;14:2197-205.
- 9. Cabuk D, Basaran G, Teomete M, et al. Clinical outcome of Turkish metastatic breast cancer patients with currently

available treatment modalities--single center experience. Asian Pac J Cancer Prev 2014;15:117-22.

- Staren ED, Salerno C, Rongione A, et al. Pulmonary resection for metastatic breast cancer. Arch Surg 1992;127:1282-4.
- 11. Meimarakis G, Angele M, Staehler M, et al. Evaluation of a new prognostic score (Munich score) to predict longterm survival after resection of pulmonary renal cell carcinoma metastases. Am J Surg 2011;202:158-67.
- Yhim HY, Han SW, Oh DY, et al. Prognostic factors for recurrent breast cancer patients with an isolated, limited number of lung metastases and implications for pulmonary metastasectomy. Cancer 2010;116:2890-901.
- Monteiro A, Arce N, Bernardo J, et al. Surgical resection of lung metastases from epithelial tumors. Ann Thorac Surg 2004;77:431-7.
- Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. Int J Surg 2010;8:336-41.
- Wells GA, Shea B, O'Connell D, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. In: 2008. Available online: http://www.ohri.ca/programs/clinical\_ epidemiology/oxford.asp
- Tierney JF, Stewart LA, Ghersi D, et al. Practical methods for incorporating summary time-to-event data into metaanalysis. Trials 2007;8:16.
- Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. Biometrics 1994;50:1088-101.
- Viechtbauer W. Conducting Meta-Analyses in R with the metafor Package. J Stat Software 2010;36:1-45. Available online: http://www.jstatsoft.org/v36/i03/paper
- Chen F, Fujinaga T, Sato K, et al. Clinical features of surgical resection for pulmonary metastasis from breast cancer. Eur J Surg Oncol 2009;35:393-7.
- 20. Ludwig C, Stoelben E, Hasse J. Disease-free survival after resection of lung metastases in patients with breast cancer. Eur J Surg Oncol 2003;29:532-5.
- Yoshimoto M, Tada K, Nishimura S, et al. Favourable long-term results after surgical removal of lung metastases of breast cancer. Breast Cancer Res Treat 2008;110:485-91.
- 22. Lanza LA, Natarajan G, Roth JA, et al. Long-term survival after resection of pulmonary metastases from carcinoma of the breast. Ann Thorac Surg 1992;54:244-7; discussion 248.
- 23. Meimarakis G, Rüttinger D, Stemmler J, et al. Prolonged overall survival after pulmonary metastasectomy in patients

with breast cancer. Ann Thorac Surg 2013;95:1170-80.

- 24. Welter S, Jacobs J, Krbek T, et al. Pulmonary metastases of breast cancer. When is resection indicated? Eur J Cardiothorac Surg 2008;34:1228-34.
- McDonald ML, Deschamps C, Ilstrup DM, et al. Pulmonary resection for metastatic breast cancer. Ann Thorac Surg 1994;58:1599-602.
- 26. Simpson R, Kennedy C, Carmalt H, et al. Pulmonary resection for metastatic breast cancer. Aust N Z J Surg 1997;67:717-9.
- 27. Friedel G, Pastorino U, Ginsberg RJ, et al. Results of lung metastasectomy from breast cancer: prognostic criteria on the basis of 467 cases of the International Registry of Lung Metastases. Eur J Cardiothorac Surg 2002;22:335-44.
- Tanaka F, Li M, Hanaoka N, et al. Surgery for pulmonary nodules in breast cancer patients. Ann Thorac Surg 2005;79:1711-4; discussion 1714-5.
- 29. Kycler W, Laski P. Surgical approach to pulmonary metastases from breast cancer. Breast J 2012;18:52-7.
- 30. Rena O, Papalia E, Ruffini E, et al. The role of surgery in the management of solitary pulmonary nodule in breast cancer patients. Eur J Surg Oncol 2007;33:546-50.
- Friedel G, Linder A, Toomes H. The significance of prognostic factors for the resection of pulmonary metastases of breast cancer. Thorac Cardiovasc Surg 1994;42:71-5.
- 32. Planchard D, Soria JC, Michiels S, et al. Uncertain benefit from surgery in patients with lung metastases from breast carcinoma. Cancer 2004;100:28-35.
- Martini N, McCormack PM. Evolution of the surgical management of pulmonary metastases. Chest Surg Clin N Am 1998;8:13-27.
- Pastorino U, Buyse M, Friedel G, et al. Long-term results of lung metastasectomy: prognostic analyses based on 5206 cases. J Thorac Cardiovasc Surg 1997;113:37-49.
- 35. Diaz-Canton EA, Valero V, Rahman Z, et al. Clinical course of breast cancer patients with metastases confined to the lungs treated with chemotherapy. The University of Texas M.D. Anderson Cancer Center experience and review of the literature. Ann Oncol 1998;9:413-8.

**Cite this article as:** Fan J, Chen D, Du H, Shen C, Che G. Prognostic factors for resection of isolated pulmonary metastases in breast cancer patients: a systematic review and meta-analysis. J Thorac Dis 2015;7(8):1441-1451. doi: 10.3978/j.issn.2072-1439.2015.08.10