

Characterization of a new subtype of allergen in *dermatophagoides farinae* – Der f 28

Jian-Li Lin^{1*}, Yuan-Yuan Wang^{1*}, Xiao-Jun Xiao¹, Yu-Lan Wu¹, Bao-Qing Sun², An-Jian Gao¹, Zhi-Gang Liu¹, Jing Li², Ping-Chang Yang¹, Xiao-Yu Liu¹

¹State Key Laboratory of Respiratory Disease for Allergy, School of Medicine, Shenzhen University, Shenzhen 518060, China; ²State Key Laboratory of Respiratory Disease, The First Affiliated Hospital, Guangzhou Medical University, Guangzhou 510120, China

Contributions: (I) Conception and design: J Lin, X Liu; (II) Administrative support: Z Liu, P Yang, J Li; (III) Provision of study materials or patients: All authors; (IV) Collection and assembly of data: J Lin, Y Wang, X Xiao; (V) Data analysis and interpretation: J Lin, Y Wu, B Sun; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

*These authors contributed equally to this work.

Correspondence to: Dr. Xiao-Yu Liu and Dr. Ping-Chang Yang. Shenzhen University, Room 722 of Medical School Bldgn, 3688 Nanhai Blvd, Shenzhen 518060, China. Email: lxy0901@163.com and pcy2356@szu.edu.cn; Dr. Jing Li. State Key Laboratory of Respiratory Disease, The First Affiliated Hospital, Guangzhou Medical University, 151 Yan Jiang Road, Guangzhou 510120, China. Email: jingli1016@vip.163.com.

Background: House dust mites (HDMs) are the major sources of indoor allergens which induce asthma, dermatitis, rhinitis, and some other allergic diseases. Close to 30 sub-allergens have been identified.

Methods: Through analyzing the full genome sequence of dust mite, a new allergen whose primary structure belongs to the heat shock protein family was identified. The sequence of this allergen was determined by cDNA cloning. The allergenicity was assayed by skin prick test, Western-blot and enzyme-linked immunosorbent assay (ELISA).

Results: r-Der f 28 bound to serum IgE from mite allergic patients. Positive responses to r-Der f 28 were shown in 11.5% by skin prick testing from 26 DM-allergic patients. Airway hyperresponsiveness, serum specific IgE and IL-4 were significantly increased in allergic asthma mouse model sensitized to r-Der f 28.

Conclusions: Der f 28 is a new subtype of allergen in *dermatophagoides farinae*.

Keywords: *Dermatophagoides farinae*; Der f 28; western-blotting; enzyme-linked immunosorbent assay (ELISA); skin prick test; allergic asthma; mouse model

Submitted Oct 13, 2015. Accepted for publication Oct 16, 2015.

doi: 10.3978/j.issn.2072-1439.2015.10.43

View this article at: <http://dx.doi.org/10.3978/j.issn.2072-1439.2015.10.43>

Introduction

A significant increase in the prevalence of allergic diseases has been an important public health problem in the world (1,2). House dust mites (HDMs) are globally important sources of indoor allergens (3), responsible for the sensitization of more than 50% of allergic patients (4,5). At present, *antigen SIT* (specific immunotherapy) with HDM extracts is the most efficient treatment for *dust mite* allergic diseases. However, the crude extracts of HDM not only contain inflammatory molecules, such as ceramides, kallikreins and the ever popular endotoxin, but also include allergens,

which have been proved having side effects in the clinical treatment (6). Specific immunotherapy with HDM extracts is an efficient treatment but severe side-effects may occur in the course of treatment. Therefore, recombinant allergens have been regarded as a substitute for crude mite extracts used in clinical immunotherapy which may effectively improve the efficacy and safety of house dust-mites-specific immunotherapy.

A large number of studies have been conducted to understand the structural properties, chemical and biological of dust mite allergens. Two groups of mite allergens (group 1 and 2) have been proven to be the major

allergens in dermatophagoides farinae. They are a cysteine protease and an epididymal protein, respectively. More than 80% of humans with HDM allergy have IgE antibody to the group 1 and more than 90% of the group 2 (7-10). However, about 20% of patients do not have IgE antibody to the two group allergens (7). Close to 30 sub-allergens have been recognized and characterized in *D. farina* (7,11). Heat shock proteins are allergens in *Hordeum vulgare*, *Malassezia globosa* and HDM (12-15). Yet, the allergenicity of the heat shock proteins in *D. farina* has not been defined. Therefore, it is necessary to characterize the *D. farinae*-derived heat shock protein in the initiation of allergic responses.

In this study, we cloned, expressed and purified the heat shock proteins from HDM, which was nominated as Der f 28 according to the published series of subtypes of HDM allergens. The allergenicity of Der f 28 was tested with a mouse model. The results showed that the r-Der f 28 was capable of inducing allergic asthma in mice.

Materials and methods

Mice

Six-to-eight-week-old female BALB/c mice, weighting 18-20 g, were obtained from Southern Medical University Laboratory Animal Center (Guangzhou, China). The mice were exposed to a 12 h light/dark cycle, maintained under conditions of constant humidity (60%) and temperature (24 °C), and provided tap water to drink, as well as SPF level standard mouse food. The experimental procedures were approved by the Institutional Ethics Committee at Shenzhen University (Shenzhen, China).

Sera

The sera of patients with allergic disorders were obtained from the First Affiliated Hospital of Guangzhou Medical University. The Sera from non-allergic individuals were used as normal controls. The using human sera in the present study was approved by the Human Ethic Committee at Shenzhen University.

Obtaining the encoding genes of Der f 28 and sequence analysis

The gene sequence of Der f 28 was obtained by blast with the whole genome of *Dermatophagoides farinae*. Total RNA was isolated from the homogenates of mite with

Trizol (Invitrogen, Carlsbad, CA, USA) following the manufacturer's instructions. The cDNA was synthesized with a SMARTer™ PCR cDNA Synthesis Kit (Clontech Laboratories, Mountain View, CA, USA) following the manufacturer's protocol. Briefly, 1 µg total RNA in 3.5 µL double-distilled H₂O was mixed with 1 µL 3' SMART CDS Primer IIA and then incubated serially at 72 °C for 3 min and at 42 °C for 2 min. The reaction was transferred to a 5.5 µL aliquot of master mix containing 2 µL 5× First-strand buffer, 0.25 µL DTT (100 mM), 1 µL dNTP (10 mM), 1 µL SMARTer IIA oligonucleotide (12 µM), 0.25 RNase inhibitor, and 1 µL SMARTScribe™ reverse transcriptase (RT, 100 U). The RT reaction was allowed to proceed at 42 °C for 1 h, followed by heating at 70 °C for 10 min, and then a transfer to a 90 µL aliquot of PCR master mix containing 10 µL 10× advantage 2 PCR buffer, 2 µL 50× dNTP (10 mM), 2 µL 5' PCR Primer2A (12 µM), and 2 µL 50× advantage 2 polymerase. The PCR conditions were as follows: 95 °C for 1 min followed by 15 cycles at 95 °C for 15 s, 65 °C for 30 s, and 68 °C for 3 min. The quality of the cDNA was confirmed by a Bioanalyzer.

The resulted Der f 28 sequence was converted to an amino acid sequence. Compared with the amino acids sequence of sixteen different sub-cloned heat shock protein in GenBank, molecular evolutionary tree was constructed by software MEGA5.

Expression and purification of the recombinant Der f 28

The PCR products were ligated into pMD19-T vector (TaKaRa, Dalian, China), the constructed plasmids were transformed into *E.coli* top10. The positive clones were sequenced by BGI (Shenzhen, China). The pMD19-T vector with Der f 28 product was digested. Der f 28 fragments were recycled from the gel and ligated into the pET44-a expression vector. Initial cloning was performed with *E.coli* strain Top10. DNA extracted from positive clones was digested with the BamH I and Hind III restriction enzymes and followed by sequencing. The plasmids were transformed into *E. coli* BL21 cells for expression. The cells were grown in LB-medium supplemented with 50 µg/mL ampicillin to an absorbance of 0.6 at 600 nm. Expression of Der f 28 was induced by adding isopropyl-D-thiogalactopyranoside (IPTG) to LB medium at a final concentration of 0.1 mM and the incubation was continued for 4 h at 37 °C. Cells were harvested by centrifugation and resuspended in 50 mM Tris-HCl, 100 mM NaCl, pH7.5. The cells were sonicated at an

amplitude of 38% for 5 min (1 second pulse on and 1 second pulse off) followed by centrifugation at 10,000 rev/min at 4 °C for 20 min. Finally the recombinant protein was purified by affinity chromatography using the Ni⁺ as a solid phase. After vigorous washing with a washing buffer (50 mM Tris, 40 mM imidazole and 0.5 M NaCl, pH8), the protein was eluted slowly using an elution buffer (50 mM Tris, 0.3 M imidazole and 0.2 M NaCl, pH8). The elution peak was collected and purified by molecular sieve chromatography.

Skin prick test (SPT) of restructured dust mites r-Der f 28

SPTs with purified and endotoxin removed r-Der f 28 was dissolved in pH7.4 phosphate buffer (50 mM PB, 100 mM NaCl). Glycerin was added to 50% final concentration. The sample concentration was 0.01 mg/mL of histamine phosphate (0.1%) or used saline as a negative control. The results were checked 20 minutes after SPT. The judgments of the result: if the prick spot became a wheal and fleck surrounding the wheal, it was positive (+). 4+: the response was stronger than histamine control; 3+: the response was almost the same as histamine control; 2+: the response was weaker than histamine, but stronger than negative control; 1+: the response was significantly weaker than histamine, but slightly stronger than the negative control; negative: No response. The approval to conduct these studies was obtained from the ethics committee of the hospital. All participants were provided written, informed consent for participating in this study.

Immunoblotting analysis with sera from allergic patients to dust mites

The r-Der f 28 protein was fractioned by SDS-PAGE (sodium dodecyl sulfate polyacryl-amide gel electrophoresis) and electro-transferred onto a nitrocellulose membrane (Bio-Rad, Hercules, CA, USA). The membrane was blocked for 2 hours with 3% bovine serum albumin (BSA) at room temperature, and then incubated with the sera from patients sensitized to dust mite. After washing three times with TBS-T (Tris-buffered saline containing triton X-100), the membrane was incubated with the secondary antibodies (mouse anti human IgE) for 60 minutes at room temperature. In the end, after washing with TBS-T for three times, the membrane was developed with a DAB kit (Invitrogen, Carlsbad, CA, USA).

Enzyme-linked immunosorbent assay (ELISA)

Sera IgE antibodies specific to Der f 28 were measured by indirect ELISA. The microtiter plates were coated overnight at 4 °C with at 100 ng/well in Carbonate buffered solution (pH9.5). Then blocked at 37 °C with 200 µL 3% BSA in PBS for 120 min. The serum (diluted to 5 times) or BSA (using as a negative control) was then added to each well (100 µL/well) and incubated for 60 min at room temperature. The plates were then incubated with peroxidase-labeled goat anti-human IgE (1:2,000) for 60 min at 37 °C. Each incubating step was followed by three washes with PBST. The color was developed by adding tetramethylbenzidine (TMB) (100 µL/well) and stopped by the addition of 2M H₂SO₄ (50 µL/well). The plates were read by an ELx808 absorbance microplate reader (BioTek, Shanghai, China) at 450 nm.

Development of a mouse model of allergic asthma

Eighteen mice were divided into three groups: (I) HDM group; (II) Der f 28 group; (III) PBS group. In HDM group and Der f 28 group, mice were injected with 100 µg extracts from dermatophagoides farinae or r-Der f 28 group with 4 mg aluminum hydroxide as adjuvants, respectively. PBS was used in the control group. The sensitization was performed on day 0, 3 and 7. From day 14, the challenge was performed once a day for a week. HDM and Der f 28 group was challenged with 50 µg crude extract of dermatophagoides farinae or Der f 28 protein, respectively. Following the challenge, Airway hyperresponsiveness to methacholine was measured with the unrestrained whole-body plethysmography (WBP). The mice were subjected to progressively increasing doses of methacholine (0, 6.25, 12.5, 25, 50, 100 mg/mL), and the Penh value was recorded. Bronchoalveolar lavage (BAL) and different cell counts were performed as previously reported (16).

Analysis of IL-4 and specific IgE in serum

Blood samples were collected from the retrobulbar venous plexus of mice by removing eyeball after the last challenge. The serum was isolated by centrifugation at 5,000 rpm for 20 min. The IgE and IL-4 were detected by ELISA with commercial reagent kits according to the manufacturers' instructions.




Figure 3 Expression and purification of Der f 28. (A) SDS-PAGE analysis of the protein expressed from the pET-44a(+)-Der f 28 recombinant plasmid in *E. coli* BL21 cells. Lane M, Protein Marker; Lane 1, *E. coli* BL21 transformed with pET-Der f 28 without IPTG induction; Lane 2, *E. coli* BL21 transformed with pET-Der f 28 with IPTG induction; (B) the peak figure of recombinant protein Der f 28 purified by molecular sieve chromatography. Longitudinal axis (mAu) for protein absorption peak; horizontal axis (mL) for volume; blue curve for protein absorption peak; gray curve for conductance value; (C) SDS-PAGE analysis of the final purified r-Der f 28 from *E. coli* BL21 cells. Lane M, Protein Marker; Lane 1, r-Der f 28 purified by molecular sieve chromatography.

The expression and purification of recombinant Der f 28 protein

In order to obtain r-Der f 28 protein for the subsequent experiments, we cloned Der f 28 cDNA into pET-44a(+), expressed and purified r-Der f 28. After the His-tag Der f 28 were successfully expressed by induction with IPTG (Figure 3A), the high purity of r-Der f 28 was obtained after

Table 1 The result of skin prick tests (SPT)

Subject	Gender/age	Diagnosis	Net wheal size (mm), level	
			DME	r-Der f 28
1	Female/9	AR, FA	7, +++	0
2	Female/72	BA, DA	2, +	2, +
3	Male/17	AR, BA	14, +++	0
4	Female/28	AR, BA	2, +	0
5	Male/62	AR, U, DA	2.25, +	0
6	Female/84	BA, FA	2.5, ++	0
7	Male/11	AR, BA	9, +++	2.5, +
8	Male/47	AR, BA	3.5, ++	0
9	Male/45	AR	2, +	0
10	Female/52	AR, FA	1.5, +	0
11	Male/20	BA, FA	1.5, +	0
12	Female/20	BA	2, ++	0
13	Female/13	AR	1.25, ++++	0
14	Female/66	AR, BA, DA	9, +++	4, ++
15	Female/64	BA	2.25, +	0
16	Female/41	AR, BA	3.5, ++	0
17	Female/44	AR, BA	4.5, +++	0
18	Male/15	BA	3, +	0
19	Female/24	AR, BA, U, FA, DA	8.5, +++	0
20	Female/10	AR, BA	8.5, +++	0
21	Female/18	AR, BA, FA, DA	10, +++	0
22	Male/54	AR, BA, U	1.5, +	0
23	Female/41	AR, BA, DA	10.5, ++++	0
24	Female/13	AR, BA	7, +++	0
25	Male/17	AR, BA, U, DA, FA	0.25, +	0
26	Female/31	U, DA	9.5, ++++	0

Positive, ≥ 1 ; Negative, 0. AR, allergic rhinitis; BA, bronchial asthma; U, urticaria; FA, food allergy; DA, drug allergy.

purification by affinity chromatography and molecular sieve chromatography (Figure 3B,C).

The allergic activity of r-Der f 28

We next determine the allergenicity of r-Der f 28. The results (Table 1) showed that 3/26 allergic patients had positive reactions to r-Der f 28, accounting for 11.5%. To determine the allergenicity of r-Der f 28, immunoblotting was performed using the serum from DME-allergic patients. The result demonstrated that the serum from allergic




Figure 4 Immunological characterization of *Dermatophagoides Farinae* Der f 28. (A) Immunoblotting analysis of specific IgE reactivity to allergen r-Der f 28 in the serum from r-Der f 28 positive individuals. Lanes 1, serum from allergic subjects; (B) the specific IgE reactivity to allergen r-Der f 28 by ELISA. P1-P3, the serum from r-Der f 28 positive patients; C1 and C2, the serum from healthy subjects. ELISA, enzyme-linked immunosorbent assay.

patients reacted to the r-Der f 28. IgE-binding bands with molecular weight were around 110 kD (Figure 4A), which was further confirmed by ELISA. Compared with the serum from healthy subjects, the IgE reactivity of r-Der f 28 from allergic patients increase significantly higher ($P < 0.01$) (Figure 4B). The results suggest that r-Der f 28 is an allergen in the dust mite allergen family.

Development of a mouse model of allergic asthma with r-Der f 28

An r-Der f 28-induced mouse model of allergic asthma was developed in our study (Figure 5A). As an important diagnostic index of asthma, airway hyperresponsiveness was assessed by methacholine-induced airflow obstruction using the FinePointe Whole-body Plethysmograph. As shown by Figure 5B, the airway hyperresponsiveness of the mice in the r-Der f 28 group is weaker than that in the HDM group, while the PBS group showed the lowest reaction. As shown by Figure 5C,D, the r-Der f 28 specific IgE and IL-4 were produced by the sensitized mice. Comparing with the normal group, those in the HDM group and the r-Der f 28 (** $P < 0.01$) group were significantly higher than that in the normal group. The total cells and eosinophils in BAL

were observed and counted with Wright Giemsa staining under an optical microscope. The proportion of eosinophils in HDM group and r-Der f 28 group was higher than that in control group (Figure 5E). The result was further confirmed by the histopathologic examination. In normal lung tissue, the contour of bronchial tube was clear and there was few inflammatory cell infiltration around the bronchial submucosa. In HDM group and r-Der f 28 group, a plenty of exudate collections in the airway lumen; the structure of the bronchial mucosa was partially damaged and inflammatory cell infiltration was apparent (Figure 5F-H). The results indicate that HDM extract and r-Der f 28 induce allergic pulmonary inflammation.

Discussion

Application of molecular biology technology to obtain recombinant allergens can provide new insight and new strategies for the development of specific immunotherapy for allergic diseases. Cumulative reports have demonstrated that it is diverse in the allergens of HDM, many mite allergens have not been identified and characterized (17). Heat-shock proteins (HSPs) are the most abundant and soluble intracellular proteins. In cell organisms, invertebrates and vertebrates, they perform a number of housekeeping functions that are essential for cell survival (18). In recent years, a growing number of studies have shown that HSPs play an important role in inflammatory processes of bronchial asthma, and has been reported to be an allergen in *Hordeum vulgare*, *Malassezia globosa* and so on (12-15).

In this study, through cloning, expression, and purification of full length recombinant heat shock protein (r-Der f 28), we identified that r-Der f 28 was an allergen in *D. Farinae*. Using SPT techniques, we proved that Der f 28 is an allergen of *D. farinae*. Der f 28 reacted to the sera from 11.5% of patients with *D. farinae* allergy (Table 1). The immunoblotting assays demonstrated that the serum IgE from allergic patients reacted to r-Der f 28 protein. The IgE reactivity of r-Der f 28 in the serum from the r-Der f 28 allergic patients increased by more than 8 folds compared with healthy subjects. Moreover, a well characterized mouse model of allergic airways disease induced by r-Der f 28 was developed successfully. Cytokines secreting by Th2 cells, such as IL-4, play an important role in the lung allergic inflammation (19). This study found that mice exposed to r-Der f 28 showed significantly higher levels of Th2 cytokine IL-4 and specific IgE. We also evaluated the effect of r-Der f 28 on AHR, a hallmark of asthma. Penh values




Figure 5 Establishment of mouse model of allergic asthma by induction with r-Der f28. (A) The protocol for mouse model of r-Der f 28-induced allergic asthma; (B) airway hyper-responsiveness were measured using the unrestrained whole-body plethysmography. The Penh values were recorded when the mice were stimulated with the increasing concentration of methacholine after the challenge; (C,D) the level of IL-4 and sIgE in Serum were detected by ELISA; (E) the number of eosinophil cell was counted in the BALF; (F-H) the pathologic changes of pulmonary tissue (hematoxylin-eosin stain, 100 \times). F, the PBS group; G, the r-Der f 28 group; H, the DME group; ELISA, enzyme-linked immunosorbent assay.

were assessed with whole-body plethysmography 24 h after the last r-Der f 28 challenge. Our result showed that the airway hyperresponsiveness of the mice in the r-Der f 28 group is weaker than that in HDM group, while PBS group showed the lowest reaction. Histological study show r-Der f 28 in asthma group induced extensive infiltration of inflammatory cells such as eosinophils around the alveoli, central bronchi and blood vessels, and marked goblet cell metaplasia and mucin production in epithelial cells.

In summary, the present data provide the evidence that Der f 28 is a new sub-group of HDM allergen, which may be useful for HDM allergy diagnosis and therapy.

Acknowledgements

Funding: This study was supported by grants from the Natural Science Foundation of China (No. 91442118 and 31400786), Guangdong Provincial University International-Hong Kong, Marco, Taiwan Scientific Innovation Projects (No. 2012gjh0009), Guangdong Engineering Technology Research Center Project (No. 2013158925), Shenzhen Scientific Technology Project International Cooperative Project (GJHZ20130408174112021), Shenzhen Scientific Technology Basic Research Projects (No. JCYJ20120613100657482, JCYJ20120613173233810,

JCYJ20140828163633992) and Shenzhen Nanshan District Research Development Project (No. KC2012JSYB0003A).

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

References

1. Arlian LG. House-dust-mite allergens: a review. *Exp Appl Acarol* 1991;10:167-86.
2. Platts-Mills TA, Thomas WR, Aalberse RC, et al. Dust mite allergens and asthma: report of a second international workshop. *J Allergy Clin Immunol* 1992;89:1046-60.
3. Tovey ER, Chapman MD, Platts-Mills TA. Mite faeces are a major source of house dust allergens. *Nature* 1981;289:592-3.
4. Vrtala S, Huber H, Thomas WR. Recombinant house dust mite allergens. *Methods* 2014;66:67-74.
5. An S, Shen C, Liu X, et al. Alpha-actinin is a new type of house dust mite allergen. *PLoS One* 2013;8:e81377.
6. Valenta R, Linhart B, Swoboda I, et al. Recombinant allergens for allergen-specific immunotherapy: 10 years anniversary of immunotherapy with recombinant allergens. *Allergy* 2011;66:775-83.
7. Thomas WR, Smith WA, Hales BJ. The allergenic specificities of the house dust mite. *Chang Gung Med J* 2004;27:563-9.
8. Van Der Veen MJ, Jansen HM, Aalberse RC, et al. Der p 1 and Der p 2 induce less severe late asthmatic responses than native *Dermatophagoides pteronyssinus* extract after a similar early asthmatic response. *Clin Exp Allergy* 2001;31:705-14.
9. Trombone AP, Tobias KR, Ferriani VP, et al. Use of a chimeric ELISA to investigate immunoglobulin E antibody responses to Der p 1 and Der p 2 in mite-allergic patients with asthma, wheezing and/or rhinitis. *Clin Exp Allergy* 2002;32:1323-8.
10. Meyer CH, Bond JF, Chen MS, et al. Comparison of the levels of the major allergens Der p I and Der p II in standardized extracts of the house dust mite, *Dermatophagoides pteronyssinus*. *Clin Exp Allergy* 1994;24:1041-8.
11. Le Mao J, Mayer CE, Peltre G, et al. Mapping of *Dermatophagoides farinae* mite allergens by two-dimensional immunoblotting. *J Allergy Clin Immunol* 1998;102:631-6.
12. Chiung YM, Lin BL, Yeh CH, et al. Heat shock protein (hsp 70)-related epitopes are common allergenic determinants for barley and corn antigens. *Electrophoresis* 2000;21:297-300.
13. Chuang JG, Su SN, Chiang BL, et al. Proteome mining for novel IgE-binding proteins from the German cockroach (*Blattella germanica*) and allergen profiling of patients. *Proteomics* 2010;10:3854-67.
14. Ishibashi Y, Kato H, Asahi Y, et al. Identification of the major allergen of *Malassezia globosa* relevant for atopic dermatitis. *J Dermatol Sci* 2009;55:185-92.
15. Aki T, Fujikawa A, Wada T, et al. Cloning and expression of cDNA coding for a new allergen from the house dust mite, *Dermatophagoides farinae*: homology with human heat shock cognate proteins in the heat shock protein 70 family. *J Biochem* 1994;115:435-40.
16. Chang YS, Kim YK, Jeon SG, et al. Influence of the Adjuvants and Genetic Background on the Asthma Model Using Recombinant Der f 2 in Mice. *Immune Netw* 2013;13:295-300.
17. Li GF, Wang Y, Zhang ZS, et al. Identification of immunodominant Th1-type T cell epitopes from *Schistosoma japonicum* 28 kDa glutathione-S-transferase, a vaccine candidate. *Acta Biochim Biophys Sin (Shanghai)* 2005;37:751-8.
18. Srivastava P. Roles of heat-shock proteins in innate and adaptive immunity. *Nat Rev Immunol* 2002;2:185-94.
19. Broide DH, Stachnick G, Castaneda D, et al. Immunostimulatory DNA mediates inhibition of eosinophilic inflammation and airway hyperreactivity independent of natural killer cells in vivo. *J Allergy Clin Immunol* 2001;108:759-63.

Cite this article as: Lin JL, Wang YY, Xiao XJ, Wu YL, Sun BQ, Gao AJ, Liu ZG, Li J, Yang PC, Liu XY. Characterization of a new subtype of allergen in *dermatophagoides farinae*—Der f 28. *J Thorac Dis* 2015;7(10):1842-1849. doi: 10.3978/j.issn.2072-1439.2015.10.43