



Clinical characteristics and determinants of mortality in coronavirus disease 2019 (COVID-19) patients on an intensive care unit—a retrospective explorative 1-year all-comers study

Wiebke Brücker¹, Amir Abbas Mahabadi², Annette Hüschen¹, Jan Becker³, Sebastian Daehnke³, Stefan Möhlenkamp^{1^}

¹Clinic of Cardiology and Intensive Care Medicine, Bethanien Krankenhaus Moers, Moers, Germany; ²Department of Cardiology and Vascular Medicine, West German Heart and Vascular Center, University Hospital Essen, University Medicine Essen, Essen, Germany; ³Clinic of Anaesthesiology and Intensive Care Medicine, Bethanien Krankenhaus Moers, Moers, Germany

Contributions: (I) Conception and design: W Brücker, AA Mahabadi, S Daehnke, S Möhlenkamp; (II) Administrative support: AA Mahabadi, S Daehnke, S Möhlenkamp; (III) Provision of study materials or patients: W Brücker, A Hüschen, J Becker, S Daehnke, S Möhlenkamp; (IV) Collection and assembly of data: W Brücker, A Hüschen, J Becker, S Daehnke, S Möhlenkamp; (V) Data analysis and interpretation: W Brücker, AA Mahabadi, S Möhlenkamp; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Stefan Möhlenkamp, MD, FACC, FESC. Clinic of Cardiology and Intensive Care Medicine, Bethanien Krankenhaus Moers, Bethanienstrasse 21, 47441 Moers, Germany. Email: stefan.moehlenkamp@bethanienmoers.de.

Background: Clinical outcome in patients with coronavirus disease 2019 (COVID-19) requiring treatment on intensive care units (ICU) remains unfavourable. The aim of this retrospective study was to exploratively identify potential predictors of unfavourable outcome in ICU patients diagnosed with COVID-19.

Methods: In all patients with COVID-19 (n=50) or severe acute respiratory syndrome coronavirus type 2 (SARS-CoV-2) as comorbidity (n=11) at our ICU we assessed clinical, respiratory and laboratory parameters with a potential role for outcome. Main outcome variables were intubation and mortality rates.

Results: Between March 2020 and March 2021, 573 patients were hospitalized with SARS-CoV-2 infection. Of these, 61 patients (10.6%, 44.3% women) aged 66.4±13.3 were admitted to ICU. A proportion of 73.8% of patients had moderate or severe acute respiratory distress syndrome (ARDS). COVID-19 patients differed clinically from those with SARS-CoV-2 as comorbidity, such as severe heart or renal failure or sepsis as the leading cause of ICU admission, despite similar mortality rates (44.0% vs. 45.5%, P>0.5). Among COVID-19 patients, those who died had more often severe ARDS (91% vs. 46%, P=0.001), longer non-invasive ventilation (NIV) therapy prior to ICU (6.3±5.9 vs. 2.5±2.0 days, P=0.046), and higher interleukin-6 (IL-6) and lactate dehydrogenase (LDH) values as compared to survivors. In multivariable analysis, NIV duration ≥5 days on admission [odds ratio (OR): 42.20, 95% confidence interval (CI): 1.22 to >99, P=0.038] and IL-6 [OR: 4.08, 95% CI: 1.16–14.33, P=0.028] remained independently predictive of mortality. In worsening tertiles of partial pressure of oxygen (pO₂)/inspiratory oxygen fraction (FiO₂) on admission (≥161.5, 96.5 to <161.5, <96.5) we observed a stepwise increase in intubation rates (P=0.0034) and mortality rates (P=0.031).

Conclusions: As inflammation, ARDS severity and longer NIV duration prior to ICU are associated with intubation and mortality rates, prognosis appears to be largely determined by disease severity. Whether NIV aggravates ARDS or if it indicates lack of recovery independent from type of ventilation, or both should be clarified in a prospective trial.

Keywords: Coronavirus disease 2019 (COVID-19); non-invasive ventilation (NIV); invasive ventilation; intubation; mortality

[^] ORCID: 0000-0002-1490-7979.

Submitted Oct 29, 2021. Accepted for publication Apr 07, 2022.

doi: 10.21037/jtd-21-1713

View this article at: <https://dx.doi.org/10.21037/jtd-21-1713>

Introduction

Infection with severe acute respiratory syndrome coronavirus type 2 (SARS-CoV-2) may cause coronavirus disease 2019 (COVID-19) and is associated with a high mortality in patients admitted to intensive care units (ICU) (1,2). Mortality risk has been linked with several parameters, including advanced age, male gender, body mass index (BMI), multi-organ failure, and associated laboratory abnormalities (3-5). Care of these critically ill patients is challenging (6,7). The goal of therapy is mainly to avoid complications, to prevent disease deterioration and to support recovery (1,2). Regarding respiratory support, the use of non-invasive ventilation (NIV) has evolved during the 1st and 2nd wave of the pandemic (8,9), with initially high rates of early intubation partly due to concerns of viral transmission to physicians and health care staff. At our hospital, NIV has early on been proposed as the main type of respiratory support to avoid intubation and mechanical ventilation (10,11). Therefore, many patients admitted to our hospital were initially referred to a "COVID-19 intermediate care unit" with high rates of NIV (10,11), unless they had concomitant diseases other than COVID-19 that necessitated ICU therapy. This led to identification of two distinct groups of patients, i.e., those with COVID-19 and those with SARS-CoV-2 infection as comorbidity, e.g., with sepsis, severe heart or renal failure, etc., as the leading cause for ICU admission. In some patients, respiratory function deteriorated and they were escalated to invasive mechanical ventilation (IMV) according to guidelines (1,12,13). Despite all efforts, mortality rates appeared to be high. It is therefore the aim of this exploratory study to examine the clinical course of patients admitted to our ICU in this setting and to identify predictors of intubation and mortality. We present the following article in accordance with the STROBE reporting checklist (available at <https://jtd.amegroups.com/article/view/10.21037/jtd-21-1713/rc>).

Methods

Study design, data collection and ethics

We conducted a single-center retrospective observational

study that per protocol included all patients admitted to our ICU between March 2020 and March 2021 (*Figure 1*). All parameters reported in this study were measured as part of clinical routine. We searched our hospital database for all patients that were coded as having SARS-CoV-2 infection and that were admitted to our ICU. A SARS-CoV-2 infection was determined by polymerase chain reaction (PCR) and supported by clinical criteria, laboratory values, and chest computed tomography (CT). We distinguished patients admitted due to COVID-19, from those admitted with SARS-CoV-2 infection as comorbidity, i.e., primarily due to acute heart failure, renal failure, etc. without direct relation to SARS-CoV-2. Precautions of viral transmission and the same medical protocol were applied in all patients. The study complies with the Declaration of Helsinki (as revised in 2013), and was approved by the ethics committee of University Duisburg-Essen (#21-9911-BO). Due to the retrospective nature of the study and blinding of study data, the need for informed consent was waived.

Inclusion and exclusion criteria and admission criteria to ICU

All patients transferred to our ICU entered the database (*Figure S1*). We excluded one patient who was transferred to our clinic for weaning from intubation after recovery from COVID-19. Patients that acquired SARS-CoV-2 at our hospital after ICU stay were also excluded from the database. No other patient was excluded. Patients were admitted to ICU using established criteria such as hemodynamic or metabolic instability, including elevated troponin, N-terminal pro-B-type natriuretic peptide (NT-proBNP), and/or D-dimer values, sepsis, renal failure, reduced Glasgow coma scale (GCS), etc. Hypoxemic awake COVID-19 patients were transferred to ICU in case of respiratory failure with CO₂-elevation despite NIV-therapy, or complications such as subcutaneous or mediastinal emphysema.

Study definitions and measurements

COVID-19 was present when patients developed respiratory failure and other symptoms typically associated with




Figure 1 Incidence in the district of Wesel, federal state of North-Rhine Westfalia (green line), and patients hospitalized with SARS-CoV-2 infection or COVID-19 (blue bars) and admitted to ICU (red bars) between 03/2020 and 03/2021. SARS-CoV-2, severe acute respiratory syndrome coronavirus type 2; COVID-19, coronavirus disease 2019; ICU, intensive care unit.

COVID-19 and required therapy. ARDS was determined as per Berlin criteria (14). Accordingly, ARDS categories are based upon worst partial pressure of oxygen (pO_2)/inspiratory oxygen fraction (FiO_2) values despite maximum respiratory support measured during ICU stay: mild: $pO_2/FiO_2 = 201$ – 300 mmHg, moderate: $pO_2/FiO_2 = 101$ – 200 mmHg, and severe: $pO_2/FiO_2 \leq 100$ mmHg. NIV was coded irrespective of interface, mode and ventilator type employed. On ICU, ventilator settings were modified clinically depending on blood gas analysis and work of breathing. In patients on IMV, pO_2 and FiO_2 were measured. In patients on NIV (Stellar 100 or 150, Resmed, Germany or Evita XL or V600, Dräger, Germany), pO_2 was measured. FiO_2 was estimated based on established tabulation for converting oxygen insufflation to FiO_2 (15–17) (see Table S1). Duration of NIV in days prior to ICU was determined based on date of first initiation of NIV during hospital admission until date of ICU admission. In patients with oxygen insufflation using oxygen cannula (ASID BONZ, Germany), FiO_2 was estimated from O_2 -flow using established tabulation (18,19) (see Table S2). pO_2 was measured by blood gas analysis. In few patients without arterial line, pO_2 was estimated from pulse oximetry-based O_2 -saturation and converted to pO_2 using tabulation (see Table S3). Septic shock was defined according to the 2016 Third International Consensus Definition for Sepsis and Septic Shock (20). Sepsis organ failure assessment (SOFA) score and Charlson Comorbidity Index (CCI) were calculated as published (21–23).

COVID-19 therapy

Standard therapy comprised i.v. dexamethasone in COVID-19 ARDS after May 2020. Anticoagulation regimen was modified according to recommendations considering comorbidities. High-flow nasal oxygen—if applied—was used alternating with NIV. Prone positioning was routinely applied in intubated patients with $pO_2/FiO_2 < 150$ mmHg and was repeated in responders. Awake NIV patients were placed in prone position depending on pO_2/FiO_2 and clinical judgement. Crystalloids were given restrictively and diuretics applied as hemodynamically tolerated. A macrolide was routinely prescribed and additional antibiotics given as needed. Intravenous opioids were administered to facilitate NIV tolerance as needed. National COVID-19 guidelines were applied in all ICU patients (1,12,13). Conscious vigilant normocapnic patients without need for catecholamines usually continued to receive NIV for initial respiratory support when admitted to ICU. Intubation and further escalation to extracorporeal membrane oxygenation (ECMO) therapy was a clinical decision of the treating physician.

Comorbidities, complications, follow-up

Comorbidities were defined as present, when being established prior to admission on ICU. Specifically, chronic kidney disease was defined as repeatedly measured glomerular filtration rate (GFR) < 60 mL/min prior to present hospitalization. Chronic heart failure was defined

as known left ventricular ejection fraction <50% or pre-existing heart failure diagnosis and medication. Chronic pulmonary disease was summarized by asthma, chronic obstructive pulmonary disease (COPD), silicosis, and asbestosis. Diabetes was defined as HbA1c \geq 6.5% or antidiabetic medication. Hypertension was defined by previous prescription of antihypertensive medication or pre-existing diagnosis of hypertension. Atherosclerotic cardiovascular disease (ASCVD) summarized as established diagnoses of coronary artery disease, peripheral artery disease or cerebrovascular disease. Additional comorbidities had all been diagnosed prior to ICU admission and are listed in Table S4.

Complications were determined using established clinical criteria and include encephalopathy, increasing confusion and/or seizure; pulmonary embolus or other thromboembolic events including intestinal ischemia; septic shock, disseminated intravascular coagulopathy (DIC); acute renal failure; myocardial infarction; acute heart failure; pneumothorax; mediastinal or cutaneous emphysema; and cardio-pulmonary resuscitation (CPR).

For all-cause in-hospital mortality, database was closed after all included patient had either died or left hospital alive, including patients that had been transferred for ECMO therapy to tertiary specialized ECMO centers.

Statistical analysis

Continuous variables are reported as mean \pm standard deviation (SD) or median and interquartile range (IQR). Discrete variables are given in frequency and percentiles. Baseline characteristics are specified for the overall cohort as well as stratified by patients admitted due to COVID-19 infection *vs.* patients with SARS-CoV-2 infection as comorbidity. In the subgroup of patients admitted due to COVID-19, continuous variables were compared using two-sided *t*-test or Man-Whitney U test and discrete variables using a two-sided Fischer's exact test. Subgroups of patients with and without intubation and with and without mortality were compared using identical statistical methods in an exploratory analysis. Patients with missing values in subgroups analyses were excluded from these analyses. Univariate and multivariable logistic regression analyses were performed for the association of clinical characteristics with mortality. As lactate dehydrogenase (LDH) and interleukin-6 (IL-6) were not normally distributed, logarithmic transformation was performed. For continuous variables, odds ratio (OR) and confidence intervals (CI) were

depicted per one SD increase. Multivariable model included all variables with significant association in univariate analysis. Given the high co-linearity, severe ARDS and intubation could not be included within the same model. Therefore, we provided separate analyses, including either severe ARDS or intubation into the model. All analyses were performed using SAS software (Version 9.4, SAS Institute Inc.). A *P* value of <0.05 indicated statistical significance.

Results

Study population

During the first year of the pandemic, *i.e.*, between March 2020 and March 2021, 573 patients had been hospitalized and coded SARS-CoV-2-positive at any time during hospital stay. Of these, 61 patients (10.6%, 44.3% women) aged 66.4 ± 13.3 (range, 17–92 years) were admitted to ICU. During the first (03/2020–09/2020) and second 6 months (10/2020–03/2021), *n*=11 (18%) and *n*=50 (82%) patients respectively, were admitted. Clinical characteristics and laboratory findings are shown in Tables 1,2. On admittance, ARDS was present in 79% of patients, with more than half of patients having severe ARDS. Lowest pO_2/FiO_2 during ICU stay averaged 123 ± 89 mmHg. Comorbidities were present in 56 patients (92%). Half of the patients were intubated and overall mortality was 44.3% (Table 1). Maximum duration on ICU was 75 days.

Intubation, complications and mortality on ICU in the entire cohort

One patient had been intubated for 10 days prior to being transferred to our clinic. IMV was initiated at our ICU in *n*=30 patients (49.2%), *i.e.*, on day 1 in *n*=9 patients (30.0%), on days 2–4 in *n*=13 (43.3%) and on day \geq 5 in *n*=8 (26.7%) patients. Three patients were intubated on days 13, 15, and 20 at our ICU. pO_2/FiO_2 just prior to intubation averaged 94 ± 65 mmHg. While on ICU, 43 patients (70.5%) experienced complications, mostly encephalopathy/confusion, heart failure, acute renal failure, or septic shock (Table 1). CPR was necessary in 14 patients (23.0%). Seven patients died with a do not intubate/do not resuscitate (DNI/DNR)-order, four with COVID-19 and three with SARS-CoV-2. These patients were included in the analysis because they initially wished to be and were treated, and DNI/DNR was consented at end stage disease. Mortality in the entire cohort was 44.3% (Table 1). Mortality on IMV in the entire

Table 1 Clinical characteristics of patients admitted to ICU due to COVID-19 or with SARS-CoV-2 as a comorbidity

Variables	All patients
N (%)	61 (100.0)
Age (years)	66.4±13.3
Sex, n (% women)	27 (44.3)
BMI (kg/m ²)	31.1±7.4
Any comorbidity (n (%))	56 (91.8)
Hypertension	34 (55.7)
Diabetes	27 (44.3)
ASCVD	26 (42.6)
Chronic kidney disease	11 (18.0)
SOFA score on admission	5.6±2.9
SOFA score after 24 h	7.0±3.8
Charlson Comorbidity Index	3.9±2.4
pO ₂ /FiO ₂ (mmHg)	
On admission	178±115
After 24 h	158±93
Lowest value	123±89
Any ARDS, n (%)	48 (78.7)
Mild	3 (4.9)
Moderate	12 (19.7)
Severe	33 (54.1)
Any complication, n (%)	43 (70.5)
CPR	14 (23.0)
Encephalopathy/confusion	17 (27.9)
Heart failure	14 (23.0)
Acute renal failure	12 (19.7)
Septic shock	10 (16.4)
Length of ICU stay, median days (IQR)	6 (2; 14)
Intubation, n (%)	31 (50.8)
Mortality, n (%)	27 (44.3)

Data were presented as mean ± standard deviation if not otherwise specified. ICU, intensive care unit; COVID-19, coronavirus disease 2019; SARS-CoV-2, severe acute respiratory syndrome coronavirus type 2; BMI, body mass index; ASCVD, atherosclerotic cardiovascular disease; SOFA, sepsis organ failure assessment; pO₂, partial pressure of oxygen; FiO₂, inspiratory oxygen fraction; ARDS, acute respiratory distress syndrome; CPR, cardio-pulmonary resuscitation; IQR, interquartile range.

Table 2 Main laboratory findings on admission in patients admitted to ICU due to COVID-19 or with SARS-CoV-2 as a comorbidity

Variables (units)	Normal range	All patients
N (%)	–	61 (100.0)
Lymphocytes (%)	20–40	11.9±9.3
CRP (mg/dL)	<0.5	13.2±9.1
PCT (ng/mL)	<0.5	0.40 (0.15; 0.85)
IL-6 (pg/mL)	<7	148 (55; 303)
D-dimer (µg/mL)	<0.5	3.5±3.9
hs-troponin (pg/mL)	<14	41.0 (21.5; 102.5)
NT-proBNP (pg/mL)	<125	1,702 (422; 4,430)
LDH (U/L)	<250	486 (412; 648)
Creatinine (mg/dL)	0.5–0.9	1.7±1.8

Data were presented as mean ± standard deviation or median days (IQR) if not otherwise specified. ICU, intensive care unit; COVID-19, coronavirus disease 2019; SARS-CoV-2, severe acute respiratory syndrome coronavirus type 2; CRP, C-reactive protein; PCT, procalcitonin; IL-6, interleukin-6; hs, high sensitive; NT-proBNP, N-terminal pro-B-type natriuretic peptide; LDH, lactate dehydrogenase; IQR, interquartile range.

cohort was 64.5% (n=20/31).

Patients admitted due to COVID-19 or with SARS-CoV-2 as comorbidity

We identified 50 patients (82%) that were admitted because of COVID-19, while 11 patients (18%) were admitted with SARS-CoV-2 as comorbidity (*Table 3*). The two sub-groups differed significantly in several clinically important aspects: patients admitted due to COVID-19 had lower CCIs, lower pO₂/FiO₂, and higher rates of ARDS and of intubation (*Table 3*). Mortality rates, SOFA scores and duration on ICU were similar (*Table 3*). Moreover, patients admitted due to COVID-19 had higher LDH and CRP values, and lower lymphocytes (*Table 4*). Among patients admitted due to COVID-19, the rate of intubation and mortality increased with increasing ARDS severity (*Tables 5,6*). In contrast, only 27% of patients with SARS-CoV-2 as comorbidity were intubated (*Table 3*).

Rates and predictors of intubation and mortality in COVID-19 patients

Intubation and mortality rates are given in *Tables 5,6*. They

Table 3 Clinical characteristics in patients admitted to ICU due to COVID-19 and with SARS-CoV-2 as a comorbidity

Variables	Patients admitted due to COVID-19	Patients with SARS-CoV-2 as comorbidity	P value
N (%)	50 (82.0)	11 (18.0)	–
Age (years)	66.3±13.4	67.2±13.7	0.83
Sex, n (% women)	23 (46.0)	4 (36.4)	0.74
BMI (kg/m ²)	31.4±7.4	30.1±7.7	0.61
Any comorbidity, n (%)	45 (90.0)	11 (100.0)	0.57
Hypertension	29 (58.0)	5 (45.5)	0.51
Diabetes	23 (46.0)	4 (36.4)	0.74
ASCVD	17 (34.0)	9 (81.8)	0.006
Chronic kidney disease	5 (10.0)	6 (54.5)	0.003
SOFA score on admission	5.6±2.7	5.5±3.7	0.92
SOFA score after 24 h	7.1±3.7	6.6±3.8	0.69
Charlson Comorbidity Index	3.7±2.2	5.2±2.6	0.051
pO ₂ /FiO ₂ (mmHg)			
On admission	151±93	302±127	<0.0001
After 24 h	137±72	253±118	0.009
Lowest value	101±58	126±129	0.009
Any ARDS, n (%)	45 (90.0)	3 (27.3)	<0.0001
Mild	3 (6.0)	0 (0)	1.00
Moderate	9 (18.0)	3 (27.3)	0.68
Severe	33 (66.0)	0 (0)	<0.0001
Any complication, n (%)	35 (70.0)	8 (72.7)	1.00
CPR	12 (24.0)	2 (18.2)	1.00
Encephalopathy/confusion	13 (26.0)	4 (36.4)	0.48
Heart failure	10 (20.0)	4 (36.4)	0.26
Acute renal failure	8 (16.0)	4 (36.4)	0.20
Septic shock	6 (12.0)	4 (36.4)	0.07
Length of ICU stay, median (IQR) (days)	5.5 (2; 14)	7 (2; 19)	0.99
Intubation, n (%)	28 (56.0)	3 (27.3)	0.11
Mortality, n (%)	22 (44.0)	5 (45.5)	1.00

Data were presented as mean ± standard deviation if not otherwise specified. ICU, intensive care unit; COVID-19, coronavirus disease 2019; SARS-CoV-2, severe acute respiratory syndrome coronavirus type 2; BMI, body mass index; ASCVD, atherosclerotic cardiovascular disease; SOFA, sepsis organ failure assessment; pO₂, partial pressure of oxygen; FiO₂, inspiratory oxygen fraction; ARDS, acute respiratory distress syndrome; CPR, cardio-pulmonary resuscitation; IQR, interquartile range.

increased with increasing ARDS severity as estimated by pO₂/FiO₂ (Figure 2). Among intubated patients with COVID-19, 64.3% (n=18/28) died, including all nine

patients on ECMO or iLA-active (Xenios, Germany).

Patients who were intubated during ICU stay, had more often severe ARDS, lower pO₂/FiO₂ values on admission,

Table 4 Main laboratory findings on admission in patients admitted to ICU due to COVID-19 and with SARS-CoV-2 as a comorbidity

Variables (units)	Normal range	Patients admitted due to COVID-19	Patients with SARS-CoV-2 as comorbidity	P value
N (%)	–	50 (82.0)	11 (18.0)	–
Lymphocytes (%)	20–40	10.8±8.6	19.0±11.6	0.028
CRP (mg/dL)	<0.5	14.6±9.3	6.3±4.1	0.0001
PCT (ng/mL)	<0.5	0.4 (0.2; 0.8)	0.4 (0.1; 4.0)	0.89
IL-6 (pg/mL)	<7	147 (55; 355)	224 (37; 267)	0.80
D-dimer (µg/mL)	<0.5	3.5±4.0	4.0±3.6	0.77
hs-troponin (pg/mL)	<14	38.0 (20.5; 92.5)	63.5 (25.0; 132.0)	0.41
NT-proBNP (pg/mL)	<125	1,681 (426; 3,965)	3,664 (263; 7,326)	0.58
LDH (U/L)	<250	538 (424; 688)	281 (199; 405)	0.004
Creatinine (mg/dL)	0.5–0.9	1.7±1.9	1.7±0.9	0.98

Data were presented as mean ± standard deviation or median days (IQR) if not otherwise specified. ICU, intensive care unit; COVID-19, coronavirus disease 2019; SARS-CoV-2, severe acute respiratory syndrome coronavirus type 2; CRP, C-reactive protein; PCT, procalcitonin; IL-6, interleukin-6; hs, high sensitive; NT-proBNP, N-terminal pro-B-type natriuretic peptide; LDH, lactate dehydrogenase; IQR, interquartile range.

after 24 h and at any time during ICU stay, and had slightly higher SOFA-scores after 24 h. These patients had also more often non-invasive respiratory support prior to ICU-admission than patients that were not intubated (Tables 5,6). CCI, duration of NIV and duration of hospital stay prior to ICU admission were slightly higher and longer (Tables 5,6). Intubated patients also differed in several laboratory findings in that they had higher IL-6 and LDH values on admission, but slightly lower NT-proBNP values (Tables 5,6).

Patients that died had more often severe ARDS (91%) and had lower pO_2/FiO_2 values on admission, after 24 h and lower values at any time of ICU stay than survivors (Tables 5,6). Patients that died were about 5 years older, and had been hospitalized longer, had more often NIV-therapy and longer duration of NIV therapy prior to ICU admission (Tables 5,6). Likewise, among patients who were escalated to intubation, those that died had longer total days on NIV, i.e., prior to ICU and in addition during ICU stay, than those who survived (7.8 ± 5.6 vs. 4.1 ± 3.0 days, $P=0.063$). Patients that died also differed in several laboratory findings: again, IL-6- and LDH values were much higher, and NT-proBNP values were slightly lower.

In univariate analysis, LDH, IL-6, presence of severe ARDS, intubation, and duration of NIV ≥ 5 days prior to ICU were associated with mortality. In multivariable logistic regression analysis, IL-6 and ≥ 5 days of NIV prior to ICU

remained independent predictors of mortality (Table 7), but ORs for severe ARDS and LDH remained about 3-fold elevated.

Discussion

At our institution, 10.6% of patients with COVID-19 or SARS-CoV-2 infection were admitted to ICU. Among COVID-19 patients on ICU, intubation rate was 57%, and 45% of patients died. These numbers are in line with previous reports as summarized in current guidelines (1,3). Roedel *et al.* reported a comparatively favorable overall ICU mortality of 35% in patients included until early June 2020 in a German cohort, but outcome was not reported for ARDS categories separately (24). Mortality of intubated patients was 44% (24), again lower compared to our study (i.e., 64.3%), possibly attributable to later intubation in our cohort of patients with mostly severe ARDS.

In our retrospective explorative study, severe ARDS estimated by pO_2/FiO_2 , (hyper-)inflammation reflected in elevated IL-6, and long disease duration and poor recovery from respiratory failure evidenced by long NIV-therapy prior to and during ICU stay were associated with intubation and death. Within the time course of clinical SARS-CoV-2 manifestation from early viral infection to severe hyperinflammatory ARDS (25), our intubation and mortality rates are subject to confounding by indication but

Table 5 Determinants of intubation and mortality in patients admitted to ICU due to COVID-19

Variables	All Patients	Intubation			Survival status		
		Yes	No	P value	Died	Survived	P value
N (%)	50 (100.0)	28 (56.0)	22 (44.0)	–	22 (44.0)	28 (56.0)	–
Age (years)	66.3±13.4	66.3±10.4	66.2±16.7	0.99	69.1±12.0	64.1±14.2	0.19
Sex, n (% women)	23 (46.0)	8 (28.6)	15 (68.2)	0.01	8 (36.4)	15 (53.6)	0.26
BMI (kg/m ²)	31.4±7.4	31.7±7.2	30.9±7.9	0.70	32.6±7.8	30.4±7.1	0.29
ARDS, n (%)							
No ARDS	5 (10.0)	0	5 (22.7)	0.001	0	5 (17.9)	0.059
Mild ARDS	3 (6.0)	1 (3.6)	2 (9.1)	0.58	0	3 (10.7)	0.25
Moderate ARDS	9 (18.0)	1 (3.6)	8 (36.4)	0.007	2 (9.1)	7 (25.0)	0.27
Severe ARDS	33 (66.0)	26 (92.9)	7 (31.8)	<0.0001	20 (90.9)	13 (46.4)	0.001
pO ₂ /FiO ₂ (mmHg)							
On admission	151±93	114±68	198±102	0.002	107±43	185±108	0.001
After 24 h	137±72	114±46	167±88	0.015	104±32	164±84	0.001
Lowest value	101±58	79±40	128±66	0.004	75±22	120±70	0.003
SOFA score							
On admission	5.6±2.9	6.2±2.7	4.8±2.5	0.06	6.2±2.8	5.1±2.5	0.17
After 24 h	7.0±3.8	8.1±3.5	5.9±3.7	0.04	8.0±3.5	6.4±3.8	0.12
CCI	3.9±2.4	3.7±2.1	3.6±2.5	0.85	4.1±2.3	3.3±2.2	0.18
NIV prior to ICU, n (%)	23 (46.0)	16 (57.1)	7 (31.8)	0.09	13 (59.1)	10 (35.7)	0.15
Days NIV prior to ICU	4.7±4.9	5.1±5.4	3.6±3.7	0.35	6.3±5.9	2.5±2.0	0.046
Days NIV ≥5 prior to ICU, n (%)	7 (14.0)	5 (17.9)	2 (9.1)	0.44	6 (27.3)	1 (3.6)	0.035
Hospitalization prior to ICU, median (IQR) (days)	1 (0; 4)	2 (0; 4)	0 (0; 5)	0.32	2 (0; 7)	0 (0; 3)	0.083

Data were presented as mean ± standard deviation if not otherwise specified. ICU, intensive care unit; COVID-19, coronavirus disease 2019; BMI, body mass index; ARDS, acute respiratory distress syndrome; pO₂, partial pressure of oxygen; FiO₂, inspiratory oxygen fraction; SOFA, sepsis organ failure assessment; CCI, Charlson Comorbidity Index; NIV, non-invasive ventilation; IQR, interquartile range.

may also in part reflect comparatively long NIV-therapy in our patient cohort.

ARDS severity and IL-6 have early on been established as being prognostically relevant (3). Our study extends these findings in that long duration on NIV prior to ICU admission was also associated with mortality and may need to be added to the list of prognostically relevant parameters. Karagiannidis *et al.* report a mortality of 50% in patients that were initially treated with NIV, which is similar to the 52% mortality observed on IMV (8). This finding is consistent with a meta-analysis in almost 9,000 critically ill COVID-19 patients, where use of high flow

nasal oxygen (HFNO) or NIV and timing of intubation had little effect if any on morbidity and mortality (26). However, Karagiannidis *et al.* also observed that mortality increased continuously the longer patients were on NIV before requiring IMV (8). Mortality was as high as 75% in patients on NIV for 5 days or longer prior to intubation, which is consistent with our findings. Unfortunately, markers of inflammation and indices of ARDS severity were not reported. A caution regarding NIV in moderate or severe ARDS also derives from the Lung Safe Study, where patients on NIV with pO₂/FiO₂ <150 mmHg had a higher ICU mortality compared to patients on IMV (27).

Table 6 Laboratory values on admission in patients with COVID-19 by intubation and mortality

Variables	All patients	Intubation	No intubation	P value	Patients died	Patients survived	P value
Lymphocytes (%)	10.8±8.6	8.9±8.3	13.1±8.6	0.10	9.4±8.4	11.9±8.7	0.33
CRP (mg/dL)	14.6±9.3	16.4±9.5	12.3±8.6	0.13	14.5±7.0	14.6±10.8	0.96
PCT (ng/mL)	0.4 (0.2; 0.8)	0.4 (0.2; 0.9)	0.4 (0.2; 0.8)	0.83	0.4 (0.2; 0.6)	0.4 (0.2; 0.9)	0.29
IL-6 (pg/mL)	147 (55; 355)	183 (89; 487)	99 (33; 175)	0.021	191 (139; 484)	93 (26; 183)	0.009
D-dimer (µg/mL)	3.5±4.0	3.8±3.2	3.0±4.9	0.52	3.7±3.1	3.3±4.6	0.77
hs-troponin (pg/mL)	38 (21; 93)	34 (22; 72)	57 (14; 149)	0.76	35 (25; 61)	51 (12; 160)	0.82
NT-proBNP (pg/mL)	1,681 (426; 3,965)	1,301 (379; 3,368)	1,851 (469; 5,517)	0.09	1,082 (279; 2,886)	1,851 (727; 5,517)	0.09
LDH (U/L)	538 (424; 688)	629 (449; 771)	470 (410; 545)	0.026	640 (537; 796)	449 (416; 607)	0.008
Creatinine (mg/dL)	1.7±1.8	1.6±2.0	1.9±1.9	0.56	1.4±1.3	2.0±2.3	0.29

Data were presented as mean ± standard deviation or median days (IQR) if not otherwise specified. COVID-19, coronavirus disease 2019; CRP, C-reactive protein; PCT, procalcitonin; IL-6, interleukin-6; hs, high sensitive; NT-proBNP, N-terminal pro-B-type natriuretic peptide; LDH, lactate dehydrogenase; IQR, interquartile range.




Figure 2 Intubation rates and mortality rates stratified by tertiles of pO₂/FiO₂ (mmHg) on admission in patients with COVID-19. pO₂, partial pressure of oxygen; FiO₂, inspiratory oxygen fraction; COVID-19, coronavirus disease 2019.

Recently, Wendel Garcia *et al.* studied outcome in different strategies of early respiratory support in critically ill COVID-19 patients. Their data also suggest that NIV should be avoided due to an elevated ICU mortality risk (28). In contrast, Daniel *et al.* report a reduced mortality when NIV was employed as the initial intervention in COVID-19 patients (29). Yet again, patients that were escalated to intubation had a (non-significant) 39% increase in mortality compared to patients that were initially intubated (29). Our data thus add to the currently limited and heterogeneous evidence on NIV in moderate to severe COVID-19 ARDS.

There are at least two possible explanations for our findings. First, NIV may have aggravated disease progression in our patients attributable to “patient self-inflicted lung injury” (P-SILI) (30,31). This concept suggests that increased respiratory drive and breathing effort induces lung injury due to uncontrolled swings in transpulmonary pressure and hence increase lung stress in the aerated compartment of the baby lung. The positive end-expiratory pressure (PEEP) that is applied during NIV may not prevent injury caused by recurrent alveolar (hyper-)inflation and deflation (32). In addition, a marked decrease in pleural pressure may increase vascular transmural pressure and vascular permeability, contributing to alveolar and interstitial pulmonary oedema (32), beyond the increased vascular permeability caused by SARS-CoV-2 itself (1,2). Especially patients with pO₂/FiO₂ <200 mmHg may be at risk of NIV failure (16,32), which is consistent with our findings. On the other hand, Tobin *et al.* argue in a series of publications that there is insufficient evidence for the concept of P-SILI (33-35). They question its role for the progression of respiratory failure in COVID-19 and caution against pre-emptive liberal use of intubation and mechanical ventilation (33-35). Indeed, pressure support via NIV reduces respiratory effort, relieves dyspnea, and ameliorates oxygenation, especially in high PEEP settings (32,36), but prospective outcome studies in COVID-19 patients comparing NIV and IMV are limited.

Second, the observation of impaired prognosis in patients being longer on NIV may indicate poor recovery from ARDS. Escalation of respiratory support from NIV to IMV is usually referred to as “NIV-failure”. The

Table 7 Univariate and multivariable associations of variables with mortality in patients with COVID-19

Variables	Univariate		Multivariable Model 1		Multivariable Model 2	
	OR (95% CI)	P value	OR (95% CI)	P value	OR (95% CI)	P value
Age (per 1 SD)	1.51 (0.81–2.80)	0.16	–	–	–	–
Male sex	2.02 (0.64–6.33)	0.23	–	–	–	–
CCI (per 1 SD)	1.52 (0.82–2.84)	0.18	–	–	–	–
LDH (log-values per 1 SD)	3.85 (1.37–10.80)	0.011	2.72 (0.65–11.40)	0.17	3.25 (0.80–13.24)	0.10
IL-6 (log-values per 1 SD)	3.18 (1.32–7.69)	0.010	4.08 (1.16–14.33)	0.028	4.13 (1.21–14.10)	0.024
Presence of severe ARDS	11.5 (2.26–59.0)	0.003	3.35 (0.26–42.97)	0.35	–	–
Intubation	8.10 (2.14–30.65)	0.002	–	–	2.57 (0.44–14.89)	0.29
Duration NIV \geq 5 days prior to ICU	10.1 (1.1–91.8)	0.040	42.20 (1.22 to >999)	0.038	32.76 (1.02 to >999)	0.049

Model 1: including severe ARDS, LDH (log-transformed), IL-6 (log-transformed), and duration NIV \geq 5 days prior to ICU admission; Model 2: including intubation, LDH (log-transformed), IL-6 (log-transformed), and duration NIV \geq 5 days prior to ICU admission. COVID-19, coronavirus disease 2019; OR, odds ratio; CI, confidence interval; SD, standard deviation; CCI, Charlson Comorbidity Index; LDH, lactate dehydrogenase; IL-6, interleukin-6; ARDS, acute respiratory distress syndrome; NIV, non-invasive ventilation; ICU, intensive care unit.

terminology “NIV-failure” in this context emphasizes the type of respiratory support on outcome. However, it should be recognized that “late failure” in COVID-19 also occurs on IMV, as reflected by long intubation times, escalation to ECMO therapy, and death due to respiratory failure. At present it is unclear if patients who have an unfavourable clinical course on NIV, should be regarded as “patients with NIV-failure” or as “patients with delayed or no recovery from lung failure”. The focus should currently be on both, i.e., (I) on the type of ventilation and (II) on disease severity and recovery from it. To date, prospective trials in COVID-19 patients in moderate or severe ARDS comparing timing of NIV and IMV are lacking. Several trials on the prognostic role of NIV in COVID-19 have been initiated (37), but results are pending.

Considering our data, evidence from other studies, and published opinion, it appears safe to state that patients with worsening respiratory failure and $pO_2/FiO_2 < 200$ mmHg, that do not improve or even deteriorate clinically after 4–5 days of NIV, are candidates for very careful ICU surveillance. Whether outcome in these patients is better with intubation and IMV, remains to be shown.

An additional finding unrelated to ventilation support is that COVID-19 patients differed from the 18% SARS-CoV-2 comorbid patients in several clinically relevant aspects, despite similar mortality. Mortality in our patients without COVID-19 may have been worsened due to SARS-CoV-2 infection, as in other cohorts (38). From such few patients, it is difficult to draw conclusions for clinical

practice. Yet, they appear to deserve special attention regarding diagnostic work up, surveillance and therapy. Patients with SARS-CoV-2 as comorbidity have previously not explicitly been excluded or mentioned (24,39,40). A separate analysis may help to better understand determinants of outcome in patients with both SARS-CoV-2 or COVID-19-induced ARDS.

This work has some limitations. It is a comparatively small single-center study. Therefore, absolute numbers as well as intubation and mortality rates in the different categories must be interpreted with caution. It is an advantage, though, that we had clinical, respiratory and laboratory values for each individual patient in a cohort with high NIV rates and late intubation, and all patients had completed ICU stay.

We could not include data of patients that were treated outside our ICU. It is thus expected that overall hospital mortality in each category of ARDS severity is lower than reported here. Our findings should nonetheless be relevant to ICU physicians treating patients in similar settings.

In patients on NIV, FiO_2 could not be measured but had to be estimated from tabulations. However, the tables used in this study have also been used in large trials (16,17), and have been validated (15). Using alternative estimates would have overestimated FiO_2 and hypoxemia severity (15).

NIV, that comprises CPAP, bilevel CPAP, HFNO, NIV via helmet or face mask, etc., was coded irrespective of interface, mode and ventilator type employed. We used NIV via face-mask in most and HFNO in few patients

(10,11). Our data can therefore not be transferred to other modes of non-invasive respiratory support.

We have not reported details on ventilator settings and supporting medication. These data are stored automatically in our hospital Krankenhaus Informationssystem (KIS) (ORBIS KIS, Dedalus Health Care, Bonn, Germany). Yet, in previous studies, this information contained little prognostic information (25), and principles of lung-protective ventilation taking into account specific COVID-19 pathophysiology, were applied (1,6,7,12,13).

During the pandemic, treatment recommendations have been modified (1,12,13). For the period of this study, the modifications mainly pertained to ventilation support strategies, corticoid therapy and anticoagulation. Throughout the study period, we were restrictive with IMV, and used NIV below pO_2/FiO_2 thresholds suggested in guidelines. Early on, we used i.v. corticosteroids in COVID-19 patients with ARDS, as suggested by Surviving Sepsis Campaign guidelines (41). Even though anticoagulation was not mentioned in the initial guidelines (12), all patients were individually treated with prophylactic, half-therapeutic or full-dose unfractionated or low molecular weight heparin depending on risk factors such as overweight or elevated D-dimers (1,7,13). The degree to which such variation in therapy over time may have affected outcome is difficult to estimate. Interestingly, a decrease of early IMV from 75% in the first period to 37% in the second period of the pandemic did not reduce overall mortality (8).

In summary, in this 1-year all comers study, we found a clinical difference between patients with SARS-CoV-2 as comorbidity and COVID-19 patients. This should be considered in future analyses. In COVID-19 patients, prognosis appears to be largely determined by ARDS severity and the degree of accompanying inflammation. Especially patients ≥ 5 days on NIV appear to have a very poor prognosis. Our data indicate that patients with severe COVID-19 hypoxemia that do not improve on NIV during the first days, should be considered candidates for invasive ventilation to reduce work of breathing and maybe P-SILI. Yet, it is currently unclear if long duration on NIV aggravates the disease process or if it indicates lack of recovery, or both. The prognostic role of NIV and best timing of intubation for outcome should be clarified in prospective trials.

Acknowledgments

We gratefully acknowledge the hard work of all nursing

and physician staff on our ICU in their care for COVID-19 and non-COVID-19 patients alike. We are indebted to Dr. Jeibmann (Bethanien Hospital Moers) who contributed the data for *Figure 1*.

Funding: None.

Footnote

Reporting Checklist: The authors have completed the STROBE reporting checklist. Available at <https://jtd.amegroups.com/article/view/10.21037/jtd-21-1713/rc>

Data Sharing Statement: Available at <https://jtd.amegroups.com/article/view/10.21037/jtd-21-1713/dss>

Peer Review File: Available at <https://jtd.amegroups.com/article/view/10.21037/jtd-21-1713/prf>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://jtd.amegroups.com/article/view/10.21037/jtd-21-1713/coif>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The study complies with the Declaration of Helsinki (as revised in 2013), and was approved by the ethics committee of University Duisburg-Essen (#21-9911-BO). Due to the retrospective nature of the study and blinding of study data, the need for informed consent was waived.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References


1. Kluge S, Janssens U, Welte T, et al. Empfehlungen zur stationären Therapie von Patienten mit COVID-19. AWMF-Register-Nr. 113/001. (S3-Leitlinie, 02/21)

- Available online: https://www.grc-org.de/files/ArticleFiles/document/113-0011_S3_Empfehlungen-zur-stationaeren-Therapie-von-Patienten-mit-COVID-19__2021-02.pdf
2. COVID-19 Treatment Guidelines Panel. Coronavirus Disease 2019 (COVID-19) Treatment Guidelines. National Institutes of Health. Available online: <https://www.covid19treatmentguidelines.nih.gov/> (last accessed 02 April 2021)
 3. Wu C, Chen X, Cai Y, et al. Risk Factors Associated With Acute Respiratory Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China. *JAMA Intern Med* 2020;180:934-43.
 4. Richardson S, Hirsch JS, Narasimhan M, et al. Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA* 2020;323:2052-9.
 5. Ruan Q, Yang K, Wang W, et al. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive Care Med* 2020;46:846-8.
 6. Gattinoni L, Chiumello D, Caironi P, et al. COVID-19 pneumonia: different respiratory treatments for different phenotypes? *Intensive Care Med* 2020;46:1099-102.
 7. Möhlenkamp S, Thiele H. Ventilation of COVID-19 patients in intensive care units. *Herz* 2020;45:329-31.
 8. Karagiannidis C, Hentschker C, Westhoff M, et al. Observational study of changes in utilization and outcomes in mechanical ventilation in COVID-19. *PLoS One* 2022;17:e0262315.
 9. Crimi C, Noto A, Cortegiani A, et al. Noninvasive respiratory support in acute hypoxemic respiratory failure associated with COVID-19 and other viral infections. *Minerva Anestesiol* 2020;86:1190-204.
 10. Voshaar T, Dellweg D, Hetzel M. Empfehlung zur Behandlung respiratorischer Komplikationen bei akuter Virusinfektion außerhalb der Intensivstation. Published by Verband Pneumologischer Kliniken. Available online: https://www.vpnemo.de/fileadmin/pdf/VPK_Empfehlung_neu_21.03.2020.pdf
 11. Voshaar T, Stais P, Köhler D, et al. Conservative management of COVID-19 associated hypoxaemia. *ERJ Open Res* 2021;7:00134-2021.
 12. Kluge S, Janssens U, Welte T, et al. Recommendations for critically ill patients with COVID-19. *Med Klin Intensivmed Notfmed* 2020;115:175-7.
 13. Kluge S, Janssens U, Spinner CD, et al. Clinical Practice Guideline: Recommendations on Inpatient Treatment of Patients with COVID-19. *Dtsch Arztebl Int* 2021. [Epub ahead of print]. doi: 10.3238/arztebl.m2021.0110.
 14. ARDS Definition Task Force; Ranieri VM, Rubenfeld GD, et al. Acute respiratory distress syndrome: the Berlin Definition. *JAMA* 2012;307:2526-33.
 15. Coudroy R, Frat JP, Girault C, et al. Reliability of methods to estimate the fraction of inspired oxygen in patients with acute respiratory failure breathing through non-rebreather reservoir bag oxygen mask. *Thorax* 2020;75:805-7.
 16. Frat JP, Thille AW, Mercat A, et al. High-flow oxygen through nasal cannula in acute hypoxemic respiratory failure. *N Engl J Med* 2015;372:2185-96.
 17. SRLF Trial Group. Hypoxemia in the ICU: prevalence, treatment, and outcome. *Ann Intensive Care* 2018;8:82.
 18. Deutsche Sepsis Gesellschaft (DSG). Available online: <https://www.sepsis-gesellschaft.de/sepsisdefinition-und-kodierung/> (last query on August 20, 2021)
 19. Deutsche Sepsis-Gesellschaft (DSG). "Sepsis 3" und Kodierung der "Sepsis" gemäß ICD-10-GM 2020. Available online: <https://www.sepsis-gesellschaft.de/sepsisdefinition-und-kodierung/>
 20. Singer M, Deutschman CS, Seymour CW, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA* 2016;315:801-10.
 21. Vincent JL, Moreno R, Takala J, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine. *Intensive Care Med* 1996;22:707-10.
 22. Lambden S, Laterre PF, Levy MM, et al. The SOFA score-development, utility and challenges of accurate assessment in clinical trials. *Crit Care* 2019;23:374.
 23. Charlson ME, Pompei P, Ales KL, et al. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987;40:373-83.
 24. Roedl K, Jarczak D, Thasler L, et al. Mechanical ventilation and mortality among 223 critically ill patients with coronavirus disease 2019: A multicentric study in Germany. *Aust Crit Care* 2021;34:167-75.
 25. Pfeifer M, Ewig S, Voshaar T, et al. Position Paper for the State-of-the-Art Application of Respiratory Support in Patients with COVID-19. *Respiration* 2020;99:521-42.
 26. Papoutsis E, Giannakoulis VG, Xourgia E, et al. Effect of timing of intubation on clinical outcomes of critically ill patients with COVID-19: a systematic review and meta-analysis of non-randomized cohort studies. *Crit Care* 2021;25:121.

27. Bellani G, Laffey JG, Pham T, et al. Noninvasive Ventilation of Patients with Acute Respiratory Distress Syndrome. Insights from the LUNG SAFE Study. *Am J Respir Crit Care Med* 2017;195:67-77.
28. Wendel Garcia PD, Aguirre-Bermeo H, Buehler PK, et al. Implications of early respiratory support strategies on disease progression in critical COVID-19: a matched subanalysis of the prospective RISC-19-ICU cohort. *Crit Care* 2021;25:175.
29. Daniel P, Mecklenburg M, Massiah C, et al. Non-invasive positive pressure ventilation versus endotracheal intubation in treatment of COVID-19 patients requiring ventilatory support. *Am J Emerg Med* 2021;43:103-8.
30. Brochard L, Slutsky A, Pesenti A. Mechanical Ventilation to Minimize Progression of Lung Injury in Acute Respiratory Failure. *Am J Respir Crit Care Med* 2017;195:438-42.
31. Windisch W, Weber-Carstens S, Kluge S, et al. Invasive and Non-Invasive Ventilation in Patients With COVID-19. *Dtsch Arztebl Int* 2020;117:528-33.
32. Grieco DL, Menga LS, Eleuteri D, et al. Patient self-inflicted lung injury: implications for acute hypoxemic respiratory failure and ARDS patients on non-invasive support. *Minerva Anestesiol* 2019;85:1014-23.
33. Tobin MJ, Laghi F, Jubran A. Caution about early intubation and mechanical ventilation in COVID-19. *Ann Intensive Care* 2020;10:78.
34. Tobin MJ, Jubran A, Laghi F. Noninvasive strategies in COVID-19: epistemology, randomised trials, guidelines, physiology. *Eur Respir J* 2021;57:2004247.
35. Tobin MJ. Basing Respiratory Management of COVID-19 on Physiological Principles. *Am J Respir Crit Care Med* 2020;201:1319-20.
36. L'Her E, Deye N, Lellouche F, et al. Physiologic effects of noninvasive ventilation during acute lung injury. *Am J Respir Crit Care Med* 2005;172:1112-8.
37. Dobler CC, Murad MH, Wilson ME. Noninvasive Positive Pressure Ventilation in Patients With COVID-19. *Mayo Clin Proc* 2020;95:2594-601.
38. Doglietto F, Vezzoli M, Gheza F, et al. Factors Associated With Surgical Mortality and Complications Among Patients With and Without Coronavirus Disease 2019 (COVID-19) in Italy. *JAMA Surg* 2020;155:691-702.
39. Auld SC, Caridi-Scheible M, Blum JM, et al. ICU and Ventilator Mortality Among Critically Ill Adults With Coronavirus Disease 2019. *Crit Care Med* 2020;48:e799-804.
40. Doidge JC, Gould DW, Ferrando-Vivas P, et al. Trends in Intensive Care for Patients with COVID-19 in England, Wales, and Northern Ireland. *Am J Respir Crit Care Med* 2021;203:565-74.
41. Alhazzani W, Möller MH, Arabi YM, et al. Surviving Sepsis Campaign: guidelines on the management of critically ill adults with Coronavirus Disease 2019 (COVID-19). *Intensive Care Med* 2020;46:854-87.

Cite this article as: Brücker W, Mahabadi AA, Hüschen A, Becker J, Daehnke S, Möhlenkamp S. Clinical characteristics and determinants of mortality in coronavirus disease 2019 (COVID-19) patients on an intensive care unit—a retrospective explorative 1-year all-comers study. *J Thorac Dis* 2022;14(5):1319-1331. doi: 10.21037/jtd-21-1713

Supplementary



*: including n = 6 who died on ECMO in external hospital and n = 3 who died on ward after discharge from ICU

Figure S1 Admission and discharge of patients to and from ICU. One patient was excluded because the patient was transferred for weaning after survived COVID-19 infection. COVID-19, coronavirus disease 2019; ECMO, extracorporeal membrane oxygenation; ICU, intensive care unit; IMC, intermediate care unit.

Table S1 Tabulation for converting oxygen insufflation to FiO₂ in patients on NIV using the “3% formula”: FiO₂ (%) = 21% + flow (L/min) × 3%. pO₂ was measured to calculate pO₂/FiO₂ (Coudroy *et al.* 2020; Frat *et al.* 2015)

Non-invasive ventilation (NIV)	
O ₂ flow (L/min)	Estimated FiO ₂ (%)
1	24
2	27
3	30
4	33
5	36
6	39
7	42
8	45
9	48
10	51
11	54
12	57
13	60
14	63
15	66

FiO₂, inspiratory oxygen fraction; pO₂, partial pressure of oxygen.

Table S2 Tabulation for converting oxygen insufflation to FiO₂ in patients on nasal cannula (Marx R *et al.* 2013; DSG: <https://www.sepsis-gesellschaft.de/sepsisdefinition-und-kodierung/>)

Nasal cannula	
O ₂ flow (L/min)	Estimated FiO ₂ (%)
1	24
2	28
3	32
4	36
5	40
6	44

DSG, Deutsche Sepsis Gesellschaft; FiO₂, inspiratory oxygen fraction.

Table S3 Tabulation to estimate pO₂ from O₂ saturation (SO₂) (Marx R *et al.* 2013; DSG: <https://www.sepsis-gesellschaft.de/sepsisdefinition-und-kodierung/>)

SO ₂ (%)	Estimated pO ₂ (mmHg)
80	44
81	45
82	46
83	47
84	49
85	50
86	52
87	53
88	55
89	57
90	60
91	62
92	65
93	69
94	73
95	79
96	86
97	96
98	112
99	145

DSG, Deutsche Sepsis Gesellschaft.

Table S4 List of diagnoses established prior to ICU admission in COVID-19 (n=50)

Diagnoses	Percentage
Atrial fibrillation	26%
Obstructive sleep apnea syndrome (OSAS)	18%
Psychiatric and/or neurological diseases	18%
Malignancies/tumor	6%
Liver steatosis and/or cirrhosis	4%
Deep vein thrombosis (DVT) and/or pulmonary embolus	4%
Peptic ulcer	4%
Pituitary insufficiency	2%
Systemic lupus erythematosus	2%

COVID-19, coronavirus disease 2019; ICU, intensive care unit.