

## STARD 2015

Section and topic	No	Item	Reported on Page Number/Line Number	Reported on Section/Paragraph
<b>Title or abstract</b>				
	1	Identification as a study of diagnostic accuracy using at least one measure of accuracy (such as sensitivity, specificity, predictive values, or AUC)	Page 1/Line 2-3	Title section. Paragraph 1.
	2	Structured summary of study design, methods, results, and conclusions (for specific guidance, see STARD for Abstracts)	Page 1-1/Line 20-57	Abstract section. Paragraph 1.
<b>Introduction</b>				
	3	Scientific and clinical background, including the intended use and clinical role of the index test	Page 3-4/Line 75-115	Introduction section. Paragraph 1-3.
	4	Study objectives and hypotheses	Page 4-5/Line 116-150	Introduction section. Paragraph 4.
<b>Methods</b>				
Study design	5	Whether data collection was planned before the index test and reference standard were performed (prospective study) or after (retrospective study)	Page 5-6/Line 156-176	Materials and Methods section Study objects section. Paragraph 1-4.
Participants	6	Eligibility criteria for participants	Page 6/ Line 1685-177	Study objects section. Paragraph 3-4.
	7	Settings and locations where the data were collected	Page 5/Line 156-162	Study objects section. Paragraph 1.
	8	Where and when potentially eligible participants were identified (setting, location and dates)	Page 5/Line 156-162	Study objects section. Paragraph 1.

	9	Whether participants formed a consecutive, random or convenience series	Page 2-3/Line 95-100	Yes
Test methods	10a	Index test, in sufficient detail to allow replication	Page 3-6/Line 143-209	Materials and Methods section. Paragraph 8-15.
	10b	Reference standard, in sufficient detail to allow replication	Page 3/Line 116-141	Materials and Methods section. Paragraph 8-12.
	11	Rationale for choosing the reference standard (if alternatives exist)	Page 3/Line 116-141	Materials and Methods section. Paragraph 8-12.
	12a	Definition of and rationale for test positivity cut-offs or result categories of the index test, distinguishing pre-specified from exploratory	Page 2-6/Line 95-209	Materials and Methods section. Paragraph 8-15.
	12b	Definition of and rationale for test positivity cut-offs or result categories of the reference standard, distinguishing pre-specified from exploratory	Page 2-6/Line 95-209	Materials and Methods section. Paragraph 8-18.
	13a	Whether clinical information and reference standard results were available to the performers or readers of the index test	Page 6/Line 211-215	Yes. Paragraph 18-22.
	13b	Whether clinical information and index test results were available to the assessors of the reference standard	Page 6/Line 211-215	Yes. Paragraph 18-22.
Analysis	14	Methods for estimating or comparing measures of diagnostic accuracy	Page 6/Line 218-224	Statistical methods section . Paragraph 18-22.
	15	How indeterminate index test or reference standard results were handled	Page 6/Line 218-224	Statistical methods section . Paragraph 18-22.
	16	How missing data on the index test and reference standard were handled	Page 6/Line 218-224	Statistical methods section. Paragraph 18-22.

	17	Any analyses of variability in diagnostic accuracy, distinguishing pre-specified from exploratory	Page 6-10/Line 180-300	Materials and Methods section Paragraph 5-28
	18	Intended sample size and how it was determined	Page 10/Line 302-310	Materials and Methods section /Paragraph 29
<b>Results</b>				
Participants	19	Flow of participants, using a diagram	Page 10-11/Line 314-322	Results section /P 1
	20	Baseline demographic and clinical characteristics of participants	Page 11/Line 324-370	Results section Paragraph 2-5
	21a	Distribution of severity of disease in those with the target condition	Page 6/Line 324-370	Results section Paragraph 2-5
	21b	Distribution of alternative diagnoses in those without the target condition	Page 6/Line 324-370	Results section Paragraph 2-5
	22	Time interval and any clinical interventions between index test and reference standard	Page 6/Line 324-370	Results section Paragraph 2-5
Test results	23	Cross tabulation of the index test results (or their distribution) by the results of the reference standard	Page 6/Line 324-370	Results section Paragraph 2-5
	23	Estimates of diagnostic accuracy and their precision (such as 95% confidence intervals)	Page 6/Line 324-370	Results section Paragraph 2-5
	25	Any adverse events from performing the index test or the reference standard		no
<b>Discussion</b>				
	26	Study limitations, including sources of potential bias, statistical uncertainty, and generalisability	Page 12-14/Line 372-426	Discussion section Paragraph 1-5

	27	Implications for practice, including the intended use and clinical role of the index test	Page 12-13/Line 309-351	Discussion section
<b>Other information</b>				
	28	Registration number and name of registry	None.	None.
	29	Where the full study protocol can be accessed	None.	None.
	30	Sources of funding and other support; role of funders	None.	None.

## AIM

STARD stands for “Standards for Reporting Diagnostic accuracy studies”. This list of items was developed to contribute to the completeness and transparency of reporting of diagnostic accuracy studies. Authors can use the list to write informative study reports. Editors and peer-reviewers can use it to evaluate whether the information has been included in manuscripts submitted for publication.

## Explanation

A **diagnostic accuracy study** evaluates the ability of one or more medical tests to correctly classify study participants as having a **target condition**. This can be a disease, a disease stage, response or benefit from therapy, or an event or condition in the future. A medical test can be an imaging procedure, a laboratory test, elements from history and physical examination, a combination of these, or any other method for collecting information about the current health status of a patient.

The test whose accuracy is evaluated is called **index test**. A study can evaluate the accuracy of one or more index tests. Evaluating the ability of a medical test to correctly classify patients is typically done by comparing the distribution of the index test results with those of the **reference standard**. The reference standard is the best available method for establishing the presence or absence of the target condition. An accuracy study can rely on one or more reference standards.

If test results are categorized as either positive or negative, the cross tabulation of the index test results against those of the reference standard can be used to estimate the **sensitivity** of the index test (the proportion of participants with the target condition who have a positive index test), and its **specificity** (the proportion without the target condition who have a negative index test). From this cross tabulation (sometimes referred to as the contingency or “2x2” table), several other accuracy statistics can be estimated, such as the positive and negative **predictive values** of the test. Confidence intervals around estimates of accuracy can then be calculated to quantify the statistical **precision** of the measurements.

If the index test results can take more than two values, categorization of test results as positive or negative requires a **test positivity cut-off**. When multiple such cut-offs can be defined, authors can report a receiver operating characteristic (ROC) curve which graphically represents the combination of sensitivity and specificity for each possible test positivity cut-off. The **area under the ROC curve** informs in a single numerical value about the overall diagnostic accuracy of the index test.

The **intended use** of a medical test can be diagnosis, screening, staging, monitoring, surveillance, prediction or prognosis. The **clinical role** of a test explains its position relative to existing tests in the clinical pathway. A replacement test, for example, replaces an existing test. A triage test is used before an existing test; an add-on test is used after an existing test.

Besides diagnostic accuracy, several other outcomes and statistics may be relevant in the evaluation of medical tests. Medical tests can also be used to classify patients for purposes other than diagnosis, such as staging or prognosis. The STARD list was not explicitly developed for these other outcomes, statistics, and study types, although most STARD items would still apply.

## DEVELOPMENT

This STARD list was released in 2015. The 30 items were identified by an international expert group of methodologists, researchers, and editors. The guiding principle in the development of STARD was to select items that, when reported, would help readers to judge the potential for bias in the study, to appraise the applicability of the study findings and the validity of conclusions and recommendations. The list represents an update of the first version, which was published in 2003. More information can be found on <http://www.equator-network.org/reporting-guidelines/stard>.

Article information: <https://dx.doi.org/10.21037/jtd-23-190>

\*As the checklist was provided upon initial submission, the page number/line number reported may be changed due to copy editing and may not be referable in the published version. In this case, the section/paragraph may be used as an alternative reference.