

The impact of decreasing cutoff values for maximal oxygen consumption (VO₂max) in the decision-making process for candidates to lung cancer surgery

Gaetano Rocco^{1,2}, Tindaro Gatani^{1,2}, Massimo Di Maio^{1,2}, Ilernando Meoli^{1,2}, Antonello La Rocca^{1,2}, Nicola Martucci^{1,2}, Carmine La Manna^{1,2}, Francesco Stefanelli^{1,2}

¹Department of Thoracic Surgery and Oncology, Division of Thoracic Surgery and Service of Physiopathology, National Cancer Institute, Naples, Italy; ²Division of Respiratory Physiopathology, Monaldi Hospital, Naples, Italy

ABSTRACT

Background: Maximal oxygen consumption (VO₂max) is considered a decisive test for risk prediction in patients with borderline cardiopulmonary reserve. Guidelines have adopted decreasing VO₂max cut-off values to define operability within acceptable mortality and morbidity limits. We wanted to investigate how the adoption of decreasing VO₂max cut-off-values assessment contributed to better select lung surgery candidates.

Methods: One hundred and nineteen consecutive surgical candidates have been prospectively analyzed as a sample population. Preoperative work-up included spirometry and transfer factor (DLco); irrespective of the spirometric values, these patients were subjected to VO₂max assessment. Surgical eligibility was decided by the same surgeon throughout the series. In the postoperative period, overall mortality and the occurrence of any, major or minor complications was recorded and graded according to the Common Terminology Criteria for Adverse Events v.4.3.

Results: Three arbitrary cut-offs were introduced at 15, 14 and 12 mL·kg⁻¹·min⁻¹. Notably, 15 and 12 mL·kg⁻¹·min⁻¹ correlated with percentage VO₂max values of 50% and 35% of predicted (P<0.0001 and 0.0079), respectively. Accordingly, the patients were subdivided into groups in which the prevalence of postoperative morbidity was recorded. The groups were homogeneous as to age, BMI, preoperative absolute and percentage FEV1 and DLco. In the Cox proportionate-hazards multivariate analysis, VO₂max less than 35% (P=0.0017) and CTCAE >2 (P=0.0457) emerged as significant predictors of survival after surgery. Conversely on logistic regression analysis, age over 70 years (P=0.03) and pneumonectomy (P=0.001), but not VO₂max cut-off values, were significant predictors of major (CTCAE >2) morbidity.

Conclusions: Since VO₂max is increasingly used to contribute to risk prediction for the individual patient, surgeons need to be advised that the concept of a definitive, generalized cut-off value for VO₂max is probably a contradiction in terms. Patient-specific VO₂max values are more likely to contribute to risk assessment since they may reflect the primarily affected component among the determinants of maximal oxygen consumption. Whether patient-specific VO₂max should be routinely used by surgeons to define operability for borderline patients needs further evaluation.

KEY WORDS

Maximal oxygen consumption; lung surgery; lung cancer

J Thorac Dis 2013;5(1):12-18. DOI: 10.3978/j.issn.2072-1439.2012.12.04

Introduction

In spite of the results from the analysis of multiinstitutional databases and the current guidelines focused on risk assessment for lung resection (1,2), the surgeon's perception of operability may at times not coincide with a borderline or adverse outcome of preoperative work-up (3). Selected patients have been offered surgery based on the surgeon's clinical instinct which still represents a useful adjunct to spirometric evaluation. Cardiopulmonary exercise testing is increasingly used to refine

Corresponding to: Gaetano Rocco, MD, FRCSEd, FETCS. Department of Thoracic Surgery and Oncology, Division of Thoracic Surgery, National Cancer Institute, Pascale Foundation, Via Semmola 81, 80131, Naples, Italy. Email: Gaetano.Rocco@btopenworld.com.

Submitted Nov 17, 2012. Accepted for publication Dec 18, 2012.
Available at www.jthoracdis.com

ISSN: 2072-1439

© Pioneer Bioscience Publishing Company. All rights reserved.

the selection of surgical candidates (3-5). Decreasing cut-off values of maximal oxygen consumption (VO_2max) have been proposed to avoid denying a surgical option to lung cancer patients (4). We investigated the possible correlation between the progressive reduction of VO_2max cut-offs and postoperative survival and morbidity after lung cancer surgery.

Material and methods

Between January 2008 and December 2010, 119 consecutive candidates to pulmonary resection and operated on at the National Cancer Institute in Naples were subjected to VO_2max assessment as part of the routine preoperative work-up for lung cancer surgery. Cycle ergometry with incremental 10 watts RAMP (Cosmed Quark CPET, Italy) adaptable to the patient's clinical condition, was used to determine VO_2max (maximal O_2 value at plateau despite increasing workload), peak VO_2 (maximal oxygen consumption at maximal workload), VCO_2 (quantity of expired CO_2 at a given time), RQ (VCO_2/VO_2 or respiratory quotient), VO_2/HR (oxygen consumption per systole), HRR (cardiac reserve) and the anaerobic threshold.

This patient population included 80 males and 39 females with a median age of 64 years (range, 22 to 80 years). Eighty-two percent of these surgical candidates were either former or current smokers. The median body mass index (BMI) was 26 kg/m^2 (range, 17 to 50 kg/m^2). In particular, 66 of the patients in this series had a BMI of 25 or greater and were accordingly considered overweight or obese (5). Comorbidities included previous stroke in 4 patients (3.4%), concurrent significant peripheral vascular and coronary artery disease in 5 (4.2%) and 11 (9.2%) patients, respectively. In addition, 2 patients (1.7%) had atrial fibrillation whereas 61 (51.3%) had a history of hypertension under pharmacological treatment. Other comorbidities included diabetes in 17 patients (14.3%) and previous cancer history in 45 (37.8%). Thirty-eight patients (32%) had had chemotherapy or chemoradiotherapy as a part of their preoperative disease management. Of 86 patients undergoing lung resection for primary non small cell lung cancer, 11 (13%) had been subjected to neoadjuvant treatment. In the remaining patients, final pathology showed lung metastases (25 patients) and benign lesions (8 patients).

Of 119 patients, 11 (9.3%) had a preoperative FEV_1 less than 70% of predicted. The median FEV_1 absolute and percentage values were 2.29 L (range, 0.86-4.01 L) and 92.0% (range, 37.7-135.4%), respectively. In addition, the median DLco absolute and percentage values were 18.4 (range, 9.9-33.6) and 78.3% (range, 45-132%). Also, median VO_2max expressed in absolute and percentage values were $15.3 \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ (range, 8.3 - $28.8 \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$) and 62% (range, 31-129%).

Reasons for surgery included non small cell lung cancer in 80 patients (67.2%), metastases to the lung in 25 (21%) and benign lesions in the remaining 14 (11.8%). A pneumonectomy was

performed in 20 patients (10.6%), a lobectomy in 54 patients (45.4%), a bilobectomy in 5 (4.2%), and, single or multiple segmentectomy/wedge resections in 40 (33.6%). A right-sided pulmonary resection was performed in 72 patients (60.5%). The standard surgical approach for primary lung cancer was a limited (8 to 10 cm) incision in the auscultatory triangle with minimal rib spreading and no muscle division with or without videoassistance (6). The decisions on operability was made in all instances by the same board-certified surgeon (G.R.) who also performed or directly supervised the surgical procedures. When final pathology was obtained in non small cell lung cancer patients, the disease was staged as pIA in 21 patients (26.2%), pIB in 44 (55%), pIIA in 5 (6.3%), pIIB in 4 (5%), pIIIA in 5 (6.3%), and, pIIIB in 1 (1.2%).

Major and minor postoperative morbidity were attributed according to the NCI Common Terminology Criteria for Adverse Events v.4.3 (7). In addition, to understand the relationship between morbidity and VO_2max and whether a cut-off value could be predictive of postoperative complications, we arbitrarily subdivided our patient population into three groups according to the VO_2max values currently reported in the literature (12 and $15 \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$) and a conservative (i.e., closer to the safest figure) intermediate value ($14 \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$).

This routine use of cycle ergometry was funded by the Italian National Health System (SSN) within the context of an institutional project aimed at improving risk management in oncologic surgery. Informed consent was obtained in all patients to undergo cardiopulmonary exercise testing prior to considering a surgical option.

Statistical methods

Pearson's correlation and t-test were used to define correlations between continuous variables. Categorical variables were compared by chi-square tests whereas Spearman's tests were used to verify correlation. Multivariate analyses were conducted by logistic regression to define prognosticators for postoperative morbidity. In addition, Cox proportional hazard regression was conducted to evaluate survival predictors. Significance was set at $P < 0.05$. For statistical analysis, SPSS version 19 and MedCalc version 12 softwares were used. Isometric graphs were constructed by using the JMP statistical software, version 9.

Results

In this series of 119 patients, correlations between FEV_1 , $\text{FEV}_1\%$, DLco and DLco% and VO_2max along with $\text{VO}_2\text{max}\%$ were studied. Expectedly, the correlation FEV_1 -DLco appeared statistically significant ($P < 0.0001$). In addition, a definite correlation FEV_1 - VO_2max ($P = 0.001$) and DLco%- $\text{VO}_2\text{max}\%$

($P=0.0235$) was also found. Moreover, in our series, 43 patients (36.1%) did have a disjunction between absolute and percentage VO_{2max} values. Absolute VO_{2max} values of 15 and 12 $mL \cdot kg^{-1} \cdot min^{-1}$ correlated with percentage VO_{2max} values of 50% ($P<0.0001$; $\rho=0.366$, 95% CI for ρ , 0.199 to 0.512) and 35% of predicted ($P=0.0079$; $\rho=0.242$, 95% CI for $\rho=0.0652$ to 0.405), respectively.

Overall, 4 patients died within 30 days of surgery (3.3%). Median FEV_1 , $FEV_1\%$, DLco and DLco% measured in these patients were 2.15 L (range, 1.29 to 2.8 L), 92.6% (range, 80% to 132%), 17.84 (range, 10.94 to 25.7), and, 83.2% (range, 64.3% to 107%), respectively. Absolute and percentage of predicted VO_{2max} values in these patients were 10.7, 12.2, 12.1, 19.3 $mL \cdot kg^{-1} \cdot min^{-1}$ and 65%, 56%, 57%, 103%, respectively. Causes of death included pulmonary embolism in 3 patients (2 after right pneumonectomy and 1 after right upper lobectomy) and sepsis following bronchopleural fistula in 1 following right lower lobectomy. In addition, postoperative complications occurred in 55 patients.

Group 1 - cut-off: 15 $mL \cdot kg^{-1} \cdot min^{-1}$

Fifty-three patients (44.5%) had a VO_{2max} less than 15 $mL \cdot kg^{-1} \cdot min^{-1}$. No statistically significant differences were noted as to age, BMI, percentage preoperative FEV_1 and DLco between patients with VO_{2max} greater or lesser than 15 $mL \cdot kg^{-1} \cdot min^{-1}$. Patients with less than 15 $mL \cdot kg^{-1} \cdot min^{-1}$ underwent pneumonectomy in 14 (26.4%) cases, lobectomy or bilobectomy in 30 (56.7%), and, segmentectomy/wedge in 9 (16.9%). Of these patients, 21 (40%) experienced some postoperative complications. On the other hand, the remaining 32 patients with VO_{2max} less than 15 $mL \cdot kg^{-1} \cdot min^{-1}$ (60%) did not experience any complications in the postoperative course. The difference with the 66 patients with VO_{2max} greater than 15 $mL \cdot kg^{-1} \cdot min^{-1}$ was not significant ($P=0.26$). If we distinguish between patients having minor and major complications in the postoperative period of patients with VO_{2max} less than 15 $mL \cdot kg^{-1} \cdot min^{-1}$, only 5 patients (9%) had major morbidity compared to 48 (91%) who did not. Again, no statistically significant difference was noted with the patients with a VO_{2max} greater than 15 $mL \cdot kg^{-1} \cdot min^{-1}$ ($P=0.48$).

If the FEV_1 less than 40% of predicted threshold is considered, no difference was noted in the severity of complications, if any, based on the 15 $mL \cdot kg^{-1} \cdot min^{-1}$ cutoff ($P=0.21$). In particular, this cutoff did not help stratify the patient population who would have developed major complications postoperatively ($P=0.5$).

Group 2 - cut-off: 14 $mL \cdot kg^{-1} \cdot min^{-1}$

Thirty-eight (26.8%) had a VO_{2max} less than 14 $mL \cdot kg^{-1} \cdot min^{-1}$. A pneumonectomy was performed in 8 (21%), lobectomy or bilobectomy in 21 (55.3%), and segmentectomy/wedge in 9

(23.7%) patients. Of these patients, 15 (39%) experienced some postoperative complications. On the other hand, the remaining 23 patients with VO_{2max} less than 14 $mL \cdot kg^{-1} \cdot min^{-1}$ (61%) did not experience any complications in the postoperative course. The difference with the 81 patients with VO_{2max} greater than 14 $mL \cdot kg^{-1} \cdot min^{-1}$ was not significant ($P=0.38$). When we compared patients with VO_{2max} less than 14 $mL \cdot kg^{-1} \cdot min^{-1}$ developing minor and major complications postoperatively, only 3 patients (8%) had major complications while 35 (92%) did not. Again, no statistically significant difference was noted compared to the patients with a VO_{2max} greater than 14 $mL \cdot kg^{-1} \cdot min^{-1}$ ($P=0.37$).

Group 3 - cut-off: 12 $mL \cdot kg^{-1} \cdot min^{-1}$

Fifteen patients (12.6%) had a VO_{2max} less than 12 $mL \cdot kg^{-1} \cdot min^{-1}$; they underwent a pneumonectomy in 2 (13.3%) cases, lobectomy or bilobectomy in 7 (46.7%), and, segmentectomy/wedge in 6 (40%). Of these patients, 7 (47%) experienced some postoperative complications. On the other hand, the remaining 8 patients with VO_{2max} less than 12 $mL \cdot kg^{-1} \cdot min^{-1}$ (53%) did not experience any complications in the postoperative course. The difference with the 104 patients with VO_{2max} greater than 12 $mL \cdot kg^{-1} \cdot min^{-1}$ was not significant ($P=0.91$). If we distinguish between patients having minor and major complications in the postoperative period of patients with VO_{2max} less than 12 $mL \cdot kg^{-1} \cdot min^{-1}$, only 1 patient (7%) had major morbidity compared to 14 (93%) who did not. Again, no statistically significant difference was noted compared to patients with a VO_{2max} greater than 12 $mL \cdot kg^{-1} \cdot min^{-1}$ ($P=0.51$).

Pearson's correlation tests were carried out by plotting decreasing values of FEV_1 and DLco against different cutoff levels of VO_{2max} expressed both in $mL \cdot kg^{-1} \cdot min^{-1}$ and percentage of predicted. As expected, the categorical distinction (greater than or less than 15, 14 and 12 $mL \cdot kg^{-1} \cdot min^{-1}$) made it possible to separate our patient population in two risk groups according to each VO_{2max} cut-off value. If one looks at the diagrams contained in the figure in sequence (Figure 1), the upper parts of the diagrams show an increasing "osmosis" of high risk patients in the lower risk groups as we set the VO_{2max} cut-offs to decreasing values. Also, it clearly appears that a VO_{2max} less than 12 $mL \cdot kg^{-1} \cdot min^{-1}$ (or 35% of predicted) may contribute to the refinement of the stratification by clinical instinct albeit not all patients are finally allocated to newly determined risk class groups. In fact, the patients' data falling into the low FEV_1 -low DLco category (extreme sides of the scatterplots) become few and far between as the VO_{2max} cut-off values decreased to less than 12 $mL \cdot kg^{-1} \cdot min^{-1}$. When isometric graphs were constructed to study the correlation between FEV_1 and DLco at different absolute and percentage VO_{2max} values, the scattered distribution of VO_{2max} performance data was an obvious finding (Figure 2). Notably, patients with normal spirometry and/or DLco were found to have inconsistent




Figure 1. Scatterplot graphs of the correlation between FEV₁-DLco values and VO₂max operability cut-offs of 119 patients subjected to lung surgery; A. patient distribution according to a cut-off value of 15 mL·kg⁻¹·min⁻¹; B. patient distribution according to a cut-off value of 14 mL·kg⁻¹·min⁻¹; C. patient distribution according to a cut-off value of 12 mL·kg⁻¹·min⁻¹. Best values of FEV₁ and DLco converge under the arrow. Impaired function expressed on the abscissas by values decreasing towards the ordinates. Patient selection was refined by the adoption of decreasing VO₂max cut-off values (see text).

VO₂max values (Figure 2).

Correlation between VO₂max and FEV₁ and DLco in patients with VO₂max greater or less than 15 mL·kg⁻¹·min⁻¹

Of the 66 patients with measured VO₂max greater than 15 mL·kg⁻¹·min⁻¹ (i.e., normal preoperative risk), all (100%) had also an FEV₁ >40% of predicted. In addition, 61 of these patients (92%) had also a DLco greater than 40% of predicted. Conversely, of the 53 patients with measured VO₂max less than 15 mL·kg⁻¹·min⁻¹ (i.e., moderate to high preoperative risk), 52 (98%) had an FEV₁ greater than 40% of predicted whereas only 47 (88%) had a preoperative DLco greater than 40% of predicted.

Surgery for lung cancer

Of the 119 patients in this series, 86 were resected for primary

lung cancer. In this group, the patients subjected to wedge resection had a median VO₂max of 14.9 mL·kg⁻¹·min⁻¹ (range, 10.8-20 mL·kg⁻¹·min⁻¹); when expressed in percentage, the median VO₂max was 64.6% (range, 35-115%). The patients subjected to lobectomy had a median VO₂max of 16 mL·kg⁻¹·min⁻¹ (range, 8.32-28.89 mL·kg⁻¹·min⁻¹); the corresponding median percentage value was 68.3% (range, 42-129%). Pneumonectomy patients had a median VO₂max of 15.6 mL·kg⁻¹·min⁻¹ (range, 11.5-22.9 mL·kg⁻¹·min⁻¹); the corresponding median percentage value was 61.1% (range, 39-82%). When the same analysis as the lung resection candidates was carried out to correlate decreasing cutoff values of VO₂max with postoperative morbidity in lung cancer patients, no differences were noted in all three subgroups with regard to either major or minor complications.

Variables entered in the multivariate analyses were age (70 years cut-off), gender, BMI (25 kg/m² cut-off), hemoglobin levels,




Figure 2. Isometric graphs indicating how VO₂max absolute and percentage values contributed to the selection of surgical candidates. The contribution of VO₂max for those patients with “abnormal” FEV₁ and DLco (i.e., lower than 70% and 50%, respectively) is characterized by zones marked by arrows. Zones characterized by nuances of blue (VO₂max <50% or 15 mL·kg⁻¹·min⁻¹) are present also in correspondence of “normal” FEV₁ and DLco.

Table 1. Multivariate analysis on 119 patients of the predictors of major complications [CTCAE ≥grade 3 (7)].

Variable	Coefficient	Std. error	P	Odds ratio	95% CI
Age over 70 years	1.23845	0.57162	0.0303	3.4503	1.1253 to 10.5784
Pneumonectomy	1.95367	0.60548	0.0013	7.0545	2.1531 to 23.1132

smoking history, percent FEV₁ (70% cut-off), percent DL_{CO} (50% cut-off), CPET expressed by VO₂max above mentioned cut-off values, including 35% and 50% cut-off, type of operation (pneumonectomy *vs.* others), morbidity expressed by CTCAE grading system (7) (CTCAE grade 3 cut-off).

In this setting, when preoperative indicators were considered, multivariate analysis demonstrated that age and cardiorespiratory performance (i.e., absolute VO₂max values) correlated with postoperative morbidity at large (Table 1). In addition, age over 70 years and type of operation emerged as significant predictors of major [CTCAE ≥grade 3 (7)] postoperative complications (Table 2).

A 93.6% overall actuarial 5 year survival was computed with the Kaplan-Meier method. The Cox proportionate-hazards multivariate analysis demonstrated that both the percentage VO₂max value greater than 35% and CTCAE >3 emerged as significant predictors of postoperative 90-day survival (Table 3). Overall, the presence of a discrepancy between absolute and percentage VO₂max values was not a significant predictor of either 30 days mortality (P=0.509) or major morbidity (P=0.849).

Discussion

In clinical practice, the quest for the perfect indicator of

postoperative morbidity is far from over (8-11). In the meantime, the selection of surgical candidates is inspired by current guidelines which usually enable the surgeons to offer a surgical option to the majority of patients (1,2). In some circumstances, experienced surgeons tend to apply their “clinical instinct” to offer a surgical option despite potentially adverse spirometric figures (1,2). Concomitantly, in recent years, guidelines have been characterized by decreasing spirometric cutoff values and have introduced the concept of assessing the diffusing capacity and the cardiopulmonary performance in selected patients with borderline function (1,2,13). In this setting, cardiopulmonary exercise testing (CPET) has been extensively used and often referred to as the ultimate indicator of postoperative morbidity (14-16). Interestingly, high technology CPET has suffered the same fate as FEV₁ (and DLco) inasmuch as decreasing cut-offs have been proposed to define “the unacceptable surgical risk” (2). Recently, the ESTS-ERS guidelines have identified this cut-off in 10 mL·kg⁻¹·min⁻¹ or 35% of the predicted VO₂max value (2). Based on the review of the available literature, the limit for operability was lowered to 10 mL·kg⁻¹·min⁻¹ (or <35%) from the previous one (15 mL·kg⁻¹·min⁻¹) (2). Our series of surgical patients operated on prior to publication of these guidelines confirmed that the morbidity rates did not differ

Table 2. Predictors of major morbidity [CTCAE >3 (7)].

Variable	Coefficient	Std. error	P	Odds ratio	95% CI
Hb	0.38818	0.21663	0.0732	1.4743	0.9642 to 2.2542
Age over 70 years	1.17037	0.58031	0.0437	3.2232	1.0335 to 10.0522
Pneumonectomy	2.05372	0.62415	0.0010	7.7969	2.2942 to 26.4977
Constant	-8.1172				

Table 3. Cox proportionate-hazards analysis of functional predictors of survival on 119 patients.

Covariate	b	SE	P	Exp(b)	95% CI of Exp(b)
Age over 70 years	1.9424	1.1285	0.0852	6.9757	0.7724 to 62.9962
VO ₂ max >35%	4.6452	1.4828	0.0017	104.0849	5.7759 to 1875.6745
CTCAE (7) >3	1.8506	0.9263	0.0457	6.3634	1.0453 to 38.7375

significantly by altering the VO₂max cutoff from previously accepted values. However, we observed that the 4 patients who died within 30 days from surgery had median spirometric values within the range of normality. Nevertheless, VO₂max values were consistently around or below the accepted minimal cutoff for operability apart from one patient. However, there was a discrepancy between VO₂max absolute and percentage values in our 4 patients who died with 30 days of the operation despite the significant correlation in the rest of our patient population. This finding seems to suggest that the pathophysiology of the limitation of performance during CPET may indeed result from failure of different systems and hence entail different prognostic significance. In this setting, the mortality in our series was due to pulmonary embolism and bronchopleural fistula complicated by sepsis. Predicting such adverse events despite an optimal prophylactic and postoperative management is a difficult task. In this setting, in order not to deny a potentially curative treatment to lung cancer patients, thoracic surgeons are faced from time to time with the occasional patient who seems to escape risk management guidelines. In those circumstances, the “clinical instinct” comes into play and surgical eligibility is warranted despite borderline - or even adverse - function tests. In this setting, anthropometric measurements, location of bronchial obstruction, different protocol criteria or technologies for CPET, origin of exercise limitation (i.e., deconditioning, primary heart or lung condition, peripheral arteriopathy), nutritional profile - all may impact the accuracy of VO₂max estimation (17,18).

We believe that there are limitations to this study. This is a preliminary work which needs to be supported by a currently ongoing, prospective analysis comparing patients accepted for surgery to excluded, non surgical candidates. Also, the surgical risk should be stratified between central and peripheral lesions since type of operation and induction treatment play a major role in determining major morbidity. Nevertheless, this is a retrospective analysis of prospectively entered relatively limited

sized series of patients operated on directly or under the supervision of a single surgeon. By spirometric data, the majority of these patients did present “normal” functional (spirometric) values - a concept of normality which needs careful revision since it does not entail total risk avoidance. Indeed, only a minority of high risk patients do require extraordinary clinical and objective evaluation. The aim of this preliminary report was to understand whether do we always adhere to guidelines for risk assessment in these patients. Our study is a “snapshot” of a homogeneous clinical conduct in a time when several guidelines were published and none accurate enough to be fully adopted (12). Certainly, guidelines play a crucial role in contributing to the education of young surgeons in training, whereas experienced surgeons tend to still rely on clinical judgment especially when dealing with borderline patients (12).

In this setting, Ferguson and colleagues have clearly raised the issue of further investigating the etiology, accuracy and utility of surgeon risk estimates (12). The interpretation of patient-specific VO₂max values is more likely to contribute to risk assessment since these values may reflect the primarily affected component among the determinants of maximal oxygen consumption (18). Hence, should this concept be confirmed by larger studies, the adoption of generalized VO₂max cut-off values in risk assessment for lung surgery may in the future become a contradiction in terms. Risk scores will have to be revised taking into account the *primum movens* of exercise limitation of individualized, patient-specific, VO₂max cut-off values.

Acknowledgements

Disclosure: The authors declare no conflict of interest.

References

1. Bernard A, Rivera C, Pages PB, et al. Risk model of in-hospital mortality after pulmonary resection for cancer: a national database of the French

- Society of Thoracic and Cardiovascular Surgery (Epithor). *J Thorac Cardiovasc Surg* 2011;141:449-58.
2. Brunelli A, Charloux A, Bolliger CT, et al. ERS/ESTS clinical guidelines on fitness for radical therapy in lung cancer patients (surgery and chemoradiotherapy). *Eur Respir J* 2009;34:17-41.
 3. Bernard A, Ferrand L, Hagry O, et al. Identification of prognostic factors determining risk groups for lung resection. *Ann Thorac Surg* 2000;70:1161-7.
 4. Mishra PK, Pandey R, Shackcloth MJ, et al. Cardiac comorbidity is not a risk factor for mortality and morbidity following surgery for primary non-small cell lung cancer. *Eur J Cardiothorac Surg* 2009;35:439-43.
 5. Sinclair ME, Reber G, Schweizer A, et al. Anticoagulation by anocrod for carbon dioxide removal by extracorporeal membrane lung in the dog. *J Thorac Cardiovasc Surg* 1989;97:275-81.
 6. Smith PW, Wang H, Gazoni LM, et al. Obesity does not increase complications after anatomic resection for non-small cell lung cancer. *Ann Thorac Surg* 2007;84:1098-105; discussion 1105-6.
 7. 2010 NCI. Common Terminology Criteria for Adverse Events v 4.3 (CTCAE). Available online: http://evs.nci.nih.gov/ftp1/CTCAE/CTCAE_4.03_2010-06-14_QuickReference_5x7.pdf
 8. Brunelli A, Rocco G. Internal validation of risk models in lung resection surgery: bootstrap versus training-and-test sampling. *J Thorac Cardiovasc Surg* 2006;131:1243-7.
 9. Harpole DH Jr, DeCamp MM Jr, Daley J, et al. Prognostic models of thirty-day mortality and morbidity after major pulmonary resection. *J Thorac Cardiovasc Surg* 1999;117:969-79.
 10. Strand TE, Rostad H, Damhuis RA, et al. Risk factors for 30-day mortality after resection of lung cancer and prediction of their magnitude. *Thorax* 2007;62:991-7.
 11. Brunelli A, Berrisford RG, Rocco G, et al. The European Thoracic Database project: composite performance score to measure quality of care after major lung resection. *Eur J Cardiothorac Surg* 2009;35:769-74.
 12. Ferguson MK, Stromberg JD, Celauro AD. Estimating lung resection risk: a pilot study of trainee and practicing surgeons. *Ann Thorac Surg* 2010;89:1037-42; discussion 1042-3.
 13. Cerfolio RJ, Bryant AS. Optimal care of patients with non-small cell lung cancer reduces perioperative morbidity. *J Thorac Cardiovasc Surg* 2011;141:22-33.
 14. American Thoracic Society, American College of Chest Physicians. ATS/ACCP Statement on cardiopulmonary exercise testing. *Am J Respir Crit Care Med* 2003;167:211-77.
 15. British Thoracic Society, Society of Cardiothoracic Surgeons of Great Britain and Ireland Working Party. BTS guidelines: guidelines on the selection of patients with lung cancer for surgery. *Thorax* 2001;56:89-108.
 16. Weisman IM. Cardiopulmonary exercise testing in the preoperative assessment for lung resection surgery. *Semin Thorac Cardiovasc Surg* 2001;13:116-25.
 17. Tewari N, Martin-Ucar AE, Black E, et al. Nutritional status affects long term survival after lobectomy for lung cancer. *Lung Cancer* 2007;57:389-94.
 18. Brunelli A, Pompili C, Belardinelli R. Beyond peak VO₂: ventilatory inefficiency (VE/VCO₂ slope) measured during cardiopulmonary exercise test to refine risk stratification in lung resection candidates. *Eur J Cardiothorac Surg* 2010;38:19-20.



Cite this article as: Rocco G, Gatani T, Di Maio M, Meoli I, La Rocca A, Martucci N, La Manna C, Stefanelli F. The impact of decreasing cutoff values for maximal oxygen consumption (VO₂max) in the decision-making process for candidates to lung cancer surgery. *J Thorac Dis* 2013;5(1):12-18. DOI: 10.3978/j.issn.2072-1439.2012.12.04