



Prediction of postoperative final degree and recurrence of pectus excavatum using machine learning algorithms

Kwanyong Hyun¹, Jae Jun Kim², Won Kyu Choi³, Yoon Ho Kim³, Sang Chul Han³

¹Department of Thoracic and Cardiovascular Surgery, St. Vincent's Hospital, College of Medicine, The Catholic University of Korea, Gyeonggi-do, Republic of Korea; ²Department of Thoracic and Cardiovascular Surgery, Uijeongbu St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Gyeonggi-do, Republic of Korea; ³Department of Anesthesiology and Pain Medicine, Uijeongbu St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Gyeonggi-do, Republic of Korea

Contributions: (I) Conception and design: K Hyun, JJ Kim; (II) Administrative support: JJ Kim; (III) Provision of study materials or patients: WK Choi, YH Kim; (IV) Collection and assembly of data: K Hyun, SC Han; (V) Data analysis and interpretation: JJ Kim, K Hyun; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Jae Jun Kim, MD, PhD. Department of Thoracic and Cardiovascular Surgery, Uijeongbu St. Mary's Hospital, College of Medicine, The Catholic University of Korea, 271, Cheonbo-ro, Uijeongbu, Gyeonggi-do, Republic of Korea. Email: medkjj@hanmail.net.

Background: Chest wall re-depression after bar removal (BR) in pectus excavatum (PE) is insufficiently investigated. However, it is not easy to investigate chest wall re-depression due to its multifactorial characteristics. Herein, we investigated chest wall re-depression after BR using machine learning algorithms. To the best of my knowledge, this is the first study of chest wall re-depression after BR using machine learning algorithms.

Methods: We retrospectively reviewed 199 consecutive subjects who underwent both minimally invasive repair of pectus excavatum (MIRPE) and BR at a single hospital from March 2012 to June 2020. We investigated attributes of chest wall re-depression and risk factors for recurrence after BR, predicted final degree and recurrence of PE after BR, and suggested the optimal age at the time of MIRPE based on recurrence. Data for the chest wall re-depression were analyzed to discover differences according to age group [<10 years (early repair group; EG) *vs.* ≥ 10 years (late repair group; LG)].

Results: We observed no significant difference between the Haller index and radiographical pectus index (RPI) ($P=0.431$) and a significant correlation between Haller index and RPI ($P<0.001$). RPI significantly increased for the first 6 months after BR in both age groups (both $P<0.001$) and was maintained at 1 year after BR. RPI value of the LG were significantly higher than those of the EG for the entire period after MIRPE ($P=0.041$). Recurrence of PE in the LG was significantly more frequent than in the EG ($P<0.001$). RPI values before and after MIRPE and age group were identified as independent risk factors for recurrence after BR ($P<0.001$, $P=0.007$, and $P=0.001$, respectively). The linear regression model outperformed for final RPI with performance scores of mean squared error 0.198, root mean squared error 0.445, mean absolute error 0.336, and R^2 0.415. In addition, the logistic regression model outperformed for predicting recurrence with performance scores of 0.865 the area under the curve, 0.884 accuracy, 0.859 F1, 0.865 precision, and 0.884 recall.

Conclusions: The present study shows that machine learning algorithms can provide good estimates for postoperative results in PE. An approach integrating machine learning models and readily available clinical data can be used to create other models in the thoracic surgery field.

Keywords: Machine learning; recurrence; minimally invasive repair of pectus excavatum (MIRPE)

Submitted Sep 11, 2023. Accepted for publication Dec 01, 2023. Published online Jan 29, 2024.

doi: 10.21037/jtd-23-1430

View this article at: <https://dx.doi.org/10.21037/jtd-23-1430>

Introduction

Pectus excavatum (PE) in which the sternum and rib cage are depressed abnormally, is the most common thoracic wall deformity (1,2). Minimally invasive repair of PE (MIRPE) is a standard procedure in which metal bars are placed beneath the sternum and used to lift the depressed chest wall (3,4). Bar placement is generally maintained for 2 to 3 years to guard against re-depression of the chest wall (5,6). The optimal age for MIRPE and recurrence of PE are the most important concerns in treatment because there is a high rate of chest wall re-depression after bar removal (BR) (2,4,5,7,8). The exact mechanisms or characteristics of the chest wall re-depression remain unclear (5,9,10). Furthermore, consensus regarding the definition of recurrence, defined as significant chest wall re-depression after BR, is lacking (2,5,9,10).

In a previous study, we investigated characteristics of chest wall re-depression and risk factors of recurrence of PE after BR (9). However, accurate prediction of chest wall re-depression is crucial for preventing recurrence. The prediction modality should provide predictive numerical value as well as information about risk factors for chest wall re-depression (11-13). However, prior studies are insufficient in prediction of surgical outcomes in PE. The prediction models using machine learning for chest wall re-depression after BR are valuable for better surgical outcomes because machine learning has successfully been used to make accurate decisions and predictions using

diverse and large amounts of data. The primary aims of the present study are to investigate characteristics of chest wall re-depression and to predict final degree and recurrence of PE after BR using machine learning algorithms. To the best of our knowledge, this is the first study in which machine learning analysis was clinically used in PE. We present this article in accordance with the TRIPOD reporting checklist (available at <https://jtd.amegroups.com/article/view/10.21037/jtd-23-1430/rc>).

Methods

All consecutive subjects who underwent both MIRPE and BR at Uijeongbu St. Mary's Hospital from March 2012 to June 2020 were retrospectively reviewed in the present study. MIRPE was conducted according to routine procedures at our institution (3,9). Eligible criteria for subject enrollment were no reoperation due to bar migration, no accompanying thoracic deformity, and no previous thoracic procedures, and no thoracic traumas such as rib, sternal, or spinal fractures. Simple chest posteroanterior and both lateral X-rays were acquired every day during hospitalization and at every outpatient follow-up appointment. Chest computed tomography (CT) scans were routinely acquired for evaluations before MIRPE and on the third postoperative day after both MIRPE and BR. The radiographical pectus index (RPI) was designed to quantify the degree of chest wall depression because a complete series of chest CT scans for the Haller index was not available at outpatient basis during follow-ups (3,5,14). Similar to the Haller index calculated using chest CT, RPI is calculated by the ratio of the transverse length to the posteroanterior length at the most depressed point of the chest wall shown by lateral chest X-rays (*Figure 1*) (5,14,15). We measured the transverse and posteroanterior lengths in double-blind fashion. In the present study, recurrence of PE, or significant chest wall re-depression after BR, is defined when cases met both conditions: increased RPI value and RPI value ≥ 3.5 after BR. Examining the perioperative data during MIRPE, the parameters assumed to be associated with chest wall re-depression after BR were age at the time of MIRPE, sex, RPI values, pectus deformity type (symmetric or asymmetric), sternal angle of bar placement, and number of inserted bars. The sternal angle of bar placement was defined as the angle formed by the sternum and the pectus bar after MIRPE. Previous studies verified that RPI measured by chest X-ray can be used instead of the Haller index measured by chest CT to

Highlight box

Key findings

- Machine learning algorithms using risk factors and basic clinical data yields good performance for prediction of final radiographical pectus index (RPI) and recurrence after bar removal (BR).

What is known and what is new?

- This is the first study in which machine learning analysis is clinically used to evaluate treatment for pectus excavatum.
- RPI values before and after minimally invasive repair of pectus excavatum (MIRPE) and age at the time of MIRPE are independent risk factors for recurrence after BR.

What is the implication, and what should change now?

- Machine learning approaches are convenient and useful for predicting outcomes of MIRPE. In addition, an approach integrating machine learning models and readily available clinical data can be used to create other models in the thoracic surgery field.




Figure 1 Radiographic pectus index. The anteroposterior length between that sternum and the anterior aspect of the corresponding spine is measured at the most depressed point of chest wall. The transverse length was measured at the level of the same spine level. The radiographic pectus index is calculated by the ratio of the transverse length to the anteroposterior length at the most depressed point of chest wall shown by chest X-rays. A, the anteroposterior length; R, right side of chest posteroanterior view; L, the transverse length; Lpa, left side of chest posteroanterior view.

describe degree of PE (5,9,14). We collected and analyzed data for these clinical parameters during treatment of PE.

There is no consensus regarding recurrence after BR (7,9). In the present study, we dealt with this problem as follows. First, we investigated the patterns or attributes of chest wall re-depression presented by RPI after BR. Second, we investigated risk factors for recurrence according to our definition of PE recurrence and predicted postoperative final RPI and recurrence 1 year after BR using perioperative data during MIRPE. Third, to determine the optimal age at the time of MIRPE, we divided the subjects into two groups by age [<10 years (early repair group; EG) *vs.* ≥ 10 years (late repair group; LG)] when they underwent MIRPE and examined the relationship between age and recurrence.

Statistical analysis

All data are shown as mean \pm standard deviation (SD). Comparisons between groups were analyzed using Student's *t*-test or paired *t*-test. To investigate relationship between two quantitative continuous variables, we conducted Pearson correlation analysis. Chi-square tests were used to examine whether two categorical variables were independent in influencing the test statistic. Repeat measures analysis of variance test was performed to investigate changes of chest wall re-depression after BR. The binary logistic regression

(LoR) test (backward, stepwise approach) was performed to investigate independent influencing factors for recurrence after BR. The Statistical Package for the Social Sciences version 22.0 (SPSS, IBM Corp, Armonk, NY, USA) was used to perform statistical evaluations. A *P* value <0.05 was considered statistically significant.

Machine learning analysis

Machine learning analyses were performed to predict the final postoperative RPI and recurrence after BR. The training and validation sets were used for model learning and 10-fold stratified cross-validation was used for training and validation. Performance scores of each model were calculated and evaluated based on area under the curve (AUC), accuracy, precision, F1 score, and recall. The outperformed model was selected based on AUC. Five machine learning algorithms [linear regression (LiR), neural network (NN), AdaBoost (AB), random forest (RF), and decision tree (DT)] were performed to predict the final postoperative RPI, and five machine learning algorithms [LoR, NN, Naïve Bayes (NB), RF, and support vector machine (SVM)] were used to predict recurrence. The Orange[®] data mining toolbox in Python (Bioinformatics Lab at University of Ljubljana, Slovenia) was used for machine learning models learning (16). This open-source platform




Figure 2 Flow chart of subjects.

is a machine learning and data mining suite for data analysis through Python scripting and visual programming (16). The detailed structure for each model includes LiR [regularization type: Lasso (L1), strength $\alpha=0.0001$], NN [neurons in the hidden layers: 100, 50, 20; activation: rectified linear unit (ReLU); solver: Adam; regularization $\alpha=0.0001$, and maximum number of iterations: 200, replicable training], AB (Bae estimator: Tree; number of estimators: 50; learning rate: 1.00000; classification algorithm: SAMME.R; regression loss function: linear), RF (number of trees: 10; replicable training; do not split subsets smaller than 5), DT (induce binary tree; minimum number of instances in leaf: 2; do not split subsets smaller than 5; limit maximum tree depth: 100; and stop when majority reaches: 95%), LoR (regularization type: ridge L2; strength $C=1$), and SVM (cost: 1.0; regression loss epsilon: 0.10; kernel: radial basis function; optimization numerical tolerance: 0.0010; and optimization iteration limit: 100).

Ethical statement

The study conformed to the provisions of the Declaration of Helsinki (as revised in 2013). Uijeongbu St. Mary's Hospital Ethics Committee reviewed and approved the present study (No. UC23RISI0109). The requirement for informed consent was waived because the present study was retrospective without exposure of patient information.

Results

One hundred ninety-nine patients were enrolled into the present study (*Figure 2*). The mean age of the study participants was 11.61 ± 6.84 years at the age of MIRPE. The mean duration of bar placement and total observation were 28.5 ± 5.08 and 44.1 ± 6.63 months, respectively. One hundred fifty-seven males and 42 females were enrolled. The pectus types comprised 158 symmetric and 41 asymmetric cases. The mean Haller index and RPI before MIRPE were 4.20 ± 1.39 and 4.10 ± 2.00 , respectively. The mean RPI before BR and 1 year after BR were 2.55 ± 0.46 and 2.93 ± 0.58 , respectively. All subjects showed lower RPI 1 year after BR then before MIRPE. However, 26 of the 199 cases showed recurrence 1 year after BR. The overall clinical characteristics of the study subjects are shown in *Table 1*.

Validity of RPI for degree of PE

Degree of PE is generally described by the Haller index and repair of PE is usually considered with the Haller index ≥ 3.25 (2,5,17). To validate usage of RPI, we compared the Haller index values and RPI before MIRPE. As in previous studies, we found no significant difference between the Haller index and RPI ($P=0.431$) and a significant correlation between the Haller index and RPI ($P<0.001$) (5,9,14).

Table 1 The overall clinical characteristics of the study population

Variables	Value
Age (years) [†]	11.61±6.84
Sex	
Male	157
Female	42
Morphology type	
Symmetric	158
Asymmetric	41
Age group	
EG (<10 years old)	105
LG (≥10 years old)	94
Recurrence	
No	173
Yes	26
Initial severity of pectus excavatum	
Haller index	4.20±1.39
Radiographic pectus index	4.10±2.00
Number of bars used	
One	128
Two	69
Three	2

Values are shown as mean ± SD or number. [†], at the age of MIRPE. EG, early repair group; LG, late repair group; MIRPE, minimally invasive repair of pectus excavatum; SD, standard deviation.

Table 2 Radiographic pectus index before and after MIRPE according to age group

Age group	Radiographic pectus index		
	Before MIRPE	Immediately after MIRPE	Immediate before BR
EG (<10 years old)	3.90±1.15	2.46±0.50	2.36±0.34
LG (≥10 years old)	4.33±2.64	2.69±0.31	2.76±0.49
P	0.130	0.001	<0.001

Values are shown as mean ± SD. MIRPE, minimally invasive repair of pectus excavatum; EG, early repair group; LG, late repair group; BR, bar removal; SD, standard deviation.

Table 3 Serial changes of radiographic pectus index after BR

Age group	Radiographic pectus index					F	P [†]
	Immediate before BR	Immediate after BR	One month after BR	Six months after BR	One year after BR		
EG (<10 years old)	2.36±0.34	2.50±0.36	2.65±0.41	2.72±0.46	2.72±0.43	80.04	<0.001
LG (≥10 years old)	2.76±0.49	3.02±0.53	3.13±0.55	3.17±0.56	3.15±0.65	2.54	0.041
P	<0.001	<0.001	<0.001	<0.001	<0.001		

Values are shown as mean ± SD. [†], Pillai's trace P value. EG, early repair group; LG, late repair group; BR, bar removal.

Chest wall re-depression after BR

Serial changes of RPI were shown to describe chest wall re-depression at the time points after BR (immediate, 1 month, 6 months, and 1 year) based on age group. RPI significantly increased for the first 6 months after BR in both age groups (both $P<0.001$) but was maintained at 1 year after BR. RPI values of the LG were significantly higher than those in the EG for the entire period after MIRPE ($P=0.041$) (Tables 2,3). As in our previous study, chest wall re-depression progressed only for the first 6 months (Figure 3) (9). In addition, serial changes of RPI were investigated at subsequent time points after BR (immediate, 1 month, 6 months, and 1 year) based on recurrence. RPI values of the recurrent group (RG) were significantly higher than those of the non-recurrent group (NG) for entire period after BR ($P<0.001$). Interestingly, RPI of the RG continuously increased after BR while RPI of the NG did not increase after 6 months following BR (Table 4 and Figure 4). These findings suggest that the pattern of chest wall re-depression after BR can differ according to recurrence.

Recurrence of PE

We assume that MIRPE is best performed early, before 10 years of age. Recurrence of PE occurred in five patients in the EG and 21 cases in the LG after BR, a significant difference ($P<0.001$). Binary logistic regression tests were




Figure 3 Chest wall re-depression after bar removal based on age group. EG, early repair group; LG, late repair group; BR, bar removal.




Figure 4 Chest wall re-depression after bar removal based on recurrence. NG, non-recurrent group; RG, recurrent group; BR, bar removal.

Table 4 Serial changes of radiographic pectus index after BR according to recurrence

Group	Radiographic pectus index					F	P [†]
	Immediate before BR	Immediate after BR	One month after BR	Six months after BR	One year after BR		
EG (<10 years old)	2.45±0.35	2.64±0.41	2.75±0.41	2.80±0.40	2.77±0.37	221.38	<0.001
LG (≥10 years old)	3.18±0.63	3.50±0.52	3.72±0.54	3.82±0.62	4.00±0.66	39.68	<0.001
P	<0.001	<0.001	<0.001	<0.001	<0.001		

Values are shown as mean ± SD. [†], Greenhouse-Geisser P value. EG, early repair group; LG, late repair group; BR, bar removal.

performed to investigate independent risk factors for recurrence of PE. Recurrence of PE at 1 year after BR was defined as an event and age group, sex, RPI values before and after MIRPE, pectus type (symmetric *vs.* asymmetric), number of bars placed, and angle of sternum were included as covariates. Accordingly, RPI values before and after MIRPE and age group were identified as independent risk factors for recurrence after BR ($P < 0.001$, $P = 0.007$, and $P = 0.001$, respectively). The results of multivariate analysis of risk factors are shown in *Table 5*.

Prediction of final RPI and recurrence of PE using machine learning models

Five machine learning algorithms were performed to predict the final postoperative RPI and recurrence of PE 1 year after BR, respectively. Perioperative data on age group, sex, RPI values before and after MIRPE, pectus types (symmetric *vs.* asymmetric), number of bars used, and angle of sternum were included as covariates in the machine learning algorithms. The LiR model outperformed other

models, with performance scores of mean squared error 0.198, root mean squared error 0.445, mean absolute error 0.336, and R^2 0.415 (*Table 6*). In addition, age group, sex, RPI values before and after MIRPE, pectus types (symmetric *vs.* asymmetric), number of bars used, and angle of sternum were included as covariates for prediction of postoperative recurrence of PE. The LoR model outperformed other models for recurrence of PE, with performance scores of AUC 0.865, accuracy 0.884, F1 0.859, precision 0.865, and recall 0.884 (*Table 7, Figures 5,6*).

Discussion

Chest wall re-depression after BR is common and remains a concern despite otherwise generally excellent outcomes of MIRPE (5,7,9). This problem highlights several issues associated with MIRPE (4,18). The first regards questions about mechanism and characteristics of chest wall re-depression after BR. However, the mechanisms and characteristics of chest wall re-depression after BR remain unclear (5). Because chest wall re-depression is

Table 5 The multivariate analysis of risk factors for the recurrence of pectus excavatum after bar removal

Variables	Odds ratio	95% confidence interval	P
LG (≥ 10 years old)	11.663	2.747–48.682	0.001
RPI before MIRPE	2.970	1.717–5.138	<0.001
RPI after MIRPE	3.894	1.454–10.428	0.007

LG, late repair group; RPI, radiographic pectus index; MIRPE, minimally invasive repair of pectus excavatum.

Table 6 Prediction of final radiographic pectus index of pectus excavatum using machine learning models

Algorithms	MSE	RMSE	MAE	R ²
Random forest	0.220	0.470	0.358	0.350
Neural network	0.617	0.786	0.411	-0.821
Linear regression	0.198	0.445	0.336	0.415
AdaBoost	0.258	0.508	0.393	0.238
Decision tree	0.419	0.648	0.478	-0.237

MSE, mean squared error; RMSE, root mean squared error; MAE, mean absolute error.

Table 7 Prediction of final radiographic pectus index and recurrence of pectus excavatum using machine learning models

Algorithms	AUC	Accuracy	F1	Precision	Recall
Random forest	0.738	0.844	0.826	0.814	0.844
Support vector machine	0.701	0.874	0.820	0.890	0.874
Logistic regression	0.865	0.884	0.859	0.865	0.884
Neural network	0.834	0.889	0.871	0.873	0.889
Naïve Bayes	0.847	0.854	0.863	0.876	0.854

AUC, area under the curve.




Figure 5 Receiver operating characteristic curve for prediction of recurrence after bar removal using the logistic regression model shows that default threshold (0.5) point is 0.487. Solid line: performance line; dotted line: base line. TP, true positive; FP, false positive.

multifactorial and spatial, clinically available methods or modalities to evaluate and analyze chest wall re-depression are needed (19,20). However, comprehensive evaluations of chest wall depression are difficult (15,19,21). We examined patterns of chest wall re-depression after BR and identified risk factors for recurrence. Chest wall re-depression generally deteriorates only for the first 6 months after BR. This finding suggest that recurrence cannot be diagnosed immediately after BR, and that final diagnosis of recurrence should occur 6 months after BR. The second problem is the defining recurrence of PE, for which there is no consensus (7,9,10). We suggest a definition of recurrence of PE in the present study and create models to predict recurrence. The third problem is the question of the optimal age for MIRPE (8-10). In the present study, we found that the recurrence rate was lower in the EG than in the LG, although the




Figure 6 Nomogram model for predicting recurrence of pectus excavatum after bar removal. RPI, radiographic pectus index; Preop, preoperative; Postop, postoperative.

change of RPI after BR showed similar patterns between groups. The lower rate of recurrence in the EG was explained by RPI values before and after MIRPE and age group, which were identified as independent risk factors of recurrence after BR. RPI values before and after MIRPE was lower in the EG than in the LG, although the degree and pattern of chest wall re-depression after BR did not differ by age. These findings are explained by the reliance of MIRPE on chest wall pliability, which may weaken with age (3,8). We conclude that early repair of PE better than late repair with respect to recurrence, and the corrective effect of MIRPE was higher in the EG than in the LG. The fourth concern is prevention of recurrence of PE. To prevent chest wall re-depression and to obtain persistent repair after BR, it is necessary to determine the details of surgical technique, to understand the mechanism of chest wall re-depression, and to investigate risk factors (4,7,9,22). The details of appropriate surgical techniques have achieved a wide consensus among surgeons even if they have minor variations (4,8,22). However, the most important criterion, the degree of repair remains undefined (19). Sufficient elevation of the depressed chest wall is essential to prevent recurrence of PE because RPI values after MIRPE is the most important factor for recurrence (5). Because patients in the LG have higher RPI than those in the EG, we suggest that depressed chest walls in the LG should be lifted more than in the EG to prevent recurrence. Risk factors for recurrence and proper techniques to prevent

recurrence have been suggested in previous studies (4,9). However, the results of those studies were inconclusive, and other novel approaches to prevent recurrence are needed. The prediction of recurrence is crucial for preventing recurrence, and the prediction modality should provide information about influencing factors and predictive value for recurrence (12,16,23). Therefore, machine learning algorithms were used in the present study to estimate numerical probability values for recurrence after BR. Machine learning has successfully been used to make accurate decisions and predictions using diverse and large amounts of data (11,16). Models using machine learning algorithms can be used to predict RPI and recurrence after BR using perioperative data during MIRPE. Degree of chest wall depression immediately before and after MIRPE and basic clinical data (age, sex, number of bars used, and sternal angle) were included as variables in the machine learning algorithms (5,9). The prediction model for final RPI and recurrence after BR yielded good outcomes, which is meaningful in real clinical practice. To the best of our knowledge, this is the first study of recurrence after BR using machine learning algorithms and additional studies are required to validate our findings. The approach of integrating machine learning models and readily available clinical data can be used to develop other models for treatment of PE (11).

The present study has several limitations. First, it was a retrospectively study in a single center. The

generalizability of the models to other institutions should be studied. Second, the definition of recurrence we used might affect our results. Third, measuring degree of chest wall depression by simple radiography could result in measurement errors that could be influenced by physiologic factors, especially in the asymmetric type. Fourth, the number of subjects in the dataset was relatively small, but we used strict eligibility criteria and study design to obtain data of good quality. PE cannot be simply and strictly defined into two types (symmetric *vs.* asymmetric) because it has many variant morphologies. In addition, RPI cannot describe the complete degree of PE. The heterogeneity in PE type and incomplete descriptions of degree reflect real-world situations, and machine learning models trained with such data may be more appropriate for clinical practice. PE occupies a minor portion of case in the thoracic surgery field, and knowledge regarding recurrence of PE is valuable because recurrence can have profound effects on patients despite its characterization as a benign condition.

Conclusions

The findings of the present study showed that machine learning algorithms considering risk factors and basic clinical data yield good performance for prediction of final RPI and recurrence after BR. Such machine learning approaches can be convenient and provide accurate decisions in management of PE. Further large-scale studies are required to validate the findings of the present study and suggest surgical details for MIRPE. In addition, an approach integrating machine learning models and readily available clinical data can be used to create other models in the thoracic surgery field.

Acknowledgments

Funding: None.

Footnote

Reporting Checklist: The authors have completed the TRIPOD reporting checklist. Available at <https://jtd.amegroups.com/article/view/10.21037/jtd-23-1430/rc>

Data Sharing Statement: Available at <https://jtd.amegroups.com/article/view/10.21037/jtd-23-1430/dss>

Peer Review File: Available at <https://jtd.amegroups.com/>

[article/view/10.21037/jtd-23-1430/prf](https://jtd.amegroups.com/article/view/10.21037/jtd-23-1430/prf)

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://jtd.amegroups.com/article/view/10.21037/jtd-23-1430/coif>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The study conformed to the provisions of the Declaration of Helsinki (as revised in 2013). Uijeongbu St. Mary's Hospital Ethics Committee reviewed and approved the present study (No. UC23RISI0109). The requirement for informed consent was waived because the present study was retrospective without exposure of patient information.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Kloth K, Klohs S, Bhullar J, et al. The Epidemiology behind Pectus Excavatum: Clinical Study and Review of the Literature. *Eur J Pediatr Surg* 2022;32:316-20.
2. Janssen N, Daemen JHT, van Polen EJ, et al. Pectus Excavatum: Consensus and Controversies in Clinical Practice. *Ann Thorac Surg* 2023;116:191-9.
3. Toci GR, Davis TA, Bigelow BF, et al. Analyzing Outcomes of Nuss and Ravitch Repair for Primary and Recurrent Pectus Excavatum in Adults. *Ann Thorac Surg* 2020;110:272-5.
4. Haecker FM, Krebs TF, Kleitsch KU. Current Development of Minimally Invasive Repair of Pectus Excavatum (MIRPE). *Children (Basel)* 2022;9:478.
5. Kuyama H, Uemura S, Soh H, et al. Anterior chest wall regression after Nuss bar removal in adult patients with pectus excavatum. *Gen Thorac Cardiovasc Surg* 2021;69:1308-12.
6. Haecker FM, Hebra A, Ferro MM. Pectus bar removal -

- why, when, where and how. *J Pediatr Surg* 2021;56:540-4.
7. Wang L, Bi R, Xie X, et al. A Modified Nuss Procedure for Recurrent Pectus Excavatum of Adults. *Front Surg* 2021;8:814837.
 8. de Loos ER, Pennings AJ, van Roozendaal LM, et al. Nuss Procedure for Pectus Excavatum: A Comparison of Complications Between Young and Adult Patients. *Ann Thorac Surg* 2021;112:905-11.
 9. Cho DG, Kim JJ, Park JK, et al. Recurrence of pectus excavatum following the Nuss procedure. *J Thorac Dis* 2018;10:6201-10.
 10. Kuyama H, Uemura S, Yoshida A. Recurrence of pectus excavatum in long-term follow-up after the Nuss procedure in young children based on the radiographic Haller index. *J Pediatr Surg* 2020;55:2699-702.
 11. Daemen JHT, Heuts S, Rezazadah Ardabili A, et al. Development of Prediction Models for Cardiac Compression in Pectus Excavatum Based on Three-Dimensional Surface Images. *Semin Thorac Cardiovasc Surg* 2023;35:202-12.
 12. Tse KM, Tan LB, Lee SJ, et al. Feasibility of using computer simulation to predict the postoperative outcome of the minimally invasive Nuss procedure: Simulation prediction vs. postoperative clinical observation. *J Plast Reconstr Aesthet Surg* 2018;71:1496-506.
 13. Sakamoto Y, Yokoyama Y, Nagasao T, et al. Outcomes of the Nuss procedure for pectus excavatum in adults. *J Plast Reconstr Aesthet Surg* 2021;74:2279-82.
 14. Matsuura R, Tazuke Y, Ueno T, et al. Can plain chest X-ray replace computed tomography for the follow-up of children who have undergone the Nuss procedure? *Surg Today* 2020;50:1249-54.
 15. Silva B, Pessanha I, Correia-Pinto J, et al. Automatic Assessment of Pectus Excavatum Severity From CT Images Using Deep Learning. *IEEE J Biomed Health Inform* 2022;26:324-33.
 16. Demsar J, Curk T, Erjavec A, et al. Orange: Data Mining Toolbox in Python. *J Mach Learn Res* 2013;14:2349-53.
 17. Wei B, Gleason F. Commentary: In With The New: Three-Dimensional Surface Imaging For Pectus Excavatum. *Semin Thorac Cardiovasc Surg* 2022;34:1376-7.
 18. Abbas AE. Commentary: Funneling the funnel chest debates into appropriate management of adult pectus excavatum. *JTCVS Tech* 2022;12:220-1.
 19. Uemura S, Yoshida A, Kuyama H. Analysis of chest wall elevation after the Nuss procedure using 3D body scanning technique in patients with pectus excavatum. *Pediatr Surg Int* 2021;37:777-82.
 20. Torre M, Genova Gaia L, Calevo MG, et al. Blood metal levels after minimally invasive repair of pectus excavatum. *Interact Cardiovasc Thorac Surg* 2021;33:76-81.
 21. Coorens NA, Daemen JHT, Slump CH, et al. The Automatic Quantification of Morphological Features of Pectus Excavatum Based on Three-Dimensional Images. *Semin Thorac Cardiovasc Surg* 2022;34:772-81.
 22. Janssen N, Daemen JHT, Ashour O, et al. Nuss bar removal without straightening is a safe technique: a single center experience. *J Thorac Dis* 2022;14:3335-42.
 23. Coorens NA, Daemen JHT, Slump CH, et al. Predicting Aesthetic Outcome of the Nuss Procedure in Patients with Pectus Excavatum. *Semin Thorac Cardiovasc Surg* 2023;35:627-37.

Cite this article as: Hyun K, Kim JJ, Choi WK, Kim YH, Han SC. Prediction of postoperative final degree and recurrence of pectus excavatum using machine learning algorithms. *J Thorac Dis* 2024;16(1):311-320. doi: 10.21037/jtd-23-1430