



Grape seed proanthocyanidin extract induces apoptosis of HL-60/ADR cells via the Bax/Bcl-2 caspase-3/9 signaling pathway

Ka-Na Lin^{1,2,3^}, Wei Zhao^{1^}, Shi-Ying Huang^{1^}, Hao Li^{1,2^}

¹Department of Pharmacy, Shanghai Children's Medical Center, School of Medicine, Shanghai Jiao Tong University, Shanghai, China; ²Clinical Research Center, Shanghai Children's Medical Center, School of Medicine, Shanghai Jiao Tong University, Shanghai, China; ³Center for Brain Science, Shanghai Children's Medical Center, School of Medicine, Shanghai Jiao Tong University, Shanghai, China

Contributions: (I) Conception and design: KN Lin; (II) Administrative support: W Zhao; (III) Provision of study materials or patients: KN Lin, W Zhao; (IV) Collection and assembly of data: KN Lin, SY Huang; (V) Data analysis and interpretation: KN Lin, SY Huang, H Li; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Shi-Ying Huang. Department of Pharmacy, Shanghai Children's Medical Center, School of Medicine, Shanghai Jiao Tong University, No. 1678 Dongfang Road, Pudong Area, Shanghai, 200127, China. Email: shiying.h@outlook.com; Hao Li. Department of Pharmacy, Shanghai Children's Medical Center, School of Medicine, Shanghai Jiao Tong University, No. 1678 Dongfang Road, Pudong Area, Shanghai, 200127, China. Email: lihao19880810@hotmail.com.

Background: Our previous study detailed the direct induction of apoptosis by grape seed proanthocyanidin extract (GSPE) in a multidrug resistant human acute myeloid leukemia (AML) HL-60/adriamycin (HL-60/ADR) cell line, although the mechanism of this effect was not detailed. This study aims to elucidate the mechanism underlying GSPE-induced cell apoptosis in HL-60/ADR cells.

Methods: HL-60/ADR cells were studied to evaluate effects of GSPE (0–100 µg/mL); a 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay was employed to identify the cytotoxic effect of varying GSPE concentrations. Trypan blue staining was used to observe changes in cell viability; flow cytometry assays were used to verify apoptosis. Expression of Bax and Bcl-2 mRNA was analyzed using real-time polymerase chain reaction (PCR). Activity of caspase-3 and caspase-9 was also detected.

Results: Here, GSPE was found to inhibit HL-60/ADR cell growth and induce cell apoptosis in a dose-dependent manner. Real-time PCR findings revealed that GSPE concentrations above 75 µg/mL significantly increase expression of Bax mRNA ($P < 0.001$). GSPE concentrations above 25 µg/mL were found to significantly decrease expression of Bcl-2 mRNA ($P < 0.01$), while concentrations above 50 µg/mL were found to significantly increase caspase-3 activity after 6, 12 and 24 h ($P < 0.01$). However, only 100 µg/mL GSPE was found to significantly increase caspase-9 activity ($P < 0.001$ at 6 and 12 h; $P < 0.05$ at 24 h).

Conclusions: GSPE inhibits the proliferation of HL-60/ADR cells by the induction of apoptosis in a dose-dependent manner via the Bax/Bcl-2 caspase-3/9 signaling pathway.

Keywords: Grape seed proanthocyanidin extract (GSPE); apoptosis; acute myeloid leukemia (AML); Bax/Bcl-2; caspase

Submitted May 28, 2021. Accepted for publication Aug 09, 2021.

doi: 10.21037/tcr-21-920

View this article at: <https://dx.doi.org/10.21037/tcr-21-920>

[^] ORCID: Ka-Na Lin, 0000-0002-9374-8901; Wei Zhao, 0000-0003-1722-4521; Shi-Ying Huang, 0000-0002-4669-0057; Hao Li, 0000-0003-2832-4851.

Introduction

Acute myeloid leukemia (AML) is a hematopoietic malignancy (1,2). It is the commonest type of acute leukemia in adults with an incidence of 50 patients per 1 million and a median diagnostic age of 68 y. This condition is very rare in pediatric patients, with an incidence of about 7 per 1 million children annually (1,3,4). With the approval of novel drugs in recent years, the frontline management of AML is rapidly changing. If tolerated, however, intensive chemotherapy remains the general therapeutic choice for patients suffering newly-diagnosed AML (5). A combination of cytarabine and an anthracycline (such as daunorubicin or idarubicin) has been the therapeutic standard of AML for several decades (1). Resistance to cytotoxic chemotherapy leads to AML relapse, resulting in a very low long-term survival rate ranging from 10–15% among patients aged 60 years and older (6). In recent years, many naturally-derived products such as parthenolide, triptolide and resveratrol have been found efficacious in the targeting of multiple leukemic stem cell pathways, resulting in AML remission (7). As such, increasing attention is being paid to the role of natural products in AML treatment.

Grape seed proanthocyanidin extract (GSPE), a compound extracted from grape seeds, is composed of flavane-3-alcohol monomers, dimers, trimers, and tetramers, as well as oligomers of proanthocyanidins, catechin, epicatechin gallate and epigallocatechin (8,9). Due to its significant health-promoting effects and potential clinical value, GSPE has attracted considerable interest in the recent year. The compound has been reported to promote skeletal muscle fiber transformation resulting in improvement of fatigue (10), enhance L-cell differentiation in intestinal organoids thus promoting hormone production in the treatment of type-2 diabetes and obesity (11–13), reverse multidrug resistance to cytotoxic agents in the setting of AML (14), inhibit secretion of complement component 3 thereby relieving renal interstitial inflammation and fibrosis (15) and exert neuroprotective effects in neonatal hypoxic-ischemic brain injury (16). A randomized, double-blind, placebo-controlled clinical study documented the benefit of GSPE on maintaining vascular endothelial function and normal blood pressure in participants with prehypertension (17). Furthermore, GSPE has also been reported to exert anti-cancer effects in malignancies such as esophageal squamous carcinoma (18), hepatocellular carcinoma (19), breast cancer (20), colon

cancer (21) and AML (22). We previously documented the potential application of GSPE in combination with cytotoxic drugs for the treatment of AML (14). GSPE reverses drug resistance to both cytarabine and doxorubicin [also as known as adriamycin (ADR)] in HL-60/ADR cells via inhibition of the PI3K/Akt signaling pathway, resulting in the downregulated expression of multidrug resistance protein 1 (MDR1), multidrug resistance-associated protein 1 (MRP1) and lung resistance-related protein (LRP) (14). In addition, we previously found that GSPE directly mediates apoptosis in HL-60/ADR cells (14). Here, we aimed to elucidate the mechanism via which GSPE induces apoptosis in HL-60/ADR cells. We present the following article in accordance with the MDAR checklist (available at <https://dx.doi.org/10.21037/tcr-21-920>).

Methods

Cell lines and reagents

ADR-resistant human AML HL-60/ADR and parental HL-60 cells were purchased from the Institute of Hematology, Chinese Academy of Medical Sciences (Tianjin, China). Both HL-60 and HL-60/ADR cells were cultured in Roswell Park Memorial Institute (RPMI 1640; Gibco, Grand Island, USA) medium supplemented with 10% fetal bovine serum (FBS; purchased from Hangzhou Sijiqing Bioengineering Material Co. Ltd., Hangzhou, China) and maintained in a humidified incubator at 37 °C and 5% CO₂. Cells were routinely passaged every other day. The HL-60/ADR cells were maintained in medium containing 1,000 ng/mL ADR (Shenzhen Main Luck Pharmaceuticals Co. Ltd., Shenzhen, China) to maintain an MDR phenotype and incubated in ADR-free medium for 2 weeks prior to experimentation.

GSPE with a purity exceeding 95% was purchased from Tianjin Jianfeng Natural Product R&D Co. Ltd. (Tianjin, China; Manufacturer's content Report) while 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) was purchased from Sigma (St Louis, USA). Trizol reagents were purchased from Invitrogen (Carlsbad, USA). The Annexin V-Fluorescein Isothiocyanate/Propidium Iodide (FITC/PI) Apoptotic Cell Detection Kit was purchased from BD Biotech (Franklin Lakes, USA). The PrimeScript™ RT Reagent Kit with gDNA Eraser and TB Green® Premix EX Taq™ (Tli RnaseH Plus) were purchased from Takara Biotechnology Inc (Dalian, China). Primer sequence design and other

Table 1 Bcl-2 and Bax mRNA primer sequences

mRNA	Primer (5'→3')
Bcl-2 (Forward)	AGACCGAAGTCCGCAGAACC
Bcl-2 (Reverse)	GAGACCACACTGCCCTGTTG
Bax (Forward)	GCGACTGATGTCCCTGTCTC
Bax (Reverse)	GGCCTCAGCCCATCTTCTTC

reagents were purchased from Sangon Biotech Inc (Shanghai, China). The Caspase-3/9 Activity Detection Kit was purchased from Beyotime Biotechnology Inc (Shanghai, China).

Cell viability assay

Cells were plated in 96-well plates at 5×10^4 cells/well (100 μ L of cell suspension with a concentration of 5×10^5 cells/mL) and treated with various doses (6.25, 12.5, 25.0, 50.0, 75.0, 100.0, 125.0, 150.0 and 200.0 μ g/mL) of GSPE for 24, 48 and 72 h. The total liquid volume was 150 μ L. Then, 50 μ L of MTT solution (2 mg/mL) was added to each well at 37 °C for another 4 h. Plates were centrifuged (Eppendorf, Hamburg, Germany) for 10 min at 2,500 rpm at room temperature; the supernatant fraction was subsequently discarded. In order to dissolve formazan, DMSO was added at 100 μ L/well and mixed thoroughly. Absorbance (OD value) was read at 570 nm using a microplate reader (Bio-Rad Laboratories, Hercules, USA). All experiments were repeated 3 times. The cell viability rate was calculated according to the following formula: cell viability (%) = $OD_{570, GSPE} / OD_{570, control} \times 100\%$.

Trypan blue staining for viable cell counts

Cells were seeded in 24-well plates at 1.5×10^5 cells per well (500 μ L of cell suspension with a concentration of 3×10^5 cells/mL) and treated with different concentrations (0, 25, 50, 75, 100 μ g/mL) of GSPE for 24 h. Cell counts and viability were measured using Trypan blue staining and automated cell counting (Countstar[®] BioMarine, Shanghai Ruiyu Biotechnology Co. Ltd., Shanghai, China). Three duplicate wells were set up for each group studied and cells from each well were counted 3 times.

Apoptosis analysis by flow cytometry

Treated cells were harvested and resuspended in PBS

solution at a concentration of 1×10^6 cells/mL; 100 μ L of cells were then placed into Eppendorf tubes and mixed with 400 μ L binding buffer. After Annexin V-FITC staining for 15 min and PI staining for 5 min at room temperature in the dark, samples were analyzed using a BD FACS Verse Flow cytometer (BD Biosciences, San Jose, USA) and Flow Jo Software (Tristar, Washington DC, USA). Cells in the Q2 phase (Annexin V-FITC +/PI -) were considered to be in the late stage of apoptosis, while cells in the Q4 phase (Annexin V-FITC +/PI +) were considered to be in the early stage of apoptosis. All experiments were performed independently in triplicate.

Real-time PCR analysis for the quantification of Bcl-2 and Bax mRNA expression

Cells were seeded in 6-well plates at 1.0×10^6 cells/well (2 mL of cell suspension with a concentration of 1×10^6 cells/mL) and treated with different concentrations (0, 25, 50, 75, 100 μ g/mL) of GSPE for 24 h. Total RNA of all treated cells, including attached cells as well as cells in the supernatant, was extracted using Trizol reagent and reverse transcribed into cDNA using the RT Reagent Kit. Real-time PCR was performed using a Biosystems 7500 Fast Real-Time PCR (Bio-Rad, Laboratories, Hercules, USA) according to the manufacturer's protocol. Primer sequences are shown in Table 1. Reaction conditions were as follows: 1 cycle of 20 s at 95 °C followed by 40 cycles of 3 s at 95 °C and 15 s at 60 °C. β -actin was used as an internal control. The relative expression of genes investigated was determined by the $2^{-\Delta\Delta CT}$ method.

Caspase-3/9 activity analysis

Caspase-3 and caspase-9 activity was evaluated using the Caspase-3/9 Activity Assay Kit according to the manufacturer's protocol. Briefly, treated cells were extracted and incubated in lysis buffer on ice for 15 min. The mixture was subsequently centrifuged at 16,000 g for 15 min at 4 °C. Supernatant was collected and incubated with Ac-DEVD-pNA and Ac-LEHD-pNA for 2 h. Absorbance was measured at 405 nm using a microplate reader. Caspase-3/9 activity of each sample was quantified according to the standard curve and normalized by protein concentration.

Statistical analysis

All experiments were performed three times. Data are




Figure 1 GSPE directly inhibits HL-60 and HL-60/ADR cell proliferation. (A) Cell validity of different GSPE concentrations. (B) Effects of different GSPE concentrations (0, 6.25, 12.5, 25.0, 50.0, 75.0, 100.0, 125.0, 150.0 and 200.0 µg/mL) on the survival rate of HL-60/ADR cells cultured for 24 h as compared to controls (GSPE 0 µg/mL); (C) Effects of different durations of cell culturing and different GSPE concentrations on HL-60/ADR cell proliferation. ***, $P < 0.001$. ADR, adriamycin; GSPE, grape seed proanthocyanidin extract.

presented as means \pm SD; figures were drawn using Microsoft Excel 2019. Comparisons were performed via one-way ANOVA in GraphPad Prism 7. $P < 0.05$ was considered statically significant.

Results

GSPE inhibits proliferation of HL-60/ADR cells

Our findings indicated that HL-60/ADR cells were resistant to cytarabine, ADR, vincristine, daunorubicin, mitoxantrone, pirarubicin, homoharringtonine and etoposide (14). Here, we re-evaluated the drug resistance of HL-60/ADR cells to ADR (Figure 1A). The IC_{50} of ADR in HL-60 and HL-60/ADR cells was 2.033 and 415.7 µM, respectively, and 204.5 times higher in HL-60/ADR cells than in HL-60 cells, confirming that the HL-60/ADR cells studied here were indeed drug resistant.

To determine whether GSPE directly inhibits HL-60/ADR cellular proliferation and which concentrations are most efficacious, HL-60/ADR cells were cultured with 6.25, 12.5, 25.0, 50.0, 75.0, 100.0, 125.0, 150.0 and 200.0 µg/mL concentrations of GSPE for 24, 48 and 72 h. As shown in Figure 1B, proliferation of HL-60/ADR cells was significantly inhibited after cell culturing with GSPE concentrations above 25.0 µg/mL for 24 h as compared to control samples cultured without GSPE ($P < 0.001$). To verify whether the inhibitory effect of GSPE was irreversible, HL-60/ADR cells were cultured for 72 h with GSPE concentrations of 25, 50, 75 and 100 µg/mL (controls were cultured without GPSE). The optical density (OD) value exhibited a sustainable increase at 48 and 72 h when

the concentration of GSPE was 25 µg/mL, indicating that GSPE does not completely inhibit HL-60/ADR cellular proliferation at this concentration (Figure 1C). When GSPE concentrations were 50, 75 and 100 µg/mL, the OD value (570 nm) of HL-60/ADR cells did not increase at 48 and 72 h as compared to the OD value at 24 h, which indicating that HL-60/ADR cellular proliferation was completely inhibited at GSPE concentrations above 50 µg/mL (Figure 1C).

GSPE induces apoptosis of HL-60/ADR cells

To confirm how GSPE inhibits proliferation of HL-60/ADR cells, we first examined changes in cellular morphology. After 24 h of culturing, cells were stained with trypan blue. As GSPE concentration at which cells were cultured increased, the proportion of apoptotic cells seen on microscopy increased, while cell survival decreased (Figure 2A-2E). The survival rate of cells cultured with GSPE concentrations above 50 µg/mL significantly decreased when compared with controls ($P < 0.001$) (Figure 2F). Thus, morphological observations underscored that GSPE induces HL-60/ADR cell apoptosis.

In order to further verify the apoptotic effects of GSPE in HL-60/ADR cells, cells were cultured with GSPE concentrations of 25, 50, 75 and 100 µg/mL for 24 h prior to analysis using flow cytometry. As shown in Figure 3, flow cytometry data revealed an increase in apoptotic cells associated with increasing GSPE concentrations. Indeed, GSPE concentrations above 25 µg/mL exhibited a stronger apoptotic effect in HL-60/ADR cells when compared to




Figure 2 Effects of different GSPE concentrations on HL-60/ADR cell count under trypan blue staining. (A-E) Observations of apoptotic HL-60/ADR cells cultured with different GSPE concentrations for 24 h under microscopy after trypan blue staining: (A) 0 µg/mL; (B) 25 µg/mL; (C) 50 µg/mL; (D) 75 µg/mL; (E) 100 µg/mL; dark blue cells in each image were apoptotic HL-60/ADR cells. A size scale in the lower right corner of the image shows a 100 µm distance. (F) HL-60/ADR cells survival rates when cultured with different GSPE concentrations. ***, $P < 0.001$ when *vs.* control. GSPE, grape seed proanthocyanidin extract; ADR, adriamycin.

controls; the proportion of apoptotic cells increased in a dose-dependent manner. These findings underscore that GSPE directly induces apoptosis in HL-60/ADR cells.

GSPE induces apoptosis via the alteration of Bax and Bcl-2 mRNA expression in HL-60/ADR cells

To further explore how GSPE induces apoptosis via the classical pathway, expression of Bcl-2 and Bax mRNA in HL-60/ADR cells was studied using real-time PCR. As the concentrations of GSPE increased, so did levels of Bax mRNA expression (Figure 4A). GSPE concentrations above 75 µg/mL were found to be associated with significantly increased Bax mRNA expression ($P < 0.001$). However, Bcl-2 mRNA expression decreased as the concentrations of GSPE increased (Figure 4B). When GSPE concentrations were greater than 25 µg/mL, expression of Bcl-2 mRNA was significantly decreased ($P < 0.01$). The Bax mRNA / Bcl-2 mRNA value was found to be significantly increased when GSPE concentrations were greater than 50 µg/mL ($P < 0.01$; Figure 4C).

GSPE increases activity of caspases-3 and -9

Activity of caspases-3 and -9 were detected at 6, 12 and

24 h after culturing HL-60/ADR cells with varying GSPE concentrations. GSPE concentrations greater than 75 µg/mL were associated with significantly increased caspase-3 activity in HL-60/ADR cells cultured for 6 h ($P < 0.001$); GSPE concentrations greater than 50 µg/mL were associated with significantly increased caspase-3 activity after 12 h (50 µg/mL, $P < 0.05$; 75 µg/mL, $P < 0.01$; 100 µg/mL, $P < 0.001$; Figure 5A). After 24 h, GSPE concentrations greater than 50 µg/mL were found to be associated with significantly increased caspase-3 activity ($P < 0.001$; Figure 5A). However, only a concentration of 100 µg/mL was found to significantly increase caspase-9 activity (Figure 5B). Thus, our findings confirmed that GSPE-induced apoptosis of HL-60/ADR cells occurs via the Bax/Bcl-2 caspase 3/9 pathway.

Discussion

Resistance to cytotoxic chemotherapy remains a challenge in the treatment of cancers such as AML (1). GSPE, a flavonoid polyphenolic compound extracted from grape seeds, has been reported to exhibit antitumor activity in the setting of malignancies such as bladder, breast and lung cancers (23,24). GSPE is composed of both monomeric




Figure 3 Effects of different GSPE concentrations on HL-60/ADR cell apoptosis were detected by Annexin V-FITC/PI double staining. (A-E) Cell apoptosis in different GSPE concentrations: (A) 0 µg/mL; (B) 25 µg/mL; (C) 50 µg/mL; (D) 75 µg/mL; (E) 100 µg/mL; cells in the Q4 phase were considered to be in the early stage of apoptosis while cells in the Q2 phase were considered to be in the late stage of apoptosis. (F) Quantification of cell apoptosis in HL-60/ADR cells cultured with different GSPE concentrations. The proportion of apoptotic cells was calculated by the ratio of Q2 to Q4 quadrant cells. ***, $P < 0.001$ *vs.* control. GSPE, grape seed proanthocyanidin extract; ADR, adriamycin.




Figure 4 Effects of different GSPE concentrations on Bax and Bcl-2 mRNA expression in HL-60/ADR cells. (A) Bax mRNA expression at different GSPE concentrations (0, 25, 50, 75, 100 µg/mL). (B) Bcl-2 mRNA expression at different GSPE concentrations. (C) The ratio of Bax/Bcl-2 at different GSPE concentrations. *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$ *vs.* control (GSPE 0 µg/mL). GSPE, grape seed proanthocyanidin extract; ADR, adriamycin.

and polymeric +catechin, -epicatechin gallate and -epigallocatechin linked together via either C4-C6 or C4-C8 bonds (13,25). According to the structural classification of proanthocyanidins, their commonest constitutive units

can be divided into A-type and B-type procyanidins; GSPE is a B-type procyanidin (26). Our previous study reported that GSPE reverses the resistance of HL-60/ADR cells to cytosine, arabinoside and ADR via the inhibition of the




Figure 5 Effects of different GSPE concentrations on the activity of caspases-3 and -9 in HL-60/ADR cells. (A) Effects of different GSPE concentrations (0, 50, 75, 100 µg/mL) on the activity of caspase-3; (B) effects of different GSPE concentrations (0, 50, 75, 100 µg/mL) on the activity of caspase-9, after HL-60/ADR cells were treated with GSPE for 6, 12 and 24 h. *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$ vs. control (GSPE 0 µg/mL). GSPE, grape seed proanthocyanidin extract; ADR, adriamycin.

PI3K/Akt signaling pathway (14). Interestingly, we found that GSPE directly induces apoptosis in HL-60/ADR cells even without the addition of chemotherapeutic drugs (14). This study was carried out to elucidate how GSPE induces apoptosis in HL-60/ADR cells.

There are 3 major apoptosis-associated pathways resulting in caspase activation: the mitochondrial/apoptosome pathway, the death receptor pathway, and the CTL/NK-derived granzyme B-dependent pathway (27). Caspase activation culminates during cell apoptosis and the inflammatory response (27). Mitochondrial apoptosome-driven caspase activation is a key caspase-activating mechanism initiated by cytotoxic drugs and results in caspase-2, -3, -6, -7, -8, -9, and -10 activation during apoptosis (27). Caspases-3 and -9, however, are the most important during mitochondrial apoptosome-driven caspase activation. The intrinsic mitochondrial pathway of apoptosis is tightly controlled by Bcl-2 family proteins, including Bcl-2, Bcl-XL and MCL-1 (28). While Bcl-2 is an anti-apoptotic protein, Bax is a pro-apoptotic effector protein. Inhibition of Bcl-2 was previously reported to promote apoptosis in leukemic cells (29).

GSPE has been reported to exhibit anticancer activity in the setting of several malignancies including AML (14,22), hepatocellular carcinoma (19), bladder cancer (23), pancreatic cancer (30) and colon cancer (21). Gao *et al.* previously reported that GSPE induces apoptosis in Jurkat cells via the c-Jun NH₂-terminal (JNK) pathway by sustained JNK activation and Cip1/p21 up-regulation that finally culminates in caspase activation (31). Hu *et al.* previously reported that GSPE induces dose-dependent, mitochondria-associated apoptosis in human AML

14.2D10 cells (22). Guo *et al.* found that GSPE inhibits the proliferation of cells of the ECA109 esophageal squamous cancer cell line; apoptosis was reported to be induced via the activation of caspase-3 and attenuation of the activation of the NF-κB signaling pathway (18). In ECA 109 cells, GSPE was also found to promote Bax expression and decrease Bcl-2 expression in both time- and dose-dependent manners (18). Liu *et al.* reported that GSPE-induced apoptosis in cells of the BIU87 human bladder cancer line is associated with increased caspase-3 activity (23). GSPE was found to increase rates of apoptosis in the MIA PaCa-2 pancreatic adenocarcinoma cell line via the downregulation of Bcl-2 (30). In addition, GSPE was reported to induce apoptosis in the CaCo2 colorectal cancer cell line via a mechanism associated with the attenuation of the PI3K/PKB pathway (21). However, Ruan *et al.* (32) reported that GSPE is associated with decreased Bax expression and increased Bcl-2 expression, thus playing a critical role in the inhibition of H9C2 cell apoptosis after hypoxic culturing via the PI3K/AKT signaling pathway. Further beneficial effects exerted by GSPE include the prevention of bone loss induced by inflammation via a decrease in the phosphorylation and degradation of NF-κB inhibitor and inhibition of apoptosis via a decrease in the activity of caspases-3 and -9 (33).

Here, we confirmed that GSPE directly induces HL-60/ADR cell apoptosis via an upregulation of Bax expression, a downregulation of Bcl-2 expression and an increase in both caspase-3 and caspase-9 activity.

There were several limitations in this study. Firstly, this study was an extension of our previous work (14). We did

not obtain additional funds for a detailed exploration of the apoptotic mechanism in this study; no western blot analysis of protein concentrations was carried out. In addition, we likewise did not study a number of other important anti- and pro-apoptotic proteins due to lack of funds. Funding was insufficient to support animal experimentation and *in vivo* validation. The aforementioned limitations would certainly be eliminated if sufficient funding was obtained by our group in the future.

Conclusions

The present study demonstrated that GSPE induces apoptosis of HL-60/ADR cells via the Bcl-2/Bax caspase-3/9 pathway. GSPE can not only be used as an adjunct in the setting of combination chemotherapy to reduce drug resistance, but also to directly inhibit tumor cell proliferation. A further toxicity study of GSPE in non-tumoral proliferating cells is thus warranted to evaluate its potential uses in clinic.

Acknowledgments

We thank Professor Qi-Hua Fu and Professor Qiu-Hui Pan from the National Children's Medical Center (Shanghai) for providing us an experimental site as well as technical guidance. We would like to thank Dr. Erik Matro for his help in polishing our paper.

Funding: This work was supported by the Pudong Shanghai Science and Technology Development Fund (PKJ2016-Y38) and Shanghai "Rising Stars of Medical Talent" Youth Development Program -Youth Medical Talents – Clinical Pharmacist Program (SHWRS (2020)_087).

Footnote

Reporting Checklist: The authors have completed the MDAR checklist. Available at <https://dx.doi.org/10.21037/tcr-21-920>

Data Sharing Statement: Available at <https://dx.doi.org/10.21037/tcr-21-920>

Peer Review File: Available at <https://dx.doi.org/10.21037/tcr-21-920>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://dx.doi.org/10.21037/tcr-21-920>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013). Institutional ethical approval and informed consent were waived.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- Short NJ, Rytting ME, Cortes JE. Acute myeloid leukaemia. *Lancet* 2018;392:593-606.
- Döhner H, Weisdorf DJ, Bloomfield CD. Acute Myeloid Leukemia. *N Engl J Med* 2015;373:1136-52.
- Zjablovskaja P, Florian MC. Acute Myeloid Leukemia: Aging and Epigenetics. *Cancers (Basel)* 2019;12:103.
- Zwaan CM, Kolb EA, Reinhardt D, et al. Collaborative Efforts Driving Progress in Pediatric Acute Myeloid Leukemia. *J Clin Oncol* 2015;33:2949-62.
- Sekeres MA, Guyatt G, Abel G, et al. American Society of Hematology 2020 guidelines for treating newly diagnosed acute myeloid leukemia in older adults. *Blood Adv* 2020;4:3528-49.
- Döhner H, Estey E, Grimwade D, et al. Diagnosis and management of AML in adults: 2017 ELN recommendations from an international expert panel. *Blood* 2017;129:424-47.
- Siveen KS, Uddin S, Mohammad RM. Targeting acute myeloid leukemia stem cell signaling by natural products. *Mol Cancer* 2017;16:13.
- González-Quilen C, Grau-Bové C, Jorba-Martín R, et al. Protective properties of grape-seed proanthocyanidins in human ex vivo acute colonic dysfunction induced by dextran sodium sulfate. *Eur J Nutr* 2021;60:79-88.
- Margalef M, Pons Z, Iglesias-Carres L, et al. Gender-related similarities and differences in the body distribution of grape seed flavanols in rats. *Mol Nutr Food Res* 2016;60:760-72.
- Xu M, Chen X, Huang Z, et al. Grape seed

- proanthocyanidin extract promotes skeletal muscle fiber type transformation via AMPK signaling pathway. *J Nutr Biochem* 2020;84:108462.
11. Casanova-Martí À, González-Abuín N, Serrano J, et al. Long Term Exposure to a Grape Seed Proanthocyanidin Extract Enhances L-Cell Differentiation in Intestinal Organoids. *Mol Nutr Food Res* 2020;64:e2000303.
 12. Grau-Bové C, González-Quilen C, Terra X, et al. Effects of Flavanols on Enteroendocrine Secretion. *Biomolecules* 2020;10:844.
 13. Liu M, Yun P, Hu Y, et al. Effects of Grape Seed Proanthocyanidin Extract on Obesity. *Obes Facts* 2020;13:279-91.
 14. Lin KN, Jiang YL, Zhang SG, et al. Grape seed proanthocyanidin extract reverses multidrug resistance in HL-60/ADR cells via inhibition of the PI3K/Akt signaling pathway. *Biomed Pharmacother* 2020;125:109885.
 15. Wang K, Wei H, Zhan J, et al. GSPE alleviates renal fibrosis by inhibiting the activation of C3/ HMGB1/ TGF- β 1 pathway. *Chem Biol Interact* 2020;316:108926.
 16. Tu X, Wang M, Liu Y, et al. Pretreatment of Grape Seed Proanthocyanidin Extract Exerts Neuroprotective Effect in Murine Model of Neonatal Hypoxic-ischemic Brain Injury by Its Antiapoptotic Property. *Cell Mol Neurobiol* 2019;39:953-61.
 17. Odai T, Terauchi M, Kato K, et al. Effects of Grape Seed Proanthocyanidin Extract on Vascular Endothelial Function in Participants with Prehypertension: A Randomized, Double-Blind, Placebo-Controlled Study. *Nutrients* 2019;11:2844.
 18. Guo F, Hu Y, Niu Q, et al. Grape Seed Proanthocyanidin Extract Inhibits Human Esophageal Squamous Cancerous Cell Line ECA109 via the NF- κ B Signaling Pathway. *Mediators Inflamm* 2018;2018:3403972.
 19. Sherif AA, Abdelhalim SZ, Salim EI. Immunohistochemical and biochemical alterations following administration of proanthocyanidin extract in rats hepatocellular carcinoma. *Biomed Pharmacother* 2017;93:1310-9.
 20. Song X, Siriwardhana N, Rathore K, et al. Grape seed proanthocyanidin suppression of breast cell carcinogenesis induced by chronic exposure to combined 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and benzoapyrene. *Mol Carcinog* 2010;49:450-63.
 21. Engelbrecht AM, Mattheyse M, Ellis B, et al. Proanthocyanidin from grape seeds inactivates the PI3-kinase/PKB pathway and induces apoptosis in a colon cancer cell line. *Cancer Lett* 2007;258:144-53.
 22. Hu H, Qin YM. Grape seed proanthocyanidin extract induced mitochondria-associated apoptosis in human acute myeloid leukaemia 14.3D10 cells. *Chin Med J (Engl)* 2006;119:417-21.
 23. Liu J, Zhang WY, Kong ZH, et al. Induction of cell cycle arrest and apoptosis by grape seed procyanidin extract in human bladder cancer BIU87 cells. *Eur Rev Med Pharmacol Sci* 2016;20:3282-91.
 24. Ye X, Krohn RL, Liu W, et al. The cytotoxic effects of a novel IH636 grape seed proanthocyanidin extract on cultured human cancer cells. *Mol Cell Biochem* 1999;196:99-108.
 25. Prasain JK, Peng N, Dai Y, et al. Liquid chromatography tandem mass spectrometry identification of proanthocyanidins in rat plasma after oral administration of grape seed extract. *Phytomedicine* 2009;16:233-43.
 26. Monagas M, Quintanilla-López JE, Gómez-Cordovés C, et al. MALDI-TOF MS analysis of plant proanthocyanidins. *J Pharm Biomed Anal* 2010;51:358-72.
 27. Creagh EM, Conroy H, Martin SJ. Caspase-activation pathways in apoptosis and immunity. *Immunol Rev* 2003;193:10-21.
 28. van Delft MF, Huang DC. How the Bcl-2 family of proteins interact to regulate apoptosis. *Cell Res* 2006;16:203-13.
 29. Choi JH, Bogenberger JM, Tibes R. Targeting Apoptosis in Acute Myeloid Leukemia: Current Status and Future Directions of BCL-2 Inhibition with Venetoclax and Beyond. *Target Oncol* 2020;15:147-62.
 30. Cedó L, Castell-Auví A, Pallarès V, et al. Gallic acid is an active component for the anticarcinogenic action of grape seed procyanidins in pancreatic cancer cells. *Nutr Cancer* 2014;66:88-96.
 31. Gao N, Budhraja A, Cheng S, et al. Induction of apoptosis in human leukemia cells by grape seed extract occurs via activation of c-Jun NH2-terminal kinase. *Clin Cancer Res* 2009;15:140-9.
 32. Ruan Y, Jin Q, Zeng J, et al. Grape Seed Proanthocyanidin Extract Ameliorates Cardiac Remodelling After Myocardial Infarction Through PI3K/AKT Pathway in Mice. *Front Pharmacol* 2020;11:585984.
 33. Kwak SC, Cheon YH, Lee CH, et al. Grape Seed Proanthocyanidin Extract Prevents Bone Loss via Regulation of Osteoclast Differentiation, Apoptosis, and Proliferation. *Nutrients* 2020;12:3164.

Cite this article as: Lin KN, Zhao W, Huang SY, Li H. Grape seed proanthocyanidin extract induces apoptosis of HL-60/ADR cells via the Bax/Bcl-2 caspase-3/9 signaling pathway. *Transl Cancer Res* 2021;10(9):3939-3947. doi: 10.21037/tcr-21-920