STROBE Statement—checklist of items that should be included in reports of observational studies

Section/item	Item No	Recommendation	Reported on Page Number/Line Number	Reported on Section/Paragraph
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	Page2/Line24	Abstract/Para2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	Page2-3/Line22-50	Abstract/Para1-4
Introduction				
Background/ rationale	2	Explain the scientific background and rationale for the investigation being reported	Page4/Line57-77	Introduction/ Para1- 2
Objectives	3	State specific objectives, including any prespecified hypotheses	Page5/Line78-81	Introduction/ Para3
Methods				
Study design	4	Present key elements of study design early in the paper	Page5-6/Line87-89 and 105-110	Methods/ Para1-2
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Page5-6/Line87-105	Methods/Par a1-2
Participants	6	(a) Cohort study —Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up Case-control study —Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls Cross-sectional study —Give the eligibility criteria, and the sources and methods of selection of participants	Page5/Line89-96	Methods/Par a1
		(b) Cohort study —For matched studies, give matching criteria and number of exposed and unexposed Case-control study —For matched studies, give matching criteria and the number of controls per case	NA	NA
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	Page5-6/Line97-110	Methods/Par a2
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	Page5-6/Line97-110	Methods/Par a2
Bias	9	Describe any efforts to address potential sources of bias	NA	NA
Study size	10	Explain how the study size was arrived at	NA	NA

Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Page6/Line113-118	Methods/Par a3
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	Page6-7/Line113- 128	Methods/Para3
		(b) Describe any methods used to examine subgroups and interactions	Page7/Line127-128	Methods/Para3
		(c) Explain how missing data were addressed	NA	NA
		(d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed <i>Case-control study</i> —If applicable, explain how matching of cases and controls was addressed <i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy	NA	NA
		(e) Describe any sensitivity analyses	NA	NA
Results				
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	Page7/Line132-136	Results/Para1
		(b) Give reasons for non-participation at each stage	Page7/Line132-136	Results/Para1
		(c) Consider use of a flow diagram	Page7/Line136	Results/Para1
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	Page7/Line136-143	Results/Para1
		(b) Indicate number of participants with missing data for each variable of interest	Page7/Line132-136	Results/Para1
		(c) Cohort study —Summarise follow-up time (eg, average and total amount)	Page7/Line140-141	Results/Para1
Outcome data	15*	Cohort study—Report numbers of outcome events or summary measures over time	Page7/Line141-143	Results/Para1
		Case-control study—Report numbers in each exposure category, or summary measures of exposure	NA	NA
		Cross-sectional study—Report numbers of outcome events or summary measures	NA	NA
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Page8-9/Line145- 186	Results/Para2-4
		(b) Report category boundaries when continuous variables were categorized	Page8-9/Line158- 171	Results/Para3
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	NA	NA
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	Page9-10/Line186- 192	Results/Para4-5
	•			

Discussion				
Key results	18	Summarise key results with reference to study objectives	Page10/Line195- 201	Discussion/Para1
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	Page12/Line234- 245	Discussion/Para5
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	Page10-12/Line202- 233	Discussion/Para2-6
Generalisability	21	Discuss the generalisability (external validity) of the study results	NA	NA
Other information				
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	Page13/Line262	Acknowledgments/P ara1

^{*}Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at http://www.plosmedicine.org/, Annals of Internal Medicine at http://www.annals.org/, and Epidemiology at http://www.epidem.com/). Information on the STROBE Initiative is available at www.strobe-statement.org.

Article Information: http://dx.doi.org/10.21037/tcr-21-2114

*As the checklist was provided upon initial submission, the page number/line number reported may be changed due to copyediting and may not be referable in the published version. In this case, the section/paragraph may be used as an alternative reference.