



# Notes on the morphological features of cotyledonary dissecting leiomyoma, which is rare in clinical practice

Saya Tamura<sup>1#</sup>, Takuma Hayashi<sup>2,3^</sup>, Nobuo Yaegashi<sup>3,4</sup>, Kaoru Abiko<sup>1</sup>, Ikuo Konishi<sup>1,3</sup>

<sup>1</sup>Department of Obstetrics and Gynecology, National Hospital Organization Kyoto Medical Centre, Kyoto, Japan; <sup>2</sup>Cancer Medicine, National Hospital Organization Kyoto Medical Centre, Kyoto, Japan; <sup>3</sup>Medical R&D Promotion Project, The Japan Agency for Medical Research and Development (AMED), Tokyo, Japan; <sup>4</sup>Department of Obstetrics and Gynecology, Tohoku University Graduate School of Medicine, Miyagi, Japan

*Correspondence to:* Takuma Hayashi. National Hospital Organization, Kyoto Medical Centre, Cancer precision medicine at Kyoto University Hospital, Mukaihatake-cho, Fushimi-ku, Kyoto, Japan. Email: yoyoyo224@hotmail.com.

*Comment on:* Ye H, Qi X, Tian Y, *et al.* Case report: cotyledonoid dissecting leiomyoma in a 49-year-old woman. *Transl Cancer Res* 2022;11:4189-93.

**Keywords:** Cotyledonoid dissecting leiomyoma; uterine mesenchymal tumor; leiomyoma; leiomyosarcoma

Submitted Nov 13, 2022. Accepted for publication Feb 13, 2023. Published online Feb 20, 2023.

doi: 10.21037/tcr-22-2605

**View this article at:** <https://dx.doi.org/10.21037/tcr-22-2605>

Ye *et al.* reported the case of a 49-year-old woman with cotyledonoid dissecting leiomyoma diagnosed by surgical pathology using a paraffin-embedded postoperative tissue section (1). In routine gynecologic practice, uterine smooth muscle tumors are among the most common gynecologic tumors. Uterine leiomyoma is a benign smooth muscle tumor that arises from the myometrium and accounts for approximately 75% of all uterine tissues removed from patients (2). Uterine smooth muscle tumors occur mainly in women of reproductive age, and the prevalence of uterine leiomyoma in women up to the age of 50 years is approximately 70% (2,3). However, its incidence is low in postmenopausal women. Uterine leiomyoma proliferation affects the secretion of female hormones, and the size of uterine leiomyomas varies slightly depending on the sexual cycle. Uterine leiomyoma has a histomorphology similar to that of lipoleiomyoma, intravenous leiomyoma, and leiomyoma with bizarre nuclei (4-7). A cotyledonoid dissecting leiomyoma is a uterine leiomyoma with a very rare placental lobed tissue morphology (2) that can be misdiagnosed as a malignant uterine leiomyosarcoma because of its rarity and characteristic appearance on gross examination.

In one such case, contrast-enhanced computed


tomography (CT) revealed a mass that was continuous with the muscular layer of the uterine corpus, suggesting that it arose from there (8). However, there was no infiltration of tumor cells into the vein and myometrium. Therefore, a benign leiomyoma growing outside the uterus and exhibiting a morphology similar to that of the placental leaf is called a cotyledonoid dissecting leiomyoma (8).

A uterine leiomyoma, which extends into the uterine smooth muscle layer and broad ligament as well as extrauterinally in a beaded manner, is accompanied by marked edematous changes and macroscopically resembles a placental lobe cotyledon. Such uterine leiomyomas are, therefore, called cotyledonoid dissecting leiomyomas (4,8,9).

Ye *et al.* reported the case of a 49-year-old woman with cotyledonoid dissecting leiomyoma diagnosed by surgical pathology using a postoperative paraffin-embedded tissue section (1). The patient presented with a history of progressive constipation that lasted for 6 months and a palpable left lower abdominal mass since 1 month (1). She exhibited neither pelvic nor para-aortic lymph node enlargement. Serum tumor markers, namely, cancer antigen (CA) 125, CA19-9, carcinoembryonic antigen (CEA), and alpha-fetoprotein (AFP) levels, were normal. Transvaginal ultrasonography revealed a massive mass comprising two

<sup>#</sup> Current position: Department of Obstetrics and Gynecology, Kyoto University School of Medicine, Kyoto, Japan.

<sup>^</sup> ORCID: 0000-0002-7525-2048.



**Figure 1** Morphological characteristics of a cotyledonoid dissecting leiomyoma observed on MRI and histopathological examination. (A) Results of MRI. The T2W1 image shows a continuous, low-to-high-intensity mass lesion originating within the right muscle layer of the uterine corpus, extending outward into the uterus. The mass (encircled by a dotted line) is present between the round ligament (white arrow) and uterine artery and is enclosed by a membrane. The right ovary (white arrow) present superior to the mass is normal. The images suggest that the mass may be a degenerative uterine leiomyoma growing from within the myometrium into the broad ligament. The leiomyoma measures approximately 20 mm in size on the posterior wall of the uterine fundus. No lesions are seen in the ovaries. Significant lymphadenopathy or ascites is absent. (B) Results of histopathological examination. A soft mass approximately 120 mm in size is seen protruding from the uterine serosal surface. The cut surface of the mass is grayish white and multinodular and demonstrates a proliferation of smooth muscle cells having an island-like/alveolar-like morphology with edematous stroma. Marked infiltration of these smooth muscle cells into the uterine smooth muscle layer is observed. There is also evidence of infiltration of some proliferating smooth muscle cells into the blood vessels. No severe nuclear atypia or mitotic cells are observed. On the basis of these findings, a diagnosis of cotyledonoid dissecting leiomyoma can be considered. Surgical pathological examination reveals no malignant findings in cervical and fallopian tube tissues. Upper panel: magnification  $\times 10$ ; lower panel: magnification  $\times 40$ . (C) Illustration of the gross findings of the patient's cotyledonoid dissecting leiomyoma. MRI, magnetic resonance imaging; H&E, hematoxylin and eosin; CDL, cotyledonoid dissecting leiomyoma.

subserous fibroids measuring 9.9 cm  $\times$  6.9 cm  $\times$  6.3 cm and 8.1 cm  $\times$  6.6 cm  $\times$  6.8 cm with peripheral and internal probing blood flow signals (1). Rapid examination using intraoperative frozen sections revealed an angioleiomyoma with edema (1). Truncal CT performed 6 months after surgery revealed no abnormal findings.

Recently, we encountered a case of cotyledonoid dissecting leiomyoma. In November 2020, a 57-year-old woman visited a hospital because of abnormal vaginal bleeding, and the development of a chocolate cyst was suspected. Magnetic resonance imaging (MRI) revealed a mass in the patient's pelvis (*Figure 1*) that could not be adjudged as malignant or benign. However, ovarian cancer was suspected because a solid component was identified

on the MR image. Therefore, the patient was referred to our hospital that has a gynecological team for thorough examination. Transvaginal ultrasonography revealed a solid mass measuring 115 mm  $\times$  57 mm with an indistinct margin in the right ovary. The area of origin of this mass suggested ovarian cancer or a retroperitoneal tumor. In March 2021, the patient underwent a simple hysterectomy and bilateral salpingectomy. A degenerative uterine leiomyoma measuring 110 mm  $\times$  80 mm was found growing within the broad ligament attached to the right round ligament. No gross abnormalities were observed in the bilateral fallopian tubes and ovaries. The surgical pathological examination of the resected tissue demonstrated a cotyledonoid dissecting leiomyoma. There was no evidence of malignancy in the

endometrial tissue, cervical tissue, or bilateral oviduct tissue. The patient is currently being followed up on an outpatient basis. Similar to the detection of a suspicious malignant mass during MRI by our medical staff, other healthcare professionals must understand the characteristic appearance of a cotyledonoid dissecting leiomyoma. Informed consent was provided by the patient.

A cotyledonoid dissecting leiomyoma is an extremely rare benign uterine tumor. Its appearance on gross examination during surgical treatment is often misinterpreted as malignancy, which may result in overtreatment. Serum tumor markers such as CA125 and CA19-9 levels are elevated in many cases of gynecologic malignancies (10,11). On the contrary, uterine leiomyomas, which are benign tumors, demonstrate normal results for serum tumor marker levels. Unnecessary surgical procedures, i.e., total abdominal hysterectomy (TAH) and bilateral salpingo-oophorectomy (BSO), can be avoided in patients of reproductive age if medical staff are aware of and experienced in recognizing the appearance and characteristics of cotyledonoid dissecting leiomyomas.

All cotyledonoid dissecting leiomyomas reported to date were clinically diagnosed as benign tumors, and only one recurred after initial partial resection (12). This was observed in a 33-year-old woman who underwent an incomplete resection to preserve fertility, and recurrence was reported 5 years after the initial surgical intervention (12). However, tumor recurrence has not been reported in cases where complete resection (hysterectomy) for cotyledonoid dissecting leiomyoma was performed. In the case reported by Ye *et al.*, no evidence of recurrence was observed in the patient 18 months after hysterectomy. Recently, cases of epithelioid cotyledonoid dissociative leiomyoma variants and cotyledonoid dissecting leiomyomas with intravenous leiomyomatosis were also reported (13,14). Histologically, the neoplastic smooth muscle cells of cotyledonoid dissecting leiomyomas form disorganized bundles, contrary to the organized pattern observed in common uterine leiomyomas (15).

In many cases, differentiating between uterine leiomyoma and uterine leiomyosarcoma is difficult. Uterine leiomyoma is the most frequently occurring uterine sarcoma and accounts for 1–2% of all uterine malignant tumors (8). Most uterine leiomyosarcomas occur in women aged >40 years. Unlike uterine benign smooth muscle tumors, uterine leiomyosarcomas are frequently observed in postmenopausal women (16). In uterine mesenchymal tumors that develop from the uterine smooth muscle layer,

the rapid growth of the tumor does not immediately suggest the development of uterine leiomyosarcoma. However, in postmenopausal women who are not undergoing hormone replacement therapy, malignancy should be suspected by medical staff when uterine mesenchymal tumors are growing or recurring. However, in the case reported by Ye *et al.*, neither an epithelioid pattern nor an intravascular neoplastic component was observed (1). Therefore, recurrence after surgical treatment will presumably not occur in this patient.

In conclusion, a cotyledonoid dissecting leiomyoma is a rare variant of uterine leiomyoma (15,17). Its gross appearance and ultrasonographic features may indicate malignancy. Furthermore, it demonstrates >10 increases in mitotic activity/10 high-power fields, tumor cell necrosis, and no evidence of cellular atypia. Thus, histologically, cotyledonoid dissecting leiomyoma is a benign tumor. To date, recurrence and metastasis have not been reported in a majority of the cases. Therefore, in clinical practice, prognosis is considered favorable. Clinicians and pathologists must understand the oncologic features of cotyledonoid dissecting leiomyoma to prevent misdiagnosis of malignancy and consequent overtreatment.

### Acknowledgments

We appreciate Dr. Susumu Tonegawa (M.I.T., Cambridge, MA, USA) and Dr. Zhang W (Roche Tissue Diagnostics, Tucson, AZ, USA) for critical research assistance. We thank all medical staff for clinical research at Kyoto University School of Medicine and the National Hospital Organization Kyoto Medical Center. We appreciate Crimson Interactive Japan Co., Ltd., for revising and polishing our manuscript.

*Funding:* The clinical research of the case reported by the authors was performed with research funding from the following: Japan Society for Promoting Science for TH (Grant No. 19K09840), START-program Japan Science and Technology Agency for TH (Grant No. STSC20001), and the National Hospital Organization Multicenter clinical study for TH (Grant No. 2019-Cancer in general-02), and The Japan Agency for Medical Research and Development (AMED) (Grant No. 22ym0126802j0001), Tokyo, Japan.

### Footnote

*Provenance and Peer Review:* This article was commissioned by the editorial office, *Translational Cancer Research*. The article did not undergo external peer review.

*Conflicts of Interest:* All authors have completed the ICMJE uniform disclosure form (available at <https://tcr.amegroups.com/article/view/10.21037/tcr-22-2605/coif>). The authors have no conflicts of interest to declare.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. Informed consent was provided by the patient whom the authors reported in this study.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

- Ye H, Qi X, Tian Y, et al. Case report: cotyledonoid dissecting leiomyoma in a 49-year-old woman. *Transl Cancer Res* 2022;11:4189-93.
- Uterine leiomyoma. *Female Genital Tumours WHO Classification of Tumours*, 5th ed., Vol.4. WHO Classification of Tumours Editorial Board. World Health Organization. 2020:272-6.
- WHO classification of mesenchymal tumours of the lower genital tract. *Female Genital Tumours WHO Classification of Tumours*, 5th ed., Vol.4. WHO Classification of Tumours Editorial Board. World Health Organization. 2020:13.
- Zaloudek CJ, Hendrickson MR, Soslow RA. Mesenchymal tumors of the uterus. In Kurman R, editor. *Blaustein's Pathology of the Female Genital Tract*, 6th ed. Springer, New York, 2010:453-527.
- Intravenous leiomyomatosis. *Female Genital Tumours WHO Classification of Tumours*, 5th ed., Vol.4. WHO Classification of Tumours Editorial Board. World Health Organization. 2020:277-8.
- Watanabe K, Hayashi T, Katsumata M, et al. Development of Uterine Leiomyosarcoma During Follow-up After Caesarean Section in a Woman With Uterine Leiomyoma. *Anticancer Res* 2021;41:3001-10.
- Nishikawa S, Hayashi T, Amano Y, et al. Characteristic of Concurrent Uterine Lipoleiomyoma and Hemangioma by Algorithm of Candidate Biomarkers for Uterine Mesenchymal Tumor. *Diagnostics (Basel)* 2022;12:2468.
- Kuman RJ, Carcangiu MI, Herrington CS, et al. editors. *WHO Classification of Tumours of Female Reproductive Organs*, 4th ed, IARC Press, Lyon, 2014.
- Ip PP, Tse KY, Tam KF. Uterine smooth muscle tumors other than the ordinary leiomyomas and leiomyosarcomas: a review of selected variants with emphasis on recent advances and unusual morphology that may cause concern for malignancy. *Adv Anat Pathol* 2010;17:91-112.
- Dolscheid-Pommerich RC, Keyver-Paik M, Hecking T, et al. Clinical performance of LOCI™-based tumor marker assays for tumor markers CA 15-3, CA 125, CEA, CA 19-9 and AFP in gynecological cancers. *Tumour Biol* 2017;39:1010428317730246.
- Babacan A, Kizilaslan C, Gun I, et al. CA 125 and other tumor markers in uterine leiomyomas and their association with lesion characteristics. *Int J Clin Exp Med* 2014;7:1078-83.
- Roth LM, Kirker JA, Insull M, et al. Recurrent cotyledonoid dissecting leiomyoma of the uterus. *Int J Gynecol Pathol* 2013;32:215-20.
- Soleymani Majd H, Ismail L, Desai SA, et al. Epithelioid cotyledonoid dissecting leiomyoma: a case report and review of the literature. *Arch Gynecol Obstet* 2011;283:771-4.
- Jordan LB, Al-Nafussi A, Beattie G. Cotyledonoid hydropic intravenous leiomyomatosis: a new variant leiomyoma. *Histopathology* 2002;40:245-52.
- Weissferdt A, Maheshwari MB, Downey GP, et al. Cotyledonoid dissecting leiomyoma of the uterus: a case report. *Diagn Pathol* 2007;2:18.
- Toledo G, Oliva E. Smooth muscle tumors of the uterus: a practical approach. *Arch Pathol Lab Med* 2008;132:595-605.
- Jamal I, Gupta RK, Sinha RK, et al. Cotyledonoid dissecting leiomyoma: an uncommon form of a common disease. *Obstet Gynecol Sci* 2019;62:362-6.

**Cite this article as:** Tamura S, Hayashi T, Yaegashi N, Abiko K, Konishi I. Notes on the morphological features of cotyledonary dissecting leiomyoma, which is rare in clinical practice. *Transl Cancer Res* 2023;12(3):452-455. doi: 10.21037/tcr-22-2605