



# The efficacy of anlotinib as third-line treatment for non-small cell lung cancer by EGFR mutation status: a subgroup analysis of the ALTER0303 randomized phase 3 study

Yizhuo Zhao<sup>1</sup>, Qiming Wang<sup>2,3</sup>, Li Zhang<sup>4</sup>, Jianhua Shi<sup>5</sup>, Zhehai Wang<sup>6</sup>, Ying Cheng<sup>7</sup>, Jianxing He<sup>8</sup>, Yuankai Shi<sup>9</sup>, Weiqiang Chen<sup>10</sup>, Yi Luo<sup>11</sup>, Lin Wu<sup>11</sup>, Xiuwen Wang<sup>12</sup>, Kejun Nan<sup>13</sup>, Faguang Jin<sup>14</sup>, Jian Dong<sup>15</sup>, Baolan Li<sup>16</sup>, Fumihiko Yamaguchi<sup>17</sup>, Daniel Breadner<sup>18,19</sup>, Tatsuya Nagano<sup>20</sup>, Fumihiko Tanaka<sup>21</sup>, Hatim Husain<sup>22</sup>, Kai Li<sup>23</sup>, Baohui Han<sup>1</sup>

<sup>1</sup>Department of Respiratory Medicine, Shanghai Chest Hospital, Shanghai Jiao Tong University, Shanghai, China; <sup>2</sup>Department of Internal Medicine, Affiliated Cancer Hospital of Zhengzhou University, Zhengzhou, China; <sup>3</sup>Department of Internal Medicine, Henan Cancer Hospital, Zhengzhou, China; <sup>4</sup>Department of Respiratory Diseases, Peking Union Medical College Hospital, Beijing, China; <sup>5</sup>Department of Medical Oncology, Linyi Cancer Hospital, Linyi, China; <sup>6</sup>Department of Internal Medicine-Oncology, Shandong Cancer Hospital, Jinan, China; <sup>7</sup>Department of Thoracic Medical Oncology, Jilin Cancer Hospital, Changchun, China; <sup>8</sup>Department of Thoracic Surgery, The First Affiliated Hospital of Guangzhou Medical University, Guangzhou, China; <sup>9</sup>Department of Medical Oncology, Cancer Hospital Chinese Academy of Medical Sciences, Beijing, China; <sup>10</sup>Department of Pulmonary Medicine, Lanzhou Military General Hospital, Lanzhou, China; <sup>11</sup>Department of Medical Oncology, Hunan Cancer Hospital, Changsha, China; <sup>12</sup>Department of Chemotherapy, Qilu Hospital of Shandong University, Jinan, China; <sup>13</sup>Department of Oncology, The First Affiliated Hospital of Xi'an Jiaotong University, Xi'an, China; <sup>14</sup>Department of Respiratory Diseases, Tang Du Hospital, Xi'an, China; <sup>15</sup>First Department of Medical Oncology, Yunnan Cancer Hospital, Kunming, China; <sup>16</sup>Department of General Medicine, Capital Medical University, Beijing Chest Hospital, Beijing, China; <sup>17</sup>Department of Respiratory Medicine, Showa University Fujigaoka Hospital, Yokohama, Japan; <sup>18</sup>Division of Medical Oncology, London Regional Cancer Program at London Health Science Center, London, Canada; <sup>19</sup>Schulich School of Medicine and Dentistry at Western University, London, Canada; <sup>20</sup>Division of Respiratory Medicine, Department of Internal Medicine, Kobe University Graduate School of Medicine, Kobe, Japan; <sup>21</sup>Second Department of Surgery, University of Occupational and Environmental Health, Kitakyushu, Japan; <sup>22</sup>Division of Hematology and Oncology, Moores Cancer Center, University of California, San Diego, La Jolla, CA, USA; <sup>23</sup>Department of Thoracic Oncology, Tianjin Medical University Cancer Institute and Hospital, Tianjin, China

**Contributions:** (I) Conception and design: B Han, K Li, Q Wang, L Zhang, Z Wang, Y Cheng, J He, Y Zhao, W Chen, L Wu, X Wang, K Nan, F Jin, J Dong, B Li; (II) Administrative support: B Han, K Li, Y Zhao, Q Wang, L Zhang, J Shi, Z Wang, Y Cheng, J He, Y Shi, W Chen, Y Luo, L Wu, X Wang, K Nan, F Jin, J Dong, B Li; (III) Provision of study materials or patients: B Han, K Li, Y Zhao, Q Wang, L Zhang, J Shi, Z Wang, Y Cheng, J He, Y Shi, W Chen, Y Luo, L Wu, X Wang, K Nan, F Jin, J Dong, B Li; (IV) Collection and assembly of data: B Han, K Li, Y Zhao, Q Wang, L Zhang, J Shi, Z Wang, Y Cheng, J He, Y Shi, W Chen, Y Luo, L Wu, X Wang, K Nan, F Jin, J Dong, B Li; (V) Data analysis and interpretation: Y Zhao, K Li, B Han; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

**Correspondence to:** Dr. Baohui Han. Shanghai Chest Hospital, West Huaihai Road, No. 241 Xuhui District, Shanghai 200030, China.

Email: 18930858216@163.com; Dr. Kai Li. Department of Thoracic Oncology, Tianjin Medical University Cancer Institute and Hospital, West Huan-Hu Rd, Ti Yuan Bei, Hexi District, Tianjin 300060, China. Email: likai\_fnk@163.com.

**Background:** Anlotinib demonstrated improved overall survival (OS) and progression-free survival (PFS) compared with placebo as a third-line or subsequent therapy in patients with non-small cell lung cancer (NSCLC) in the ALTER0303 trial. The status of epidermal growth factor receptor (EGFR) mutation, different previous treatment may affect the efficacy of subsequent therapy, and we did this subgroup analysis to characterize the efficacy of anlotinib in patients with and without EGFR mutation.

**Methods:** The ALTER0303 trial was a randomized, double-blind, phase 3 study of anlotinib in patients with NSCLC who failed at least 2 lines of treatment. In the study, 138 of 437 randomized patients were EGFR mutation positive. A Cox model was used to examine the influence of previous treatment on the efficacy of anlotinib according to EGFR mutation status.

**Results:** For patients with EGFR mutation, the OS was 10.7 and 6.3 months (HR 0.59; 95% CI: 0.38–0.94, P=0.025) in the anlotinib and placebo group, respectively. The PFS was 5.6 and 0.8 months (HR 0.21; 95%

CI: 0.13–0.32,  $P < 0.0001$ ) in the anlotinib and placebo group, respectively. For patients without EGFR mutation, the OS was 8.9 months for anlotinib and 6.5 months for placebo (HR 0.73; 95% CI: 0.55–0.97,  $P = 0.029$ ), and the PFS was 5.4 months for anlotinib and 1.6 months for placebo (HR 0.29; 95% CI: 0.22–0.39,  $P < 0.0001$ ). In the anlotinib group, the OS and PFS for patients with and without EGFR mutation was 10.7 and 8.9 months (HR 0.69; 95% CI: 0.50–0.95,  $P = 0.021$ ), 5.6 and 5.4 months (HR 1.00; 95% CI: 0.75–1.34,  $P = 1.000$ ), respectively. The incidence of adverse events was similar in subgroups.

**Conclusions:** This analysis demonstrated that the benefit of anlotinib as a third-line therapy for patients with NSCLC was independent of EGFR mutation status.

**Keywords:** Anlotinib; non-small cell lung cancer (NSCLC); third-line; EGFR mutation

Submitted Feb 18, 2022. Accepted for publication May 20, 2022.

doi: 10.21037/tlcr-22-320

View this article at: <https://dx.doi.org/10.21037/tlcr-22-320>

## Introduction

Over the past decade, great progress has been achieved in the treatment of advanced non-small cell lung cancer (NSCLC). Targeted therapy determined by individual molecular profiles, including immunotherapy and anti-angiogenesis therapy, has improved survival. However, most patients will develop progressive disease after response to first- and second-line therapy, while the subsequent treatment is less hopeful and a clear standard has not been established, and there is an unmet need to improve treatment for this population.

Angiogenesis is a complex process that plays an important role in sustaining the tumor microenvironment, tumor growth, and metastatic dissemination, involving both pro-angiogenic and anti-angiogenic molecules. The vascular endothelial growth factor (VEGF) family members play a key role in angiogenesis, among which VEGFA is the main mediator of tumor-associated angiogenesis. Of the VEGF receptors that are crucial for the transmission of angiogenic signals, VEGFR-2 is the major mediator of the mitogenic, angiogenic, and permeability effects of VEGF (1). Several other pro-angiogenic factors also impact tumor angiogenesis, including platelet-derived growth factor (PDGF) and fibroblast growth factor (FGF). Cancer cells can express and secrete PDGF $\beta$  and FGF-2 at a high level, which assists in stimulating angiogenesis (2).

Treatment by targeting tumor angiogenesis has been approached through several methods, including monoclonal antibodies that block VEGF-VEGFR binding, VEGFR tyrosine kinase inhibitors (TKIs), and VEGF Trap (3). Bevacizumab and ramucirumab, which are monoclonal antibodies targeting VEGF and VEGFR, respectively, have led

to improved overall survival (OS) for NSCLC when added to standard first- and second-line chemotherapy. However, only about one third of patients respond to anti-VEGF antibodies (4,5). Previously, VEGFR TKIs failed to show significant prolongation of OS either in combination with chemotherapy as first- and second-line therapy, or in comparison with placebo as third- and fourth-line therapy, including vandetanib, sorafenib, sunitinib, axitinib, and cediranib (6).

Anlotinib is a novel multi-target TKI that inhibits VEGFR2/3, FGFR1-4, PDGFR  $\alpha/\beta$ , c-Kit, and Ret, among others (7). In a preclinical study, the inhibition of VEGFR2, PDGFR $\beta$ , and FGFR1 by anlotinib was stronger than sunitinib and sorafenib (8). Anlotinib showed higher selectivity for VEGFR family members than other VEGFR TKIs, which may indicate milder toxicity (9). In the ALTER0302 trial, a phase 2 study of anlotinib as a third-line treatment for patients with advanced NSCLC compared with placebo, progression-free survival (PFS) was prolonged significantly in the anlotinib group regardless of EGFR mutation status in the post-hoc subgroup analysis (10). In the ALTER0303 trial, anlotinib significantly improved PFS and OS as a third-line and subsequent treatment for NSCLC patients regardless of EGFR mutation status (11). In China, EGFR mutations are found in up to 45% of NSCLC patients (12). In previous studies, the response rates of chemotherapy in patients with and without EGFR mutation status were different (13), and frontline EGFR TKI may reduce the efficacy of chemotherapy in patients with EGFR mutation (14). The previous treatments of patients with and without EGFR mutation are distinct, and we analyzed the efficacy and safety of anlotinib in these two subpopulations in the ALTER0303 trial. We present the

following article in accordance with the TREND reporting checklist (available at <https://tcr.amegroups.com/article/view/10.21037/tcr-22-320/rc>).

## Methods

### *Study design and procedures*

ALTER0303 was a randomized, double-blind, placebo-controlled phase 3 study (NCT02388919), which has been published (11). The study was conducted in accordance with Declaration of Helsinki (as revised in 2013) and Good Clinical Practice requirements. The study was approved by the institutional ethics board of each site. All the patients provided written informed consent before study entry.

The main inclusion criteria included: aged 18–75 years, histologically or cytologically confirmed NSCLC, at least 2 lines of chemotherapy for patients without driver mutation, and at least 1 line of chemotherapy and 1 line of TKI therapy for patients with known driver mutations. Subjects were randomized in a 2:1 ratio with a block randomization scheme (block size of 4) to receive oral anlotinib 12 mg/d or placebo on day 1–14 of a 21-day cycle. The primary endpoint was OS, the secondary endpoints included progression-free survival (PFS), objective response rate (ORR), disease control rate (DCR) and safety. EGFR mutation status was one of the pre-stratification factors. Patients should have formal report of EGFR and ALK status, those who did not have report should provide specimens for central detection. Tumor assessment was conducted with computed tomography every 2 cycles, follow-up was done every 8 weeks until death or the data cutoff date. The efficacy of treatment was assessed by RECIST v1.1 and treatment-related adverse events were graded according to NCI CTCAE v5.5.

We compared the outcomes according to potential covariates including prior duration of targeted therapy, the duration of EGFR TKI treatment, and the time since the start of prior therapy (TSPT) in patients assessed by EGFR mutation status. We identified 18 months as the cut-off point of TSPT, as it was the median time from the start of first-line treatment to study entry. Eight months was identified as the cut-off point of time to progression with targeted therapy as it was the median treatment duration of EGFR TKI.

### *Statistical analysis*

We used the Kaplan-Meier estimates method to assess the

median PFS and OS with 95% confidence intervals (CIs). Differences in survival were assessed using the log-rank test. The proportional hazards (Cox) model was used to estimate hazard ratios (HRs) for PFS and OS with 95% CIs. To verify the relationship between EGFR mutation status and treatment efficacy, an interaction term of EGFR mutation status and the intervention of anlotinib or placebo was conducted using multiple regression model. All statistical tests were two sided, and  $P < 0.05$  was considered to be statistically significant. All analyses were performed using SAS software, version 9.4.

## Results

### *Characteristics of patients according to EGFR mutation status*

Of the 437 randomized patients, 138 (31.6%) were EGFR mutation positive, including 93 in the anlotinib group and 45 in the placebo group, while 299 (68.4%) were EGFR mutation negative, including 201 in the anlotinib group and 98 in the placebo group. Baseline characteristics were balanced between treatment arms in both the EGFR mutation positive and negative subgroups. In patients with EGFR mutation, the proportion of female and adenocarcinoma was higher, which are known clinical factors associated with higher EGFR mutation frequency (Table 1).

### *The efficacy of anlotinib in patients with and without EGFR mutation*

In the anlotinib group, the PFS for patients with and without EGFR mutation was 5.6 and 5.4 months (HR 1.00; 95% CI: 0.75–1.34,  $P=1.000$ ), the OS was 10.7 and 8.9 months (HR 0.69; 95% CI: 0.50–0.95,  $P=0.021$ ) respectively; while in the placebo group, the PFS for patients with and without EGFR mutation was 0.8 and 1.6 months (HR 1.53; 95% CI: 1.03–2.26,  $P=0.033$ ), the OS was 6.3 and 6.5 months respectively (HR 0.87; 95% CI: 0.57–1.35,  $P=0.541$ ).

In patients with EGFR mutation, the OS was 10.7 and 6.3 months (HR 0.59; 95% CI: 0.38–0.94,  $P=0.025$ ) for anlotinib and placebo, respectively. The PFS was 5.6 and 0.8 months (HR 0.21; 95% CI: 0.13–0.32,  $P < 0.0001$ ) for anlotinib and placebo, respectively. The ORR was 7.5% for anlotinib and 0% for placebo, while the DCR was 82.8% for anlotinib and 20.0% for placebo. In patients without EGFR mutation, the OS was 8.9 and 6.5 months (HR 0.73;

**Table 1** Baseline characteristics of patients with and without EGFR mutation


Characteristics	EGFR M+, n (%)		EGFR M-, n (%)	
	Anlotinib (n=93)	Placebo (n=45)	Anlotinib (n=201)	Placebo (n=98)
Gender				
Male	42 (45.2)	23 (51.1)	146 (72.6)	74 (75.5)
Female	51 (54.8)	22 (48.9)	55 (27.4)	24 (24.5)
Age, years				
≤60	55 (59.1)	31 (68.9)	98 (48.7)	59 (60.2)
61–69	34 (36.6)	11 (24.4)	91 (45.3)	30 (30.6)
≥70	4 (4.3)	3 (6.7)	12 (6.0)	9 (9.2)
Pathological diagnosis				
Adeno-carcinoma	85 (91.4)	43 (95.6)	143 (71.1)	65 (66.3)
Squamous	6 (6.5)	2 (4.4)	47 (23.4)	31 (31.6)
Others	2 (2.2)	0 (0.0)	11 (5.5)	2 (2.1)
Clinical stage				
IIIB	2 (2.1)	3 (6.7)	13 (6.5)	4 (4.1)
IV	91 (97.9)	42 (93.3)	186 (92.5)	94 (95.9)
Others	0 (0.0)	0 (0.0)	2 (1.0)	0 (0.0)
Number of metastases				
>3	44 (47.3)	23 (51.1)	79 (39.3)	39 (39.8)
≤3	49 (52.7)	22 (48.9)	122 (60.7)	59 (60.2)
Lines of prior chemotherapy				
2	55 (59.1)	26 (57.8)	112 (55.7)	52 (53.1)
3	34 (36.6)	19 (42.2)	89 (44.3)	46 (46.9)
1	4 (4.3)	0 (0.0)	0 (0.0)	0 (0.0)
Prior targeted therapy				
No	2 (2.2)	3 (6.7)	134 (66.7)	71 (72.5)
Yes	91 (97.8)	42 (93.3)	67 (33.3)	27 (27.5)
Prior radiotherapy				
No	59 (63.4)	20 (44.4)	117 (58.2)	58 (59.2)
Yes	34 (36.6)	25 (55.6)	84 (41.8)	40 (40.8)
ECOG PS				
0	22 (23.7)	6 (13.3)	37 (18.4)	16 (16.3)
1	71 (76.3)	39 (86.7)	162 (80.6)	81 (82.7)
2	0 (0.0)	0 (0.0)	2 (1.0)	1 (1.0)

**Table 1** (continued)

Table 1 (continued)

Characteristics	EGFR M+, n (%)		EGFR M-, n (%)	
	Anlotinib (n=93)	Placebo (n=45)	Anlotinib (n=201)	Placebo (n=98)
Smoking history				
Once or current smoker	1 (1.1)	2 (4.4)	12 (6.0)	8 (8.2)
None-smoker	92 (98.9)	43 (95.6)	189 (94.0)	90 (91.8)
ALK status				
Negative	93 (100.0)	45 (100.0)	193 (96.0)	95 (97.0)
Positive	0 (0.0)	0 (0.0)	5 (2.5)	2 (2.0)
Unknown	0 (0.0)	0 (0.0)	3 (1.5)	1 (1.0)

EGFR, epidermal growth factor receptor; M+, mutation positive; M-, mutation negative; ECOG PS, eastern cooperative oncology group performance status; ALK, anaplastic lymphoma kinase.



**Figure 1** Kaplan-Meier estimates of overall survival and progression-free survival in patients with and without EGFR mutation. (A) Overall survival of patients with and without EGFR mutation; (B) progression-free survival of patients with and without EGFR mutation. EGFR, epidermal growth factor receptor.

95% CI: 0.55–0.97,  $P=0.029$ ) for anlotinib and placebo, respectively. The PFS was 5.4 and 1.6 months (HR 0.29; 95% CI: 0.22–0.39,  $P<0.0001$ ) for anlotinib and placebo, respectively. In the multiple Cox regression analysis, the interaction term of OS and PFS was 0.1064 and 0.4485 respectively (Tables S1–S4). The ORR was 10.0% for anlotinib and 1.0% for placebo, while the DCR was 80.1% for anlotinib and 44.9% for placebo (Figure 1A, 1B).

#### The effect of previous targeted therapy on survival in patients with EGFR mutation

Among the 138 patients with EGFR mutation, 133 received

EGFR TKIs before study entry (91/93 in the anlotinib group and 42/45 in the placebo group). A total of 27 (20.3%) received EGFR TKIs as first-line therapy, while 106 (79.7%) received EGFR TKIs as second-line therapy. The proportion of patients who received EGFR TKIs as first-line therapy in the anlotinib group was lower (17.6% vs. 26.2%). The most frequently used EGFR TKIs were gefitinib, erlotinib, and icotinib. For patients who received TKIs as first-line therapy (16/93 in the anlotinib group, 11/45 in the placebo group), the PFS was 4.3 months for anlotinib and 0.8 months for placebo (HR 0.32; 95% CI: 0.11–0.91,  $P=0.032$ ), while the OS was 11.1 months for anlotinib and 3.5 months for placebo (HR 0.31; 95%



CI: 0.11–0.87,  $P=0.026$ ; Figure S1A,S1B). For those who received EGFR TKIs as second-line therapy (75/93 in the anlotinib group, 31/45 in the placebo group), the PFS was 5.6 months for anlotinib and 0.9 months for placebo (HR 0.2; 95% CI: 0.12–0.34,  $P<0.0001$ ), while the OS was 10.7 months for anlotinib and 7.0 months for placebo (HR 0.69; 95% CI: 0.4–1.18,  $P=0.17$ ; Figure S1C,S1D).

The median treatment duration of previous targeted therapy was 8 months. Patients who progressed from EGFR TKIs within 8 months (30 in the anlotinib group, 11 in the placebo group) had significantly longer PFS (4.5 *vs.* 0.7 months, HR 0.07; 95% CI: 0.02–0.26,  $P<0.0001$ ) and OS (10.0 *vs.* 3.4 months, HR 0.39; 95% CI: 0.17–0.9,  $P=0.03$ ) with anlotinib compared with placebo (Figure S2A,S2B). For those who progressed from EGFR TKIs exceeding 8 months (61 in the anlotinib group, 31 in the placebo group), the PFS with anlotinib was longer than with placebo (5.6 *vs.* 1.1 months; HR 0.24; 95% CI: 0.14–0.4,  $P<0.0001$ ), while no significant difference in OS was found (12.9 *vs.* 7.0 months, HR 0.62; 95% CI: 0.35–1.09,  $P=0.1$ ; Figure S2C,S2D).

### *The effect of TSPT on survival*

The median time for patients from the start of prior treatment to study entry was 18 months. In patients with EGFR mutation whose TSPT  $\leq 18$  months (28 in the anlotinib group, 14 in the placebo group), the PFS (4.3 *vs.* 0.8 months, HR 0.18; 95% CI: 0.08–0.42,  $P<0.0001$ ) was improved significantly with anlotinib compared with placebo, while the OS (6.6 *vs.* 4.4 months, HR 0.58; 95% CI: 0.27–1.28,  $P=0.18$ ) was improved without a significant difference. In patients without EGFR mutation whose TSPT  $\leq 18$  months (110 in the anlotinib group, 61 in the placebo group), the PFS (4.2 *vs.* 1.4 months, HR 0.29; 95% CI: 0.19–0.42,  $P<0.0001$ ) was improved significantly with anlotinib compared with placebo, while no difference in OS was found (7.3 and 6.3 months, HR 0.77; 95% CI: 0.54–1.1,  $P=0.15$ ; Figure S3A,S3B).

In patients with EGFR mutation whose TSPT  $> 18$  months (65 in the anlotinib group, 31 in the placebo group), the PFS (5.6 *vs.* 0.8 months, HR 0.22; 95% CI: 0.13–0.36,  $P<0.0001$ ) and OS (15.8 *vs.* 7.0 months, HR 0.55; 95% CI: 0.31–0.99,  $P=0.043$ ) were improved with anlotinib. In patients without EGFR mutation whose TSPT  $> 18$  months (91 in the anlotinib group, 37 in the placebo group), the PFS was improved (5.8 *vs.* 2.8 months, HR 0.3; 95% CI: 0.19–0.48,  $P<0.0001$ ) with anlotinib compared with

placebo, while the OS (10.5 *vs.* 8.0 months, HR 0.79; 95% CI: 0.48–1.3,  $P=0.36$ ) was prolonged without significance (Figure S3C,S3D).

### *Safety*

Treatment-related adverse events (TRAEs) were more frequently in patients treated with anlotinib than placebo in both the subgroups, such as hypertension (64.5% *vs.* 11.1% in patients with EGFR mutation, 64.7% *vs.* 15.3% in patients without EGFR mutation), elevated thyroglobulin (52.7% *vs.* 0, 40.8% *vs.* 6.1%), hand and foot syndrome (48.4% *vs.* 8.9%, 40.8% *vs.* 9.2%), elevated triglyceride (41.9% *vs.* 15.6%, 37.8% *vs.* 20.4%), proteinuria (26.9% *vs.* 8.9%, 25.9% *vs.* 15.3%), and rash (7.5% *vs.* 4.4%, 13.4% *vs.* 6.1%). The most common grade 3–4 TRAEs with anlotinib in patients with and without EGFR mutation were hypertension (9.7%, 14.9%) and elevated triglyceride (3.2%, 2.0%), which were manageable. Most of the TRAEs with anlotinib were similar in patients with and without EGFR mutation, while elevated thyroglobulin and hand and foot syndrome occurred more frequently in patients with EGFR mutation (52.7% *vs.* 40.8%). Rash occurred more frequently in patients without EGFR mutation (7.5% *vs.* 13.4%) (Table 2).

### **Discussion**

In these subgroup analyses, anlotinib demonstrated significant OS and PFS benefits in patients with and without EGFR mutation who had failed at least 2 lines of therapy. During the study recruitment, most patients with EGFR mutation still received chemotherapy as first-line therapy, due to the relatively low EGFR mutation detection rate at diagnosis several years ago. Osimertinib was not approved at that time, and patients who progressed from first-line EGFR TKIs usually received chemotherapy as second-line treatment without T790M detection. Only 19.6% of patients with EGFR mutation received EGFR TKIs as first-line therapy in this study. Approximately 60% of patients with EGFR mutation who received first-generation EGFR TKIs will develop a secondary T790M mutation on progression (15), and osimertinib has been approved as a standard second-line therapy for this population (16). For patients without T790M mutation or those with T790M mutation who do not respond to osimertinib, chemotherapy remains the primary choice, but the clinical benefit from chemotherapy may be reduced substantially after EGFR TKI treatment (14). Several anti-PD-1/PD-L1 antibodies

**Table 2** Selected treatment-related adverse events in patients with anlotinib or placebo according to EGFR mutation status

Event	EGFR M+ (n=138) (%)				EGFR M- (n=299) (%)			
	Anlotinib (n=93)		Placebo (n=45)		Anlotinib (n=201)		Placebo (n=98)	
	Any grade	Grade 3–4	Any grade	Grade 3–4	Any grade	Grade 3–4	Any grade	Grade 3–4
Hypertension	60 (64.5)	9 (9.7)	5 (11.1)	0	130 (64.7)	30 (14.9)	15 (15.3)	0
Elevated thyroglobulin	49 (52.7)	0	0	0	82 (40.8)	1 (0.5)	6 (6.1)	0
Hand and foot syndrome	45 (48.4)	2 (2.2)	4 (8.9)	0	82 (40.8)	9 (4.5)	9 (9.2)	0
Elevated triglyceride	39 (41.9)	3 (3.2)	7 (15.6)	0	75 (37.8)	4 (2.0)	20 (20.4)	0
Proteinuria	25 (26.9)	2 (2.2)	4 (8.9)	1 (2.2)	52 (25.9)	5 (2.5)	15 (15.3)	0
Rash	7 (7.5)	0	2 (4.4)	0	27 (13.4)	0	6 (6.1)	1 (1.0)

EGFR, epidermal growth factor receptor; M+, mutation positive; M-, mutation negative.

have been approved for NSCLC as second- and further-line therapies (17), but a low percentage of patients with EGFR mutations benefit from immunotherapy (18). Anti-angiogenic agents, ramucirumab and nintedanib, showed improved survival in combination with docetaxel as a second-line therapy (5,19), but neither is approved in China.

For patients with EGFR mutation who had disease progression after EGFR TKI treatment and chemotherapy, anlotinib achieved a PFS of 5.6 months and an OS of 10.7 months. For patients without EGFR mutation who progressed from 2 lines of chemotherapy, anlotinib achieved a PFS of 5.4 months and an OS of 8.9 months, which was consistent with the results in a previous phase 2 study (10). In previous second-line studies, the PFS was 2.3–3.9 months and the OS was 9.2–13.8 months with immunotherapy, the PFS and OS were 3.4–4.5 and 10.1–10.5 months respectively with the combination of anti-angiogenic agents and docetaxel (5,17,19). This is the first VEGFR TKI that prolonged OS significantly as a single agent in the third-line setting, and the results of OS and PFS were promising and comparable with those in other second-line studies. The stronger effect on VEGF receptors than other VEGF TKIs, as well as the optimal effect on FGFR1–4 and PDGFR  $\alpha/\beta$  which contribute to the resistance of anti-VEGF treatment, may explain the favorable treatment effect of anlotinib (20).

In our study, the prognosis of patients with EGFR mutation might be worse, for the PFS and OS were shorter than those without EGFR mutation in the placebo group. While in the anlotinib group, the OS of patients with EGFR mutation was significantly improve, indicating that patients with EGFR mutation might benefit more from anti-angiogenic therapy than those without EGFR mutation.

It was reported the activation of EGFR can upregulate the expression of VEGF and VEGFR-1, activate VEGFR and promote angiogenesis. The dependence of EGFR mutant tumors on VEGF may explain our results (21).

In this analysis, the benefit of PFS and OS with anlotinib was found to be consistent in both patients with and without EGFR mutation. The pathological type of most patients with EGFR mutation was adenocarcinoma. In the LUME-Lung 1 study, a significant OS benefit (12.6 *vs.* 10.3 months) was demonstrated in the docetaxel plus nintedanib group compared with the docetaxel plus placebo group in adenocarcinoma patients but not in the total study population, indicating the efficacy of anti-angiogenic therapy for non-squamous NSCLC might be better than that of squamous NSCLC (19). In the ZEPHYR study, a greater benefit in PFS was suggested for vandetanib-treated patients with EGFR mutation compared with those without mutation (22). In the MISSION study, OS was similar in the whole population, while among the 89 patients with EGFR mutations, both OS (13.9 *vs.* 6.5 m) and PFS (2.7 *vs.* 1.4 m) were significantly improved with sorafenib compared with placebo (23). These results also suggest that EGFR mutation status may correlate with the treatment effect of VEGFR TKIs. There was previous evidence that mutant EGFR may enhance VEGF expression in lung cancers, suggesting that different angiogenic mechanisms might exist between EGFR mutation positive and negative tumors (24).

The LUME-Lung 1 and REVEL studies demonstrated survival advantages for nintedanib and ramucirumab plus docetaxel in advanced NSCLC patients with TSPT <9 months, suggesting that patients with a poorer prognosis may benefit more from the addition of anti-angiogenic

therapy (18,25). Meanwhile, the improvement of OS in patients with EGFR mutation who progressed from EGFR TKIs within 8 months was significant with anlotinib. Tumors of greater invasiveness are more likely to depend on new blood vessels and benefit more from anti-angiogenic therapy. FGFR overexpression is mechanism of resistance to both EGFR-TKIs and cytotoxic chemotherapy, which is likely why the anlotinib benefit was consistent in patients whose tumors harbor EGFR mutations regardless of whether they were most recently treated with chemotherapy or an EGFR-TKI (26,27). This is also the basis for an ongoing study combining osimertinib and anlotinib for the first-line treatment of patient with advanced EGFR mutated NSCLC (28,29).

In our study, the benefit with anlotinib was consistent in subgroups, independent of EGFR mutation status and previous treatment. With the rapid progress in immunotherapy, checkpoint inhibitors have been applied as first-line therapy, alone or in combination with chemotherapy (30), which may affect the subsequent treatment choice of immunotherapy or chemotherapy. VEGF receptors are present in stromal cells and endothelial cells of the tumor microenvironment, and the action of anti-angiogenic agents on normal cells with a stable genome may be less affected by previous therapy. The ORRs were 7.5% and 10% with anlotinib in patients with and without EGFR mutation, respectively. The DCRs were 82.3% and 80.1% with anlotinib in patients with and without EGFR mutation, respectively, indicating that most patients had stable disease with anlotinib. Anti-VEGF therapy has primarily cytostatic effects and can normalize the tumor vasculature and have substantial systemic effects such as modulation of circulating pro-angiogenic and proinflammatory cytokines and cells, which stabilize the tumor size instead of shrinking it (31).

The safety profile of anlotinib was acceptable and favorable, similar to that described in previous studies (9,10). The most commonly reported TRAEs were typical VEGFR-inhibition related events and were similar in patients with and without EGFR mutation. The mild safety profile might establish the basis for the future combination of anlotinib with other EGFR TKIs such as osimertinib in patients with EGFR mutation, or combination with chemotherapy and immunotherapy in patients without EGFR mutation. The treatment of third-line EGFR mutated lung cancer continues to evolve and treatment strategies in this group now include anlotinib, docetaxel and EGFR-TKI monotherapy rechallenge with additional emerging therapies such as patritumab deruxtecan and

amivantamab (32,33).

The present study had limitations. The standards for first and second-line therapy were altered in both EGFR mutation positive and negative patients. Most EGFR mutation positive patients received first-generation TKIs as second-line therapy in our study. At the start of the study, osimertinib was not approved in China, and patients who progressed from the first-generation TKI were not required to undergo T790M testing and most of them received chemotherapy subsequently without known T790 mutation status, which does not comply with the current guidelines. Likewise, checkpoint inhibitors were not approved during the study, and the efficacy of anlotinib after immunotherapy could not be evaluated. The sample size was small in subgroups, and it was difficult to draw firm conclusions in specific populations. The control arm was placebo, and additional studies are necessary to compare anlotinib with other approved subsequent treatments. Further studies will be conducted to evaluate anlotinib as a subsequent therapy in different populations.

## Conclusions

In conclusion, anlotinib offers a third-line treatment option for patients with advanced NSCLC, regardless of EGFR mutation status and previous therapy strategies.

## Acknowledgments

The authors appreciate the academic support from the AME Lung Cancer Collaborative Group. We thank all the participating patients and their families, the clinical trial team, and all the study personnel.

*Funding:* ALTER0303 was funded by Chia-Tai Tianqing Pharmaceutical Group Co., Ltd. The funder had no role in the study design, data collection and analysis.

## Footnote

*Reporting Checklist:* The authors have completed the TREND reporting checklist. Available at <https://tlcr.amegroups.com/article/view/10.21037/tlcr-22-320/rc>

*Data Sharing Statement:* Available at <https://tlcr.amegroups.com/article/view/10.21037/tlcr-22-320/dss>

*Conflicts of Interest:* All authors have completed the ICMJE uniform disclosure form (available at <https://>



[tcr.amegroups.com/article/view/10.21037/tcr-22-320/coif](https://tcr.amegroups.com/article/view/10.21037/tcr-22-320/coif)). All authors report this study was funded by Chia-Tai Tianqing Pharmaceutical Group Co., Ltd. DB reports that AstraZeneca has paid him for speakers' fees and honoraria, but all funds went to his institution to support research. FT reports research grant from Boehringer Ingelheim Japan, Ono Pharmaceutical, Taiho Pharmaceutical, Chugai Pharmaceutical. FT also reports payment for lectures from MSD, Bristol-Meyers Squibb, Boehringer Ingelheim Japan, Ono Pharmaceutical, Johnson & Johnson, Covidien Japan, Taiho Pharmaceutical, Astra Zeneca, Chugai Pharmaceutical. The authors have no other conflicts of interest to declare.

**Ethical Statement:** The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The study was conducted in accordance with Declaration of Helsinki (as revised in 2013). The study was approved by the institutional ethics board of each site. The informed consent was taken from all the patients.

**Open Access Statement:** This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

- Ferrara N, Gerber HP, LeCouter J. The biology of VEGF and its receptors. *Nat Med* 2003;9:669-76.
- Ferrara N, Kerbel RS. Angiogenesis as a therapeutic target. *Nature* 2005;438:967-74.
- Hall RD, Le TM, Haggstrom DE, et al. Angiogenesis inhibition as a therapeutic strategy in non-small cell lung cancer (NSCLC). *Transl Lung Cancer Res* 2015;4:515-23.
- Sandler A, Gray R, Perry MC, et al. Paclitaxel-carboplatin alone or with bevacizumab for non-small-cell lung cancer. *N Engl J Med* 2006;355:2542-50.
- Garon EB, Ciuleanu TE, Arrieta O, et al. Ramucirumab plus docetaxel versus placebo plus docetaxel for second-line treatment of stage IV non-small-cell lung cancer after disease progression on platinum-based therapy (REVEL): a multicentre, double-blind, randomised phase 3 trial. *Lancet* 2014;384:665-73.
- Manzo A, Montanino A, Carillio G, et al. Angiogenesis Inhibitors in NSCLC. *Int J Mol Sci* 2017;18:2021.
- Sun Y, Niu W, Du F, et al. Safety, pharmacokinetics, and antitumor properties of anlotinib, an oral multi-target tyrosine kinase inhibitor, in patients with advanced refractory solid tumors. *J Hematol Oncol* 2016;9:105.
- Lin B, Song X, Yang D, et al. Anlotinib inhibits angiogenesis via suppressing the activation of VEGFR2, PDGFR $\beta$  and FGFR1. *Gene* 2018;654:77-86.
- Xie C, Wan X, Quan H, et al. Preclinical characterization of anlotinib, a highly potent and selective vascular endothelial growth factor receptor-2 inhibitor. *Cancer Sci* 2018;109:1207-19.
- Han B, Li K, Zhao Y, et al. Anlotinib as a third-line therapy in patients with refractory advanced non-small-cell lung cancer: a multicentre, randomised phase II trial (ALTER0302). *Br J Cancer* 2018;118:654-61.
- Han B, Li K, Wang Q, et al. Effect of Anlotinib as a Third-Line or Further Treatment on Overall Survival of Patients With Advanced Non-Small Cell Lung Cancer: The ALTER 0303 Phase 3 Randomized Clinical Trial. *JAMA Oncol* 2018;4:1569-75. Erratum in: *JAMA Oncol* 2018;4:1625.
- Han B, Tjulandin S, Hagiwara K, et al. EGFR mutation prevalence in Asia-Pacific and Russian patients with advanced NSCLC of adenocarcinoma and non-adenocarcinoma histology: The IGNITE study. *Lung Cancer* 2017;113:37-44.
- Mok TS, Wu YL, Thongprasert S, et al. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med*. 2009;361:947-57.
- Zeng Z, Yan HH, Zhang XC, et al. Reduced chemotherapy sensitivity in EGFR-mutant lung cancer patient with frontline EGFR tyrosine kinase inhibitor. *Lung Cancer* 2014;86:219-24.
- Yu HA, Arcila ME, Rekhtman N, et al. Analysis of tumor specimens at the time of acquired resistance to EGFR-TKI therapy in 155 patients with EGFR-mutant lung cancers. *Clin Cancer Res* 2013;19:2240-7.
- Mok TS, Wu YL, Ahn MJ, et al. Osimertinib or Platinum-Pemetrexed in EGFR T790M-Positive Lung Cancer. *N Engl J Med* 2017;376:629-40.
- Cortinovic D, Gregorc V, Migliorino MR, et al. New perspectives in the second-line treatment of non squamous NSCLC patients: Results from a large Italian Lung Cancer Working Group. *Crit Rev Oncol Hematol*

- 2017;109:35-41.
18. Bylicki O, Paleiron N, Margery J, et al. Targeting the PD-1/PD-L1 Immune Checkpoint in EGFR-Mutated or ALK-Translocated Non-Small-Cell Lung Cancer. *Target Oncol* 2017;12:563-9.
  19. Reck M, Kaiser R, Mellemegaard A, et al. Docetaxel plus nintedanib versus docetaxel plus placebo in patients with previously treated non-small-cell lung cancer (LUME-Lung 1): a phase 3, double-blind, randomised controlled trial. *Lancet Oncol* 2014;15:143-55.
  20. Kerbel RS. Tumor angiogenesis. *N Engl J Med* 2008;358:2039-49.
  21. Langer C, Soria JC. The role of anti-epidermal growth factor receptor and anti-vascular endothelial growth factor therapies in the treatment of non-small-cell lung cancer. *Clin Lung Cancer* 2010;11:82-90.
  22. Lee JS, Hirsh V, Park K, et al. Vandetanib Versus placebo in patients with advanced non-small-cell lung cancer after prior therapy with an epidermal growth factor receptor tyrosine kinase inhibitor: a randomized, double-blind phase III trial (ZEPHYR). *J Clin Oncol* 2012;30:1114-21.
  23. Paz-Ares L, Hirsh V, Zhang L, et al. Monotherapy Administration of Sorafenib in Patients With Non-Small Cell Lung Cancer (MISSION) Trial: A Phase III, Multicenter, Placebo-Controlled Trial of Sorafenib in Patients with Relapsed or Refractory Predominantly Nonsquamous Non-Small-Cell Lung Cancer after 2 or 3 Previous Treatment Regimens. *J Thorac Oncol* 2015;10:1745-53.
  24. Hung MS, Chen IC, Lin PY, et al. Epidermal growth factor receptor mutation enhances expression of vascular endothelial growth factor in lung cancer. *Oncol Lett* 2016;12:4598-604.
  25. Reck M, Paz-Ares L, Bidoli P, et al. Outcomes in patients with aggressive or refractory disease from REVEL: A randomized phase III study of docetaxel with ramucirumab or placebo for second-line treatment of stage IV non-small-cell lung cancer. *Lung Cancer* 2017;112:181-7.
  26. Lian Z, Du W, Zhang Y, et al. Anlotinib can overcome acquired resistance to EGFR-TKIs via FGFR1 signaling in non-small cell lung cancer without harboring EGFR T790M mutation. *Thorac Cancer* 2020;11:1934-43.
  27. Raouf S, Mulford IJ, Frisco-Cabanos H, et al. Targeting FGFR overcomes EMT-mediated resistance in EGFR mutant non-small cell lung cancer. *Oncogene* 2019;38:6399-413.
  28. He Z, Lin J, He Y, et al. The combination of EGFR-TKIs and anlotinib as a first-line therapy for EGFR-mutant advanced non-small cell lung cancer: A multicenter, single-arm, phase II clinical trial. *J Clin Oncol* 2021;39:abstr 9030.
  29. Zhou B, Gong Q, Li B, et al. Clinical outcomes and safety of osimertinib plus anlotinib for patients with previously treated EGFR T790M-positive NSCLC: A retrospective study. *J Clin Pharm Ther* 2022;47:643-51.
  30. Martinez P, Peters S, Stammers T, et al. Immunotherapy for the First-Line Treatment of Patients with Metastatic Non-Small Cell Lung Cancer. *Clin Cancer Res* 2019;25:2691-8.
  31. Crabb SJ, Patsios D, Sauerbrei E, et al. Tumor cavitation: impact on objective response evaluation in trials of angiogenesis inhibitors in non-small-cell lung cancer. *J Clin Oncol* 2009;27:404-10.
  32. Jänne PA, Baik C, Su WC, et al. Efficacy and Safety of Patritumab Deruxtecan (HER3-DXd) in EGFR Inhibitor-Resistant, EGFR-Mutated Non-Small Cell Lung Cancer. *Cancer Discov* 2022;12:74-89.
  33. Shu, C A, et al. Amivantamab plus lazertinib in post-osimertinib, post-platinum chemotherapy EGFR-mutant non-small cell lung cancer (NSCLC): Preliminary results from CHRYSALIS-2. *Ann Oncol* 2021;32:S952-S953.
- (English Language Editor: C. Betlazar-Maseh)

**Cite this article as:** Zhao Y, Wang Q, Zhang L, Shi J, Wang Z, Cheng Y, He J, Shi Y, Chen W, Luo Y, Wu L, Wang X, Nan K, Jin F, Dong J, Li B, Yamaguchi F, Breadner D, Nagano T, Tanaka F, Husain H, Li K, Han B. The efficacy of anlotinib as third-line treatment for non-small cell lung cancer by EGFR mutation status: a subgroup analysis of the ALTER0303 randomized phase 3 study. *Transl Lung Cancer Res* 2022;11(5):776-785. doi: 10.21037/tlcr-22-320

## Supplementary

**Table S1** Multiple Cox regression analysis for PFS in patients with EGFR mutation

Parameter		Parameter estimation	Standard error	P value	HR (95% CI)
Group	Anlotinib	-1.703	0.252	<.0001	0.18 (0.11-0.30)
Number of metastases	≤ 3	-0.394	0.220	0.074	0.68 (0.44-1.04)
Gender	Female	-0.410	0.219	0.061	0.66 (0.43-1.02)
Age (years)	61-69	-0.263	0.235	0.264	0.77 (0.49-1.22)
Age (years)	≥70	-1.557	0.749	0.038	0.21 (0.05-0.92)
Histological type	Others	1.470	0.746	0.049	4.35 (1.01-18.77)
Histological type	Squamous	0.310	0.411	0.451	1.36 (0.61-3.05)
Clinical Stage	IV	-0.798	0.558	0.153	0.45 (0.15-1.34)
Lines of prior chemotherapy	1	-0.033	0.769	0.966	0.97 (0.21-4.37)
Lines of prior chemotherapy	3	-0.254	0.232	0.274	0.78 (0.49-1.22)
Prior target therapy	Yes	-0.583	0.583	0.318	0.56 (0.18-1.75)
Prior radiotherapy	Yes	0.269	0.227	0.235	1.31 (0.84-2.04)
Smoking status	None-smoker	-0.829	0.769	0.281	0.44 (0.10-1.97)
ECOG PS score	1	-0.206	0.277	0.457	0.81 (0.47-1.40)

ECOG PS, Eastern Cooperative Oncology Group Performance Status; EGFR, epidermal growth factor receptor; PFS, progression free survival

**Table S2** Multiple Cox regression analysis for OS in patients with EGFR mutation

Parameter		Parameter estimation	Standard error	P value	HR (95% CI)
Group	Anlotinib	-0.638	0.259	0.014	0.53 (0.32-0.88)
Number of metastases	≤3	-1.095	0.258	<.0001	0.33 (0.20-0.56)
Gender	Female	-0.634	0.243	0.009	0.53 (0.33-0.85)
Age (years)	61-69	-0.041	0.261	0.874	0.96 (0.58-1.60)
Age (years)	≥70	-1.287	0.626	0.040	0.27 (0.08-0.94)
Histological type	Others	1.231	0.756	0.104	3.43 (0.78-15.08)
Histological type	Squamous	-0.948	0.616	0.124	0.39 (0.12-1.30)
Clinical Stage	IV	-1.143	0.707	0.106	0.32 (0.08-1.28)
Lines of prior chemotherapy	1	0.445	0.756	0.555	1.56 (0.36-6.86)
Lines of prior chemotherapy	3	-0.387	0.257	0.131	0.68 (0.41-1.12)
Prior targeted therapy	Yes	0.730	0.701	0.298	2.08 (0.53-8.20)
Prior radiotherapy	Yes	0.218	0.237	0.357	1.24 (0.78-1.98)
Smoking status	None-smoker	-1.057	0.649	0.103	0.35 (0.10-1.24)
ECOG PS score	1	0.302	0.315	0.338	1.35 (0.73-2.51)

ECOG PS, Eastern Cooperative Oncology Group Performance Status; EGFR, epidermal growth factor receptor; OS, overall survival.

**Table S3** Multiple Cox regression analysis for PFS in patients without EGFR mutation


Parameter		Parameter estimation	Standard error	P value	HR (95% CI)
Group	Anlotinib	-1.371	0.161	<.0001	0.25 (0.19-0.35)
Number of metastases	≤3	-0.506	0.143	0.0004	0.60 (0.46-0.80)
Gender	Female	-0.186	0.172	0.279	0.83 (0.59-1.16)
Age (years)	61-69	-0.479	0.150	0.001	0.62 (0.46-0.83)
Age (years)	≥70	-1.093	0.300	0.0003	0.34 (0.19-0.60)
Histological type	Others	-0.134	0.356	0.706	0.87 (0.44-1.76)
Histological type	Squamous	-0.030	0.174	0.860	0.97 (0.69-1.36)
Clinical Stage	IV	-0.0007	0.301	0.998	1.00 (0.55-1.80)
Clinical Stage	Others	-13.078	407.326	0.974	0.00 (0.00-0.00)
Lines of prior chemotherapy	3	-0.034	0.147	0.817	0.98 (0.72-1.29)
Prior targeted therapy	Yes	0.036	0.161	0.822	1.04 (0.76-1.42)
Prior radiotherapy	Yes	-0.214	0.148	0.148	0.81 (0.60-1.08)
Smoking status	None-smoker	-0.090	0.273	0.741	0.91 (0.54-1.56)
ECOG PS score	1	0.249	0.188	0.186	1.28 (0.89-1.85)
ECOG PS score	2	1.647	0.756	0.029	5.19 (1.18-22.85)

ECOG PS, Eastern Cooperative Oncology Group Performance Status; EGFR, epidermal growth factor receptor; PFS, progression free survival.

**Table S4** Multiple Cox regression analysis for OS in patients without EGFR mutation


Parameter		Parameter estimation	Standard error	P value	HR (95% CI)
Group	Anlotinib	-0.370	0.149	0.013	0.69 (0.52-0.93)
Number of metastases	≤3	-0.574	0.144	<.0001	0.56 (0.43-0.75)
Gender	Female	-0.337	0.177	0.056	0.71 (0.51-1.01)
Age (years)	61-69	0.134	0.151	0.377	1.14 (0.85-1.54)
Age (years)	≥70	-0.309	0.312	0.322	0.73 (0.40-1.35)
Histological type	Others	0.059	0.333	0.860	1.06 (0.55-2.04)
Histological type	Squamous	-0.020	0.177	0.912	0.98 (0.69-1.39)
Clinical Stage	IV	0.234	0.333	0.482	1.26 (0.66-2.43)
Clinical Stage	Others	-3.094	1.297	0.017	0.05 (0.004-0.58)
Lines of prior chemotherapy	3	-0.340	0.147	0.020	0.71 (0.53-0.95)
Prior targeted therapy	Yes	0.097	0.164	0.552	1.10 (0.80-1.52)
Prior radiotherapy	Yes	0.010	0.148	0.948	1.01 (0.76-1.35)
Smoking status	None-smoker	-0.186	0.268	0.486	0.83 (0.49-1.40)
ECOG PS score	1	0.644	0.216	0.003	1.91 (1.25-2.91)
ECOG PS score	2	3.902	0.777	<.0001	49.50 (10.80-226.98)

ECOG PS, Eastern Cooperative Oncology Group Performance Status; EGFR, epidermal growth factor receptor; OS, overall survival.




**Figure S1** Overall survival and progression-free survival in patients with EGFR mutation who received EGFR TKIs as first- or second-line therapy. (A) Overall survival in patients with EGFR mutation who received EGFR TKIs as first-line therapy; (B) progression-free survival in patients with EGFR mutation who received EGFR TKIs as first-line therapy; (C) overall survival in patients with EGFR mutation who received EGFR TKIs as second-line therapy; (D) progression-free survival in patients with EGFR mutation who received EGFR TKIs as second-line therapy. EGFR, epidermal growth factor receptor; TKIs, tyrosine kinase inhibitors.





**Figure S2** Overall survival and progression-free survival in patients with EGFR mutation who progressed from EGFR TKIs within or exceeding 8 months. (A) Overall survival in patients with EGFR mutation who progressed from EGFR TKIs within 8 months; (B) progression-free survival in patients with EGFR mutation who progressed from EGFR TKIs within 8 months; (C) overall survival in patients with EGFR mutation who progressed from EGFR TKIs exceeding 8 months; (D) progression-free survival in patients with EGFR mutation who progressed from EGFR TKIs exceeding 8 months. EGFR, epidermal growth factor receptor; TKIs, tyrosine kinase inhibitors.



**Figure S3** Overall survival and progression-free survival in patients with time since TSPT  $\leq 18$  or  $> 18$  months according to EGFR mutation status. (A) Overall survival in patients with TSPT  $\leq 18$  months according to EGFR mutation status; (B) progression-free survival in patients with TSPT  $\leq 18$  months according to EGFR mutation status; (C) overall survival in patients with TSPT  $> 18$  months according to EGFR mutation status; (D) progression-free survival in patients with TSPT  $> 18$  months according to EGFR mutation status. TSPT, the start of prior therapy; EGFR, epidermal growth factor receptor.