



Intracranial responses with selective *KRAS*-G12C inhibitors in non-small cell lung cancer

Kevin Lu, Hatim Husain

University of California, San Diego, La Jolla, CA, USA

Correspondence to: Hatim Husain, MD. Division of Medical Oncology, University of California, San Diego, 3855 Health Sciences #3011, La Jolla, CA 92093, USA. Email: hhusain@ucsd.edu.

Comment on: Sabari JK, Velcheti V, Shimizu K, *et al.* Activity of Adagrasib (MRTX849) in Brain Metastases: Preclinical Models and Clinical Data from Patients with *KRAS*G12C-Mutant Non-Small Cell Lung Cancer. *Clin Cancer Res* 2022;28:3318-28.

Keywords: Non-small cell lung cancer (NSCLC); *KRAS* G12C; adagrasib; sotorasib

Submitted Jan 15, 2023. Accepted for publication May 05, 2023. Published online May 19, 2023.

doi: 10.21037/tlcr-23-26

View this article at: <https://dx.doi.org/10.21037/tlcr-23-26>

The recent Food and Drug Administration (FDA)-approval of adagrasib (MRTX849) has added to the armamentarium of *KRAS* G12C-mutant non-small cell lung cancer (NSCLC). The approval was based on the phase 2 KRYSTAL-1 trial (NCT03785249) which demonstrated clinical benefit in this patient population (1). Adagrasib joins sotorasib (AMG-510) as the only two FDA-approved selective *KRAS*-G12C inhibitors at this time. In phase 2 trials, both adagrasib and sotorasib showed meaningful clinical activity demonstrating a systemic objective response rate (ORR) of 42.9% versus 37.1% respectively (2,3). An area of active investigation in this space surrounds the efficacy of the selective G12C inhibitors in the central nervous system and untreated intracranial lesions.

The initial KRYSTAL-1 and CodeBreak100 trials excluded patients with active, untreated brain metastasis (4). A post-hoc analysis of the CodeBreak100 trial reported intracranial disease control in 14 of 16 patients (87.5%) (5). At 2022 American Society of Clinical Oncology Annual Meeting (ASCO22), the KRYSTAL-1 phase 1b expansion cohort which enrolled active and untreated central nervous system (CNS) metastasis demonstrated an objective intracranial response and disease control rate of 31.6% and 84.2% respectively (6). Intracranial objective responses were observed in 2 of 4 patients with non-target brain lesions and 4 of 15 with target brain lesions. The KRYSTAL-1 phase 2 trial reported an intracranial ORR of 33.3% (11/33 patients) with one intracranial complete response and a

median duration of intracranial response of 11.2 months (2). A baseline CT head with contrast or brain MRI was required for study entry in both phase 2 trials.

In this translational report, Sabari *et al.* demonstrated adagrasib cerebrospinal fluid (CSF) penetration and anti-tumor activity (7). The investigators remark from their cohort of 374 patients that the incidence of brain metastasis in *KRAS*-G12C and non-G12C NSCLC patients to be around 40% which is consistent with other reports (8). Adagrasib has properties which include P-glycoprotein multi-drug resistance (MDR) pump inhibition, suggesting possible CNS penetration. To test this hypothesis, mice models were implanted with H23-Luc or LU65-Luc cell xenografts intracranially. The investigators observed that 200 mg/kg adagrasib dosing (equivalent plasma concentration to 600 mg twice daily in humans) achieved maximum CNS exposure. At 100 mg/kg twice daily dosing, brain lesion regression in animal models was observed including 2 of 5 complete intracranial responses.

In the translational report, unbound brain-to-plasma partition coefficient ($K_{p_{uu, brain}}$), predictor for blood-brain barrier (BBB) permeability, was found to be approximately 1 at 200 mg/kg and 0.2–0.4 at 100 mg/kg. For reference, the anti-EGFR tyrosine kinase inhibitor osimertinib has a $K_{p_{uu, brain}}$ of 0.39 in mice with intra-internal carotid artery injection xenograft (9). Mouse models which received intracranial implantation of xenografts could have compromised BBB integrity and permeability, and

additional model systems including intracardiac, tail vein injections, or spontaneous models would corroborate their findings.

Published sotorasib CNS activity data remains limited at this time. Two case reports describe intracranial response of untreated, active brain metastasis (10,11). In both cases, the patients were treated with sotorasib 960 mg with rapid intracranial response in less than 2 months. Additionally, here we highlight a patient case from our practice involving sotorasib and CNS control. A 74-year-old male with *KRAS* G12C-mutant oligometastatic lung adenocarcinoma involving the left frontal lobe had resection of the brain lesion followed by a course of stereotactic body radiation therapy (SBRT) and immunotherapy (pembrolizumab, carboplatin, pemetrexed) with further CNS progression after a year requiring stereotactic radiation therapy (SRT). The patient was started on sotorasib 960 mg with stable intracranial and improving thoracic cavity disease burden after 1 month. After 5 months of sotorasib initiation, the patient presented with seizures found to have vasogenic edema without new intracranial lesions on brain MRI and was dose reduced to 480 mg daily. A month after the dose reduction, the patient had new brain lesions on MRI and received further SRT. This case reinforces the importance of dose-dependent CNS control of the selective inhibitor. Some combinatorial strategies with *KRAS*-G12C inhibitors are being explored with different doses when combined with anti-PD1 inhibitors. How the effect of dose reduction will impact brain metastasis response and CNS penetration are important areas for future investigation.

The findings of Sabari *et al.* are important as they have demonstrated tumor shrinkage in preclinical models which has been consistent with intracranial responses noted in the KRYSTAL-1 trial. Sabari *et al.* reported in two patients a rapid intracranial response after two cycles of adagrasib (7). An oral presentation at ASCO22 discussed a patient with active, untreated CNS metastasis in the KRYSTAL-1 trial reported to have an intracranial partial response after 1 month and complete response after 5 months of starting adagrasib (6).

Current clinical practice for active focal brain lesions remains stereotactic radiosurgery. Immunotherapy regimens can have CNS activity. A phase 2 trial of monotherapy pembrolizumab in PDL-positive metastatic NSCLC reported a brain metastasis response of 29.7% (11/37) with a 5.7-month duration of CNS response (12). Further studies are needed to evaluate the response rate with selective G12C inhibitors in the context with or without

immunotherapy.

Additional information on the activity of *KRAS*-G12C inhibitors on untreated, active brain metastasis will be important to understand the positioning of this class of medicines in the front-line setting. Further prospective clinical studies are required to fully characterize the CNS efficacy of both adagrasib and sotorasib as approved therapies and other selective G12C inhibitors in development including GDC-6936 (Roche), JDQ443 (Novartis), LY3537982 (Eli Lilly) among others (4).

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Translational Lung Cancer Research*. The article did not undergo external peer review.

Conflicts of Interest: Both authors have completed the ICMJE uniform disclosure form (available at <https://tldr.amegroups.com/article/view/10.21037/tlcr-23-26/coif>). HH reports that he received grants or contracts from BillionToOne, Roche Diagnostics, BMS, and Lilly Oncology, and received consulting fees from Janssen, Astrazeneca, EMD Serono, Foundation Medicine, Mirati, and Amgen. The other author has no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. FDA grants accelerated approval to adagrasib for KRAS

- G12C-mutated NSCLC. FDA, 2022. [cited 2022 Dec 31]. Available online: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-accelerated-approval-adagrasib-kras-g12c-mutated-nsclc>
2. Jänne PA, Riely GJ, Gadgeel SM, et al. Adagrasib in Non-Small-Cell Lung Cancer Harboring a KRAS(G12C) Mutation. *N Engl J Med* 2022;387:120-31.
 3. Skoulidis F, Li BT, Dy GK, et al. Sotorasib for Lung Cancers with KRAS p.G12C Mutation. *N Engl J Med* 2021;384:2371-81.
 4. Palma G, Khurshid F, Lu K, et al. Selective KRAS G12C inhibitors in non-small cell lung cancer: chemistry, concurrent pathway alterations, and clinical outcomes. *NPJ Precis Oncol* 2021;5:98.
 5. Ramalingam S, Skoulidis F, Govindan R, et al. P52.03 Efficacy of Sotorasib in KRAS p.G12C-Mutated NSCLC with Stable Brain Metastases: A Post-Hoc Analysis of CodeBreak 100. *J Thorac Oncol* 2021;16:S1123.
 6. Sabari JK, Spira AI, Heist RS, et al. Activity of adagrasib (MRTX849) in patients with KRASG12C-mutated NSCLC and active, untreated CNS metastases in the KRYSTAL-1 trial. *J Clin Oncol* 2022;40:LBA9009.
 7. Sabari JK, Velcheti V, Shimizu K, et al. Activity of Adagrasib (MRTX849) in Brain Metastases: Preclinical Models and Clinical Data from Patients with KRASG12C-Mutant Non-Small Cell Lung Cancer. *Clin Cancer Res* 2022;28:3318-28.
 8. Cui W, Franchini F, Alexander M, et al. Real world outcomes in KRAS G12C mutation positive non-small cell lung cancer. *Lung Cancer* 2020;146:310-7.
 9. Ballard P, Yates JW, Yang Z, et al. Preclinical Comparison of Osimertinib with Other EGFR-TKIs in EGFR-Mutant NSCLC Brain Metastases Models, and Early Evidence of Clinical Brain Metastases Activity. *Clin Cancer Res* 2016;22:5130-40.
 10. Yeh J, Marks JA, Alzeer AH, et al. Remarkable Intracranial Response to Sotorasib in a Patient With KRASG12C-Mutated Lung Adenocarcinoma and Untreated Brain Metastases: A Case Report. *JTO Clin Res Rep* 2022;3:100428.
 11. Koster KL, Appenzeller C, Lauber A, et al. Sotorasib Shows Intracranial Activity in Patients with KRAS G12C-Mutated Adenocarcinoma of the Lung and Untreated Active Brain Metastases. *Case Rep Oncol* 2022;15:720-5.
 12. Goldberg SB, Schalper KA, Gettinger SN, et al. Pembrolizumab for management of patients with NSCLC and brain metastases: long-term results and biomarker analysis from a non-randomised, open-label, phase 2 trial. *Lancet Oncol* 2020;21:655-63.

Cite this article as: Lu K, Husain H. Intracranial responses with selective *KRAS*-G12C inhibitors in non-small cell lung cancer. *Transl Lung Cancer Res* 2023;12(6):1335-1337. doi: 10.21037/tlcr-23-26