

STROBE Statement—checklist of items that should be included in reports of observational studies

Section/item	Item No	Recommendation	Reported on Page Number/Line Number	Reported on Section/Paragraph
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1 (1) and (3)	Title
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	3/4 (36-61)	Abstract
Introduction				
Background/ rationale	2	Explain the scientific background and rationale for the investigation being reported	5/6 (66-92)	Intro Para 1-3
Objectives	3	State specific objectives, including any prespecified hypotheses	6 (90-92)	Intro Para 3
Methods				
Study design	4	Present key elements of study design early in the paper	6 (94-101)	Methods: Patient Population
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	6 (95), Ref 11	Methods: Patient Population
Participants	6	(a) Cohort study —Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up Case-control study —Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls Cross-sectional study —Give the eligibility criteria, and the sources and methods of selection of participants	6 (95), Ref 11	Methods: Patient Population
		(b) Cohort study —For matched studies, give matching criteria and number of exposed and unexposed Case-control study —For matched studies, give matching criteria and the number of controls per case	N/A	N/A
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	6 (95-109), Ref 11	Methods: Patient Population and Statistical analysis
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	6 (95-109), Ref 11	Methods: Patient Population and Statistical analysis
Bias	9	Describe any efforts to address potential sources of bias	N/A	N/A
Study size	10	Explain how the study size was arrived at	N/A	N/A
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	6 (95-109), Ref 11	Methods: Patient Population and

				Statistical analysis
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	6 (95-109), Ref 11	Methods: Patient Population and Statistical analysis
		(b) Describe any methods used to examine subgroups and interactions	N/A	N/A
		(c) Explain how missing data were addressed	N/A	N/A
		(d) Cohort study —If applicable, explain how loss to follow-up was addressed Case-control study —If applicable, explain how matching of cases and controls was addressed Cross-sectional study —If applicable, describe analytical methods taking account of sampling strategy	N/A	N/A
		(e) Describe any sensitivity analyses	N/A	N/A
Results				
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	6 (95-109), 7 (128), Ref 11	Methods: Patient Population and Results Para 1
		(b) Give reasons for non-participation at each stage	N/A	N/A
		(c) Consider use of a flow diagram	N/A	N/A
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	7/8 (128-135)	Results Para 1
		(b) Indicate number of participants with missing data for each variable of interest	N/A	N/A
		(c) Cohort study —Summarise follow-up time (eg, average and total amount)	N/A	N/A
Outcome data	15*	Cohort study —Report numbers of outcome events or summary measures over time	8 (136-138)	Results Para 2
		Case-control study —Report numbers in each exposure category, or summary measures of exposure	N/A	N/A
		Cross-sectional study —Report numbers of outcome events or summary measures	N/A	N/A
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	8 (136-138)	Results Para 2
		(b) Report category boundaries when continuous variables were categorized	N/A	N/A
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	N/A	N/A
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	8-9 (139-170)	Results Para 3-7

Discussion				
Key results	18	Summarise key results with reference to study objectives	9-12 (171-247)	Discussion Para 1-6
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	12 (241-247)	Discussion Para 7
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	13 (249-252)	Conclusion
Generalisability	21	Discuss the generalisability (external validity) of the study results	12 (241-247)	Discussion Para 7
Other information				
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	14 (271-264)	Acknowledgements

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

Article information: <https://dx.doi.org/10.21037/tlcr-23-383>

*As the checklist was provided upon initial submission, the page number/line number reported may be changed due to copy editing and may not be referable in the published version. In this case, the section/paragraph may be used as an alternative reference.