



Alectinib continuation beyond progression in *ALK*-positive non-small cell lung cancer with alectinib-refractory

Yimeng Li[#], Zhanpeng Hao[#], Yuyan Ma[#], Kaidiriye Setiwalidi, Yingming Zhang, Yujia Zhao, Xiao Fu, Xuan Liang, Zhiping Ruan, Tao Tian, Yu Yao

Department of Medical Oncology, The First Affiliated Hospital of Xi'an Jiaotong University, Xi'an, China

Contributions: (I) Conception and design: Y Yao, Z Ruan, X Liang, T Tian, X Fu; (II) Administrative support: Y Yao, T Tian; (III) Provision of study materials or patients: Y Li, Z Hao, Y Ma; (IV) Collection and assembly of data: Y Li, Y Ma, Z Hao, K Setiwalidi, Y Zhang, Y Zhao; (V) Data analysis and interpretation: Y Li, Y Ma, Z Hao; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

[#]These authors contributed equally to this work.

Correspondence to: Yu Yao, MD; Tao Tian, MD. Department of Medical Oncology, The First Affiliated Hospital of Xi'an Jiaotong University, No. 277 Yanta West Road, Xi'an 710061, China. Email: 13572101611@163.com; tiantao0607@163.com.

Background: Alectinib, a next-generation anaplastic lymphoma kinase tyrosine kinase inhibitor (ALK-TKI), has demonstrated noteworthy efficacy in the treatment of non-small cell lung cancer (NSCLC). Unfortunately, 53.3% of untreated patients receiving first-line treatment with alectinib developed resistance to alectinib. However, despite the widespread use of alectinib, studies on the efficacy and safety of continuing alectinib with other necessary therapies after progression of alectinib and possible population of benefit are still limited.

Methods: This retrospective cohort study included fifteen patients with *ALK*-positive NSCLC from nine institutions in China who experienced disease progression after first- or second-line treatment and continued to receive alectinib treatment between 2019 and 2022. This study aimed to evaluate the median progression-free survival (mPFS), objective response rate (ORR), median overall survival (mOS), and adverse events (AEs) of continuing alectinib combined with other therapies after the emergence of drug resistance.

Results: Among fifteen patients eligible for this study, all patients started continuing treatment with alectinib after oligoprogression or central nervous system (CNS) progression. The mPFS for the whole cohort receiving continuing alectinib with other necessary therapies was 8 months [95% confidence interval (CI): 4 to not applicable (NA)], with an ORR of 46.7%. The mOS was not reached. During continuing alectinib treatment, only one patient experienced grade 2 elevation of aspartate aminotransferase (AST) and serum glutamic-oxaloacetic transaminase (SGOT).

Conclusions: The continuation of alectinib treatment combined with other necessary therapies demonstrates favorable response and safety in patients with *ALK*-positive NSCLC who experienced oligoprogression or CNS progression following alectinib in first- or second-line therapy. Instead of immediately switching to another ALK-TKI, continuing alectinib combined with other necessary therapies may offer greater survival benefits to the patients.

Keywords: Alectinib; anaplastic lymphoma kinase-tyrosine kinase inhibitor (ALK-TKI); non-small cell lung cancer (NSCLC); *ALK*-positive; resistance

Submitted Dec 01, 2023. Accepted for publication Jan 11, 2024. Published online Jan 22, 2024.

doi: 10.21037/tlcr-23-798

View this article at: <https://dx.doi.org/10.21037/tlcr-23-798>

Introduction

Rearrangement of the anaplastic lymphoma kinase (*ALK*) gene is a significant oncogenic driver in patients with non-small cell lung cancer (NSCLC), and it has been confirmed to be a molecular target for this disease. Approximately 3–7% of all NSCLC cases are characterized by *ALK*-positive (1,2). With the rapid development of antitumor therapy, tyrosine kinase inhibitors (TKIs) that specifically act on the *ALK* receptor tyrosine kinase have shown notable efficacy in treating patients with *ALK*-positive NSCLC (2–8). The remarkable efficacy and safety profile of these *ALK*-TKIs has led to the accelerated development of three generations of *ALK* inhibitors in less than a decade (9–12).

Among the *ALK*-TKIs, alectinib, a next-generation *ALK*-TKI, has demonstrated remarkable efficacy in comparison to crizotinib, a first-generation *ALK*-TKI (10,13). The global ALEX study reported a median progression-free survival (mPFS) of 34.8 months and an objective response rate (ORR) of 82.9% for alectinib in untreated *ALK*-positive NSCLC patients (10,13,14). Furthermore, the Asian population-based ALESIA study

showed even better efficacy, with an mPFS of 41.6 months and an ORR of 91%. However, despite the significant clinical benefits achieved with alectinib, resistance to alectinib occurred in 53.3% of patients treated with alectinib as the first-line therapy, leading to disease progression, and more progressive disease occurred in second-line alectinib treatment (4,15–17). *ALK*-TKIs resistance mainly arises from *ALK* mutations and *ALK* amplification, such as the *G1202R* mutation (18). In addition, resistance can occur due to bypass activation or downstream pathway activation, such as *MET*-amplification, or phenotypical changes (18–20). Subsequent therapeutic decision-making will be based firstly on the type of progression and secondly on the mechanism of resistance (4,21). Progression can be divided into oligoprogression, central nervous system (CNS) progression, and systemic progression (4,17). Oligoprogression is defined as a tumor disease that progresses in a limited number of sites (17). CNS progression is defined as progression involving only the CNS. When the disease suffers progression in more sites, it is defined as systemic progression (17). It can be considered that oligoprogression or CNS progression does not necessarily indicate a complete failure of the prior TKI treatment, it may be due to heterogeneity within and between tumors. However, systemic progression often represents the failure of the prior TKI treatment (21). If systemic progression occurs, medication selections are made to target the mechanism of acquired resistance (4). According to established guidelines, patients with *ALK*-positive advanced NSCLC who experience oligoprogression or CNS progression during TKI treatment have the option to continue with their current TKI therapy combined with local therapies or switch to an alternative *ALK*-TKI (4,21). However, several studies have indicated that sequential *ALK*-TKI treatment may ultimately result in the development of highly resistant complex *ALK* mutations, which could potentially limit the survival benefits of patients (22). Fortunately, some clinical studies have shown the superiority of continuing the original *ALK*-TKI treatment combined with other treatments after the progression of next-generation *ALK*-TKI (18,23). Lin *et al.* demonstrated that in patients who underwent chemotherapy in combination with *ALK*-TKI after developing resistance to next-generation *ALK*-TKI, the mPFS was 6.8 months, which was significantly longer than the patients who received chemotherapy alone (6.8 *vs.* 3.2 months). In a Phase II clinical trial investigating non-squamous NSCLC

Highlight box

Key findings

- Alectinib continuation beyond progression of first- or second-line alectinib in patients with anaplastic lymphoma kinase (*ALK*)-positive non-small cell lung cancer (NSCLC) resulted in an 8-month median progression-free survival (mPFS) and a favorable safety profile.

What is known and what is new?

- Alectinib demonstrates outstanding efficacy and a favorable safety profile as a next-generation *ALK*-tyrosine kinase inhibitor (TKI). It could provide an mPFS of 34.8 months. However, 53.3% of patients develop resistance to this drug according to ALEX research.
- We investigated the efficacy, safety, and population most likely to benefit from continuing alectinib with other necessary therapies in *ALK*-positive NSCLC patients who developed resistance to prior alectinib therapy.

What is the implication, and what should change now?

- Treatment consisting of continuing alectinib and other necessary therapies in *ALK*-positive NSCLC patients who suffered oligoprogression or central nervous system progression using alectinib yielded favorable efficacy and good safety. In this scenario, the option of continuing alectinib therapy in combination with other necessary therapies may be favorable over switching to another TKI.

patients with combined alectinib and bevacizumab following resistance to alectinib, the study resulted in an mPFS of 3.1 months (23). The above studies suggest that even if disease progression occurs, tumors can remain responding to TKIs (18,23). This may be due to the persistence of TKI-sensitive clones (24). In this scenario, considering the circumstances, it may be an advisable choice to continue the existing TKI treatment and explore the potential benefits of combining it with other therapeutic approaches. However, despite the widespread use of alectinib, studies on the efficacy and safety of continuing alectinib with other necessary therapies after progression of alectinib and possible population of benefit are still limited.

In this study, we conducted a real-world investigation to assess the efficacy, safety, and the population most likely to benefit of continuing alectinib treatment with other necessary therapies in patients with *ALK*-positive NSCLC who have experienced disease progression following initial or second-line alectinib therapy. We present this article in accordance with the STROBE reporting checklist (available at <https://tldr.amegroups.com/article/view/10.21037/tlcr-23-798/rc>).

Methods

Patient population

We enrolled eligible patients in this study who had histologically or cytologically confirmed stage IV NSCLC and who tested positive for *ALK* using fluorescence *in situ* hybridization, immunohistochemical analysis, or next-generation gene sequencing. These patients had previously received first- or second-line alectinib therapy and experienced disease progression as assessed based on periodic and radiological surveillance. The study excluded patients who had uncertain tumor conditions and who were lost to follow-up.

Treatment

Patients received continuing alectinib treatment from the time of disease progression on prior alectinib until repeated disease progression, an unacceptable adverse event (AE), or death occurred. If necessary, additional systemic or local therapies were administered concurrently with alectinib treatment.

Study design

This retrospective cohort study included a total of fifteen patients from nine institutions in China due to the rarity of the study population. These patients experienced disease progression after first- or second-line alectinib and continued to receive alectinib treatment between 2019 and 2022. This study was approved by the Ethics Review Committee of the First Affiliated Hospital of Xi'an Jiaotong University (No. XJTU1AF2023LSK-364) and adhered to the ethical guidelines of the Helsinki Declaration (as revised in 2013). Individual consent for this retrospective analysis was waived. All participating institutions were informed and agreed with this study. The primary objective of the study was to evaluate the safety and the mPFS of the continuing alectinib, as assessed by the investigators. Secondary endpoints included the ORR of alectinib continuation and the median overall survival (mOS) of the study participants as assessed by the investigators. To assess tumor response, computed tomography (CT) or magnetic resonance imaging (MRI) was applied every 2 months and evaluated according to Response Evaluation Criteria in Solid Tumors (RECIST) v. 1.1 guidelines. AEs are recorded in terms of type and severity, and the severity of toxicity was evaluated using the Common Terminology Criteria for Adverse Events v. 5.0. The study was a single-arm cohort study that was compared with previous studies.

Statistical analysis

PFS was considered to be the time to progression or death following initial progression with alectinib. OS was calculated from the initiation of alectinib treatment to the time of death. ORR was considered to be the ORR during the continuation of alectinib. Kaplan-Meier survival curves were generated for PFS and OS. The “swimplot” package and “ggplot” package in R software were used for creating swim plots. All statistical analyses were conducted using R software version 4.2.3 and Jamovi software version 2.3.18 for Microsoft Windows 64.

Results

Patient characteristics

A total of fifteen patients were enrolled in the study

Table 1 Clinical and pathologic characteristics of the study cohort receiving alectinib for the first time (n=15)

Characteristic	Value
Sex	
Male	5 (33.3)
Female	10 (66.7)
Age at diagnosis, years	43 [17–57]
Smoking history	
Yes	1 (6.7)
None	11 (73.3)
Unknown	3 (20.0)
Histology (adenocarcinoma)	15 (100.0)
TNM stage at diagnosis (stage IV)	15 (100.0)
Brain metastases at diagnosis	4 (26.7)
Bone metastases at diagnosis	8 (53.3)
Line of prior alectinib	
1	13 (86.7)
2	2 (13.3)
Prior exposure to chemotherapy before alectinib	
Yes	2 (13.3)
None	13 (86.7)
Prior exposure to ALK-TKI before alectinib	
Yes	3 (20.0)
None	12 (80.0)
ECOG performance status	
1	8 (53.3)
2	6 (40.0)
3	0
4	1 (6.7)

Data are presented as n (%) or median [range]. TNM, tumour-nodes-metastasis; ALK-TKI, anaplastic lymphoma kinase-tyrosine kinase inhibitor; ECOG, Eastern Cooperative Oncology Group.

between 2019 and 2022. Patient characteristics at the baseline of the first alectinib treatment are presented in *Table 1*. Out of the fifteen patients, ten were female, and the median age was 43 years. Among them, only one patient had a specific history of smoking. All the patients had advanced *ALK*-positive lung adenocarcinoma. In the

baseline, there were four patients with brain metastases and eight patients with bone metastases. Thirteen patients received first-line treatment with alectinib, two patients were treated with alectinib as second-line therapy. Two patients had prior exposure to chemotherapy and three patients had previously exposed to ALK-TKI. As shown in *Table S1*, patient No. 9 received one cycle of chemotherapy after diagnosis while awaiting the result of the genetic test, followed by crizotinib in combination with cranial gamma knife radiosurgery and alectinib after the progression of the prior treatment. Patient No. 3 was treated with crizotinib for 2.5 months, which was then discontinued based on individual decision, followed by alectinib for better efficacy. Patient No. 2 received one cycle of chemotherapy during the post-diagnostic genetic test and thereafter received alectinib. Other patients who received first-line treatment with alectinib were previously untreated. The Eastern Cooperative Oncology Group (ECOG) performance status of fourteen patients ranged from 1 to 2, indicating relatively good performance status, while one patient had an ECOG performance status of 4.

Table 2 presents the baseline characteristics of patients who continued treatment with alectinib. Thirteen patients achieved partial response at the prior alectinib treatment, and two patients achieved stable disease. Seven patients started continuing alectinib after oligoprogression, while eight patients continued alectinib because of CNS progression. Six patients developed new intracranial lesions, and one patient developed new bone metastases. Fourteen patients had an ECOG performance status of 1–2, while one patient had an ECOG performance status of 3. Among the fifteen patients, ten were treated with a combination of radiotherapy and continuing alectinib treatment, two received alectinib monotherapy, and the remaining three underwent chemotherapy, bevacizumab, and radiofrequency ablation, respectively. The median follow-up time for OS was 35 months [95% confidence interval (CI): 32 to not applicable (NA)] for all patients.

Efficacy and safety

All patients exhibited measurable disease as per the RECIST criteria. At the final follow-up on May 31, 2023, eleven patients experienced progressive disease (PD), and among the eleven patients, two deaths occurred. *Figure 1* shows that the mPFS of continuing alectinib combined with other necessary therapies was 8 months (95% CI: 4 to NA), while the mOS was not reached. *Figure 2A,2B* provide

Table 2 Clinical and pathologic characteristics of the study cohort continuing to receive alectinib (n=15)

Characteristic	Value
Age at continuation of alectinib, years	44 [18–59]
Best response to prior alectinib therapy	
SD	2 (13.3)
PR	13 (86.7)
Progression type of prior alectinib	
Oligoprogression	7 (46.7)
CNS progression	8 (53.3)
New brain metastases at PD	6 (40.0)
New bone metastases at PD	1 (6.7)
ECOG performance status	
1	8 (53.3)
2	6 (40.0)
3	1 (6.7)
4	0
Combination therapy	
None	2 (13.3)
Bevacizumab	1 (6.7)
Radiation	10 (66.7)
Chemotherapy	1 (6.7)
Radiofrequency ablation	1 (6.7)

Data are presented as n (%) or median [range]. SD, stable disease; PR, partial response; CNS, central nervous system; PD, progressive disease; ECOG, Eastern Cooperative Oncology Group.

details of patients receiving continued treatment with alectinib in combination with other necessary therapies. During the continuation of alectinib, seven patients achieved partial response (PR), six had stable disease (SD), and two experienced PD, resulting in an ORR of 46.7% (Figure 2C).

These patients demonstrated a favorable safety profile during initial treatment with alectinib. One patient experienced a grade 1 skin rash and grade 1 elevation in alanine aminotransferase (ALT), while another patient experienced a grade 2 elevation in bilirubin. Both patients showed improvement with symptomatic supportive therapy, and no dose reduction or discontinuation of alectinib was necessary. Additionally, one patient underwent a dose reduction due to the development of a grade 2 skin rash. Among the patients who continued receiving alectinib, only one patient experienced a grade 2 elevation of aspartate aminotransferase (AST) and serum glutamic-oxaloacetic transaminase (SGOT), which also improved with symptomatic supportive therapy without the need for dose reduction or discontinuation.

Discussion

Our study demonstrated a cohort who received continuing alectinib combined with other therapies after prior alectinib resistance in the first- or second-line treatment, achieving an mPFS of 8 months. All patients started continuing alectinib because of oligoprogression or CNS progression. Eleven (73.3%) patients combined local treatment, and two patients received systemic therapy. Notably, two patients continued monotherapy with alectinib after progression,




Figure 1 Kaplan-Meier curves for progression-free survival (A) and overall survival (B). CI, confidence interval; NA, not applicable; PFS, progression-free survival; OS, overall survival.




Figure 2 The swimmer plot and pie plot of continuing alectinib with other therapies in patients with alectinib-refractory *ALK*-positive NSCLC. (A) Progression-free survival and progression of continuing alectinib with other therapies. (B) Progression-free survival and progression of different combination therapies with continued alectinib. (C) Best tumor response to alectinib continuation with other necessary therapies. *ALK*, anaplastic lymphoma kinase; NSCLC, non-small cell lung cancer; PD, progressive disease; PFS, progression-free survival; RFA, radiofrequency ablation; SD, stable disease; PR, partial response.

considering their favorable response to the prior alectinib treatment. Until the end of the follow-up, they had stable disease and had not experienced progression. According to guidelines, when oligoprogression or CNS progression occurs in advanced NSCLC patients on *ALK*-TKIs, patients can continue existing *ALK*-TKI combined with local therapy or switch to another *ALK*-TKI. However, sequential *ALK*-TKI treatment may ultimately result in the development of highly resistant complex *ALK* mutations, which could potentially limit patient survival benefits (22). Consequently, the potential for continuing alectinib in combination with other therapies should be explored.

Previous studies have demonstrated the persistence of

TKI-sensitive clones in lung cancers that have developed resistance to TKIs (24-29). The current definition of acquired resistance is the occurrence of disease progression after 6 months of ongoing TKI therapy (4). However, it is necessary to confirm this resistance before switching therapies for the benefit of the patients (30). It is well-known that prolonged exposure to epidermal growth factor receptor (EGFR) inhibitors can lead to the emergence of a secondary *T790M* mutation, which restores EGFR phosphorylation in the presence of TKI (31-33). Recent findings indicate that discontinuation of TKI in cell lines and patients with acquired resistance results in a gradual loss of the *T790M* mutation, allowing these cells to regain

sensitivity to EGFR-TKI (25,26). This phenomenon might be attributed to the inactivation of *T790M*-mutant cell growth, leading to the overgrowth of parental cells that carry only *EGFR*-sensitizing mutations. Another possibility is that *EGFR* alleles are located extrachromosomally in double minutes, and they can be lost from the cells without appropriate selection pressure (25,27). These observations suggest the potential reversibility of acquired resistance and give clinicians mechanistic evidence to continue the original TKI therapy. As a matter of fact, it is not only in the laboratory but also in clinical practice. Chaft *et al.* summarized case reports of gefitinib-resistant patients with *EGFR*-mutated NSCLC who were reintroduced to gefitinib, indicating that the majority of patients retained sensitivity to gefitinib, which was attributed to the presence of tumor cells that exhibit partial resistance to EGFR-TKI (24). Ultimately, increasing the TKI dosage could overcome the acquired resistance in these patients (27,28). In an American cohort of patients with relapsed *EGFR*-mutant NSCLC, some patients experienced relapse after adjuvant treatment with EGFR-TKI, but the majority still responded favorably to EGFR-TKI reintroduction (29). This further supports the aforementioned hypothesis.

In other patients with NSCLC treated with ALK-TKI, similar findings have been observed. A study conducted in the United States reported on a group of patients who remained on their initial treatment of crizotinib and erlotinib in combination with local ablative therapy after developing acquired resistance. The mPFS for this cohort was 6.2 months (34). Another American cohort of 120 patients continued on crizotinib treatment despite crizotinib resistance, and this group had a longer mOS from the first dose of crizotinib (29.6 *vs.* 10.8 months) compared to patients who discontinued treatment after acquiring resistance (35). Hong *et al.* reported on a Chinese cohort for whom crizotinib therapy was continued after resistance had developed (36). This cohort achieved an mPFS of 16 weeks, patients who received local therapy after disease progression had a significantly longer PFS ($P=0.039$), and patients with a longer PFS in prior TKI treatment also experienced a longer PFS in continuing TKI therapy (36). Similarly, in a Chinese cohort studied by Lei *et al.*, patients who continued on crizotinib treatment after experiencing CNS progression achieved an mPFS of 6.3 months (37). Most of these patients received localized CNS therapy, including whole-brain radiation therapy or stereotactic radiosurgery (37). In another cohort of patients resistant to alectinib, those who received a platinum-pemetrexed combination with ALK-TKI had a

longer duration of PFS compared to those receiving platinum or pemetrexed alone (6.8 *vs.* 3.2 months) (18). Additionally, in the phase II clinical NLCTG1501 trial, twelve alectinib-resistant patients treated with bevacizumab in combination with alectinib achieved an mPFS of 3.1 months (95% CI: 1.2–16.1) and an mOS of 24.1 months (95% CI: 8.3 to NA). These findings are in line with our study.

Alectinib is currently recommended as the first-line treatment for patients with *ALK*-positive NSCLC in international guidelines due to its broad application base. Consequently, there is a significant clinical demand for addressing resistance to this treatment (38). At present, when patients develop resistance to next-generation TKI, the decision-making to replace another TKI depends on the mechanism of acquired resistance, such as the location of the resistance within the *ALK* domain or off-target (4). If the resistance is caused by mutations such as *G1202R*, lorlatinib becomes the primary treatment option, leading to an mPFS of approximately 10 months (38–42). However, if the resistance occurs outside the kinase domain, chemotherapy is considered the best choice, resulting in an mPFS of around 3 months (18,43). According to some research findings, patients who switch to other next-generation ALK-TKI after resistance can achieve an mPFS of approximately 3 months (44,45). This suggests that switching to other next-generation ALK-TKI may have some benefits in extending disease control time in cases of resistance (44,45). The mPFS obtained in our study was 8 months. When alectinib becomes resistant again, it can still be used in combination with chemotherapy or switched to another ALK-TKI such as lorlatinib, depending on the underlying resistance mechanism. This approach has the potential to prolong OS in patients.

Certainly, the decision-making of combination therapy should be established based on the specific disease status and mechanisms of drug resistance. For instance, if an isolated brain metastatic lesion progresses, radiotherapy is usually preferred to control the brain metastatic lesion due to the existence of the blood-brain barrier (46,47). Similarly in our study, a total of 75% of patients (6/8) with CNS progression received cranial radiotherapy, and they exhibited positive responses to the treatment. Other localized lesion control can also rely on radiotherapy (48,49). When there is progression of multiple lesions or resistance occurs outside the kinase domain, despite oligoprogression, CNS progression rather than systemic progression, systemic treatments such as chemotherapy and antiangiogenesis therapy will likely need to be considered to control all

lesions (18,23). Additionally, it is worth further exploration in future studies to determine the most appropriate treatment strategy by exploring the synergistic effects of alectinib and radiotherapy, chemotherapy, or antiangiogenic drugs to make an accurate decision. Our study provides a significantly beneficial therapy for alectinib-resistant patients. It balances the treatment of TKI-sensitive clones and possible off-target TKI-insensitive clones, avoiding the generation of complex resistance and providing more possibilities for subsequent ALK-TKI such as lorlatinib. Unfortunately, in this study, it is disappointing to see that several patients had limited benefits from continuing alectinib. It is necessary to explore the biomarkers associated with the benefits of continuing alectinib to guide clinical practice more accurately (50). In the study by Ou *et al.*, patients who continued treatment with the original TKI after crizotinib resistance had a better ECOG performance status and a better ORR on prior crizotinib therapy compared to patients who switched to another TKI, which might give us some insights (35). However, we were limited by the fact that our study only included patients who continued alectinib and failed to compare the differences.

Our study has certain limitations, many of which are inherent to the retrospective study designs. First, the study had a limited sample size, consisting only of a small number of patients. Since all patients were recruited from academic institutions, the findings may not be fully representative of the broader *ALK*-positive population. Second, the mechanisms of resistance in most patients with disease progression are unknown due to the low frequency of re-biopsies, especially when progressive metastases occur in the brain, where biopsies cannot be conveniently conducted. Moreover, it is rare for re-biopsy to be performed in patients with oligoprogression, considering the existence of TKI-sensitive clones, the progression is usually due to tumor heterogeneity. Therefore, we have not been able to explore the population that might benefit from this therapy from the results of genetic tests. Furthermore, retrospective studies are prone to limitations in data collection, potentially leading to incomplete or insufficient data. Finally, the physician's decision to continue alectinib treatment might have been influenced by various factors, such as the patient's ECOG performance status, the extent of the CNS and extracranial disease, tumor genotyping, smoking history, and the patient's response and tolerability to previous ALK-TKI. These confounding factors might have influenced the observed PFS, which is the main focus of our study.

Conclusions

In our study, we found that continuing alectinib treatment combined with other necessary therapy after the oligoprogression or CNS progression of the first- or second-line alectinib in patients with *ALK*-positive NSCLC yielded a favorable response and maintained safety. Our findings suggest that instead of immediately switching to another ALK-TKI, continuing alectinib combined with other necessary therapies after the oligoprogression or CNS progression of the first- or second-line alectinib treatment may offer greater survival benefits to patients.

Acknowledgments

Funding: This work was financially supported by the Youth Program of the National Natural Science Foundation of China (No. 82002437), CSCO-Hengrui Cancer Research Foundation (No. Y-HR2019-0401), and the Medical Scientific Research Project (Medical Research Project for Young and Middle-Aged Oncologists of Lung Cancer).

Footnote

Reporting Checklist: The authors have completed the STROBE reporting checklist. Available at <https://tldr.amegroups.com/article/view/10.21037/tlcr-23-798/rc>

Data Sharing Statement: Available at <https://tldr.amegroups.com/article/view/10.21037/tlcr-23-798/dss>

Peer Review File: Available at <https://tldr.amegroups.com/article/view/10.21037/tlcr-23-798/prf>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://tldr.amegroups.com/article/view/10.21037/tlcr-23-798/coif>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. This retrospective study was conducted in accordance with the Declaration of Helsinki (as revised in 2013) and was approved by the Ethics Committee of the First Affiliated Hospital of Xi'an Jiaotong University (No. XJTU1AF2023LSK-364). Individual consent for this retrospective analysis was waived.

All participating institutions were informed and agreed with this study.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- Hallberg B, Palmer RH. The role of the ALK receptor in cancer biology. *Ann Oncol* 2016;27 Suppl 3:iii4-iii15.
- Schneider JL, Lin JJ, Shaw AT. ALK-positive lung cancer: a moving target. *Nat Cancer* 2023;4:330-43.
- Katayama R. Therapeutic strategies and mechanisms of drug resistance in anaplastic lymphoma kinase (ALK)-rearranged lung cancer. *Pharmacol Ther* 2017;177:1-8.
- Lin JJ, Riely GJ, Shaw AT. Targeting ALK: Precision Medicine Takes on Drug Resistance. *Cancer Discov* 2017;7:137-55.
- Rizzo A, Cusmai A, Giovannelli F, et al. Impact of Proton Pump Inhibitors and Histamine-2-Receptor Antagonists on Non-Small Cell Lung Cancer Immunotherapy: A Systematic Review and Meta-Analysis. *Cancers (Basel)* 2022;14:1404.
- Rizzo A. Identifying optimal first-line treatment for advanced non-small cell lung carcinoma with high PD-L1 expression: a matter of debate. *Br J Cancer* 2022;127:1381-2.
- Santoni M, Rizzo A, Mollica V, et al. The impact of gender on The efficacy of immune checkpoint inhibitors in cancer patients: The MOUSEION-01 study. *Crit Rev Oncol Hematol* 2022;170:103596.
- Mollica V, Rizzo A, Marchetti A, et al. The impact of ECOG performance status on efficacy of immunotherapy and immune-based combinations in cancer patients: the MOUSEION-06 study. *Clin Exp Med* 2023;23:5039-49.
- Solomon BJ, Mok T, Kim DW, et al. First-line crizotinib versus chemotherapy in ALK-positive lung cancer. *N Engl J Med* 2014;371:2167-77.
- Mok T, Camidge DR, Gadgeel SM, et al. Updated overall survival and final progression-free survival data for patients with treatment-naive advanced ALK-positive non-small-cell lung cancer in the ALEX study. *Ann Oncol* 2020;31:1056-64.
- Shaw AT, Bauer TM, de Marinis F, et al. First-Line Lorlatinib or Crizotinib in Advanced ALK-Positive Lung Cancer. *N Engl J Med* 2020;383:2018-29.
- Camidge DR, Kim HR, Ahn MJ, et al. Brigatinib versus Crizotinib in ALK-Positive Non-Small-Cell Lung Cancer. *N Engl J Med* 2018;379:2027-39.
- Peters S, Camidge DR, Shaw AT, et al. Alectinib versus Crizotinib in Untreated ALK-Positive Non-Small-Cell Lung Cancer. *N Engl J Med* 2017;377:829-38.
- Remon J, Pignataro D, Novello S, et al. Current treatment and future challenges in ROS1- and ALK-rearranged advanced non-small cell lung cancer. *Cancer Treat Rev* 2021;95:102178.
- Zia V, Lengyel CG, Tajima CC, et al. Advancements of ALK inhibition of non-small cell lung cancer: a literature review. *Transl Lung Cancer Res* 2023;12:1563-74.
- Desai A, Lovly CM. Strategies to overcome resistance to ALK inhibitors in non-small cell lung cancer: a narrative review. *Transl Lung Cancer Res* 2023;12:615-28.
- Pisano C, De Filippis M, Jacobs F, et al. Management of Oligoprogression in Patients with Metastatic NSCLC Harboring ALK Rearrangements. *Cancers (Basel)* 2022;14:718.
- Lin JJ, Schoenfeld AJ, Zhu VW, et al. Efficacy of Platinum/Pemetrexed Combination Chemotherapy in ALK-Positive NSCLC Refractory to Second-Generation ALK Inhibitors. *J Thorac Oncol* 2020;15:258-65.
- Dagogo-Jack I, Kiedrowski LA, Heist RS, et al. Efficacy and Tolerability of ALK/MET Combinations in Patients With ALK-Rearranged Lung Cancer With Acquired MET Amplification: A Retrospective Analysis. *JTO Clin Res Rep* 2023;4:100534.
- Urbanska EM, Sørensen JB, Melchior LC, et al. Changing ALK-TKI-Resistance Mechanisms in Rebiopsies of ALK-Rearranged NSCLC: ALK- and BRAF-Mutations Followed by Epithelial-Mesenchymal Transition. *Int J Mol Sci* 2020;21:2847.
- Campo M, Al-Halabi H, Khandekar M, et al. Integration of Stereotactic Body Radiation Therapy With Tyrosine Kinase Inhibitors in Stage IV Oncogene-Driven Lung Cancer. *Oncologist* 2016;21:964-73.
- Yoda S, Lin JJ, Lawrence MS, et al. Sequential ALK Inhibitors Can Select for Lorlatinib-Resistant Compound ALK Mutations in ALK-Positive Lung Cancer. *Cancer Discov* 2018;8:714-29.
- Watanabe S, Sakai K, Matsumoto N, et al. Phase II Trial

- of the Combination of Alectinib with Bevacizumab in Alectinib Refractory ALK-Positive Nonsquamous Non-Small-Cell Lung Cancer (NLCTG1501). *Cancers (Basel)* 2022;15:204.
24. Chaft JE, Oxnard GR, Sima CS, et al. Disease flare after tyrosine kinase inhibitor discontinuation in patients with EGFR-mutant lung cancer and acquired resistance to erlotinib or gefitinib: implications for clinical trial design. *Clin Cancer Res* 2011;17:6298-303.
 25. Chmielecki J, Foo J, Oxnard GR, et al. Optimization of dosing for EGFR-mutant non-small cell lung cancer with evolutionary cancer modeling. *Sci Transl Med* 2011;3:90ra59.
 26. Sequist LV, Waltman BA, Dias-Santagata D, et al. Genotypic and histological evolution of lung cancers acquiring resistance to EGFR inhibitors. *Sci Transl Med* 2011;3:75ra26.
 27. Ercan D, Zejnullahu K, Yonesaka K, et al. Amplification of EGFR T790M causes resistance to an irreversible EGFR inhibitor. *Oncogene* 2010;29:2346-56.
 28. Watanabe S, Tanaka J, Ota T, et al. Clinical responses to EGFR-tyrosine kinase inhibitor retreatment in non-small cell lung cancer patients who benefited from prior effective gefitinib therapy: a retrospective analysis. *BMC Cancer* 2011;11:1.
 29. Oxnard GR, Janjigian YY, Arcila ME, et al. Maintained sensitivity to EGFR tyrosine kinase inhibitors in EGFR-mutant lung cancer recurring after adjuvant erlotinib or gefitinib. *Clin Cancer Res* 2011;17:6322-8.
 30. Oxnard GR, Arcila ME, Chmielecki J, et al. New strategies in overcoming acquired resistance to epidermal growth factor receptor tyrosine kinase inhibitors in lung cancer. *Clin Cancer Res* 2011;17:5530-7.
 31. Pao W, Miller VA, Politi KA, et al. Acquired resistance of lung adenocarcinomas to gefitinib or erlotinib is associated with a second mutation in the EGFR kinase domain. *PLoS Med* 2005;2:e73.
 32. Arcila ME, Oxnard GR, Nafa K, et al. Rebiopsy of lung cancer patients with acquired resistance to EGFR inhibitors and enhanced detection of the T790M mutation using a locked nucleic acid-based assay. *Clin Cancer Res* 2011;17:1169-80.
 33. Kobayashi S, Boggon TJ, Dayaram T, et al. EGFR mutation and resistance of non-small-cell lung cancer to gefitinib. *N Engl J Med* 2005;352:786-92.
 34. Weickhardt AJ, Scheier B, Burke JM, et al. Local ablative therapy of oligoprogressive disease prolongs disease control by tyrosine kinase inhibitors in oncogene-addicted non-small-cell lung cancer. *J Thorac Oncol* 2012;7:1807-14.
 35. Ou SH, Jänne PA, Bartlett CH, et al. Clinical benefit of continuing ALK inhibition with crizotinib beyond initial disease progression in patients with advanced ALK-positive NSCLC. *Ann Oncol* 2014;25:415-22.
 36. Hong X, Chen Q, Ding L, et al. Clinical benefit of continuing crizotinib therapy after initial disease progression in Chinese patients with advanced ALK-rearranged non-small-cell lung cancer. *Oncotarget* 2017;8:41631-40.
 37. Lei YY, Yang JJ, Zhong WZ, et al. Clinical efficacy of crizotinib in Chinese patients with ALK-positive non-small-cell lung cancer with brain metastases. *J Thorac Dis* 2015;7:1181-8.
 38. Hotta K, Hida T, Nokihara H, et al. Final overall survival analysis from the phase III J-ALEX study of alectinib versus crizotinib in ALK inhibitor-naïve Japanese patients with ALK-positive non-small-cell lung cancer. *ESMO Open* 2022;7:100527.
 39. Riudavets M, Planchard D. An update on lorlatinib: a novel first line treatment for ALK-positive advanced lung cancer. *Expert Opin Pharmacother* 2023;24:291-9.
 40. Ou SI, Solomon BJ, Shaw AT, et al. Continuation of Lorlatinib in ALK-Positive NSCLC Beyond Progressive Disease. *J Thorac Oncol* 2022;17:568-77.
 41. Shaw AT, Solomon BJ, Besse B, et al. ALK Resistance Mutations and Efficacy of Lorlatinib in Advanced Anaplastic Lymphoma Kinase-Positive Non-Small-Cell Lung Cancer. *J Clin Oncol* 2019;37:1370-9.
 42. Cooper AJ, Sequist LV, Lin JJ. Third-generation EGFR and ALK inhibitors: mechanisms of resistance and management. *Nat Rev Clin Oncol* 2022;19:499-514.
 43. Fukuda A, Yoshida T. Treatment of advanced ALK-rearranged NSCLC following second-generation ALK-TKI failure. *Expert Rev Anticancer Ther* 2023;23:1157-67.
 44. Ou SI, Nishio M, Ahn MJ, et al. Efficacy of Brigatinib in Patients With Advanced ALK-Positive NSCLC Who Progressed on Alectinib or Ceritinib: ALK in Lung Cancer Trial of brigatinib-2 (ALTA-2). *J Thorac Oncol* 2022;17:1404-14.
 45. Hida T, Seto T, Horinouchi H, et al. Phase II study of ceritinib in alectinib-pretreated patients with anaplastic lymphoma kinase-rearranged metastatic non-small-cell lung cancer in Japan: ASCEND-9. *Cancer Sci* 2018;109:2863-72.
 46. Dipasquale A, Barigazzi C, Losurdo A, et al. Brain

- metastases and next-generation anticancer therapies: a survival guide for clinicians. *Crit Rev Oncol Hematol* 2023;194:104239.
47. Gruber I, Weidner K, Treutwein M, et al. Stereotactic radiosurgery of brain metastases: a retrospective study. *Radiat Oncol* 2023;18:202.
48. Tsai CJ, Yang JT, Shaverdian N, et al. Standard-of-care systemic therapy with or without stereotactic body radiotherapy in patients with oligoprogressive breast cancer or non-small-cell lung cancer (Consolidative Use of Radiotherapy to Block [CURB] oligoprogression): an open-label, randomised, controlled, phase 2 study. *Lancet* 2024;403:171-82.
49. Gits HC, Khosravi Flanigan MA, Kapplinger JD, et al. Sublobar Resection, Stereotactic Body Radiation Therapy, and Percutaneous Ablation Provide Comparable Outcomes for Lung Metastasis-Directed Therapy. *Chest* 2023. doi: 10.1016/j.chest.2023.12.013.
50. Siringo M, Gentile G, Caponnetto S, et al. Evaluation of Efficacy of ALK Inhibitors According to Body Mass Index in ALK Rearranged NSCLC Patients-A Retrospective Observational Study. *Cancers (Basel)* 2023;15:3422.

Cite this article as: Li Y, Hao Z, Ma Y, Setiwalidi K, Zhang Y, Zhao Y, Fu X, Liang X, Ruan Z, Tian T, Yao Y. Alectinib continuation beyond progression in *ALK*-positive non-small cell lung cancer with alectinib-refractory. *Transl Lung Cancer Res* 2024;13(1):152-162. doi: 10.21037/tlcr-23-798

Supplementary

Table S1 The original data of the study

ID	Sex	Age at diagnosis	Age at continuing alectinib	Smoking history	Histology	Combination therapy	TNM stage at diagnosis	Brain metastases at diagnosis	Bone metastases at diagnosis	New brain metastases at PD	New bone metastasis at PD	Prior exposure to chemotherapy before alectinib	Prior exposure to ALK-TKI before alectinib	ECOG Performance Status in prior alectinib	ECOG Performance Status in continuing alectinib	Best response to prior alectinib	Best response to continuing alectinib	Line of prior Alectinib	PFS of prior alectinib	PFS of continuation of alectinib	Survival months	Progression type of prior alectinib	Progression of continuing alectinib	Vital status	Research center
1	Female	17	18	Unkown	adenocarcinoma	None	Stage IV	None	None	Yes	None	None	None	1	1	PR	SD	1	18	8	26	oligoprogression	None	Alive	Nanfang Hospital, Southern Medical University
15	Female	48	48	None	adenocarcinoma	Radiation	Stage IV	None	Yes	None	None	None	None	1	1	SD	SD	1	5	21	26	oligoprogression	None	Alive	Tang Du Hospital
10	Male	37	39	None	adenocarcinoma	Radiation	Stage IV	None	Yes	Yes	None	None	None	1	1	PR	PR	1	20	15	35	CNS progression	None	Alive	The First Affiliated Hospital of University of Science and Technology of China
4	Female	36	36	None	adenocarcinoma	Radiofrequency ablation	Stage IV	None	Yes	None	None	None	None	4	3	PR	PD	1	6	5	33	oligoprogression	Yes	Alive	General Hospital for EmergenciesThe First Affiliated Hospital of Xi'an Jiaotong University
5	Male	56	57	Unkown	adenocarcinoma	Radiation	Stage IV	None	None	Yes	None	None	crizotinib	2	2	PR	SD	2	9	13	61	CNS progression	Yes	Alive	The First Affiliated Hospital of Xi'an Jiaotong University
13	Female	28	29	None	adenocarcinoma	Radiation	Stage IV	None	Yes	Yes	None	None	None	1	1	PR	PR	1	18	12	44	CNS progression	Yes	Alive	Fudan University Tumor Hospital
3	Male	39	40	Unkown	adenocarcinoma	Radiation	Stage IV	Yes	None	None	None	None	crizotinib	2	2	PR	PD	1	12	8	54	CNS progression	Yes	Alive	The First Affiliated Hospital of Xi'an Jiaotong University
6	Female	57	58	None	adenocarcinoma	Radiation	Stage IV	Yes	None	Yes	None	None	None	1	1	PR	PR	1	18	7	25	CNS progression	Yes	Death	Fudan University Tumor Hospital
9	Female	48	48	None	adenocarcinoma	Chemotherapy	Stage IV	Yes	Yes	None	None	1 cycle of chemotherapy during post-diagnostic genetic test	crizotinib	2	2	SD	SD	2	7	2	60	CNS progression	Yes	Alive	The First Affiliated Hospital of Xi'an Jiaotong University
8	Female	41	41	None	adenocarcinoma	Radiation	Stage IV	None	None	None	None	None	None	2	2	PR	SD	1	8	4	17	oligoprogression	Yes	Alive	The First Affiliated Hospital of Xi'an Jiaotong University
7	Female	43	44	None	adenocarcinoma	Radiation	Stage IV	None	None	Yes	None	None	None	2	2	PR	PR	1	11	4	27	CNS progression	Yes	Alive	The First Affiliated Hospital of Xi'an Jiaotong University
12	Female	52	54	None	adenocarcinoma	None	Stage IV	None	None	None	None	None	None	1	1	PR	SD	1	31	13	44	oligoprogression	None	Alive	Shaanxi Provincial Tumor Hospital
2	Female	28	29	None	adenocarcinoma	Radiation	Stage IV	None	Yes	None	Yes	1 cycle of chemotherapy during post-diagnostic genetic test	None	2	2	PR	PR	1	12	3	32	oligoprogression	Yes	Alive	Shaanxi Province People's Hospital
11	Male	49	51	Yes	adenocarcinoma	Radiation	Stage IV	None	Yes	None	None	None	None	1	1	PR	PR	1	27	2	32	oligoprogression	Yes	Death	Nanfang Hospital, Southern Medical University
14	Male	56	59	None	adenocarcinoma	Bevacizumab	Stage IV	Yes	Yes	None	None	None	None	1	1	PR	PR	1	19	9	52	CNS progression	Yes	Alive	Shantou City Central Hospital

TNM, Tumor-Node-Metastasis staging system; PD, progression disease; ALK-TKI, anaplastic lymphoma kinase tyrosine kinase inhibitor; ECOG, Eastern Cooperative Oncology Group; PFS, progression-free survival; PR, partial response; SD, stable disease; PD, progressive disease.