



A novel technique of oblique resection and reconstruction in a post-tracheotomy tracheoesophageal fistula: a case report

Xiaogang Liu, Qiuyuan Li, Wei Huang, Ming Liu, Haifeng Wang, Gening Jiang

Department of Thoracic Surgery, Shanghai Pulmonary Hospital, Tongji University School of Medicine, Shanghai, China

Correspondence to: Gening Jiang, MD. Department of Thoracic Surgery, Shanghai Pulmonary Hospital, Tongji University School of Medicine, No. 507 Zhengmin Rd., Shanghai, China. Email: geningjiang@tongji.edu.cn.

Abstract: Iatrogenic tracheoesophageal fistulae (TEF) following tracheostomy is unique in that the tracheal segment involved is too long to allow for a tension-free anastomosis in a traditional TEF repair. Repairing TEF according to traditional surgical methods will result in complicated surgical procedures and high surgical complications due to the high tension of anastomosis after the removal of a long section of trachea or even the inability to directly anastomosis. Difficulty in a direct end-to-end anastomosis results in a challenging situation where the procedure is complex and the morbidity is high. Here in this case, we reported a successful application of a novel technique of “oblique resection and reconstruction” for repair of a post-tracheotomy TEF. This technique made adequate use of autologous tracheal tissue for reconstruction and might implicate a reliable solution to the surgical management of TEF with minimum tension and morbidity. The operation of the patient went well, there were no short-term and long-term complications after the operation, and the follow-up recovered well. The operation method of oblique resection and reconstruction can greatly improve the utilization of normal tracheal tissue, reduce the tension of anastomosis, and reduce postoperative complications.

Keywords: Post-tracheotomy; tracheoesophageal fistula repair (TEF repair); tension-free anastomosis; tracheal reconstruction; case report

Received: 11 September 2020; Accepted: 21 October 2020; Published: 25 August 2021.

doi: 10.21037/ccts-20-156

View this article at: <http://dx.doi.org/10.21037/ccts-20-156>

Introduction

Surgical treatment of tracheoesophageal fistulae (TEF) is very challenging. Since the etiologies are not identical, the procedure needs to be adapted accordingly. In 1967, case reports from Malmejac, Toty, and Nicolas described early explorations in the etiology of post-tracheostomy TEF respectively (1-3). Typically, the formation of the fistula results from the sandwiching effect of the cuff of the tracheotomy tube anteriorly and the nasogastric tube posteriorly. Fistulae formed in this way are located on a different transverse plane from that of the tracheostomy stoma, with the stoma located superiorly, and the fistula inferiorly. By the standards of conventional procedures, the extent of resection should at least encompass both the fistula and the stoma. As a result, the remaining segments of the trachea become too short for a tension-free anastomosis.

In this circumstance, an anastomosis with excessive tension could result in anastomotic dehiscence in the short-term and tracheal malacia or stenosis in the long-term. How to develop a new surgical technique that minimizes the anastomotic tension remains a challenge and warrants ongoing investigation. We present the following article in accordance with the CARE reporting checklist (available at <https://ccts.amegroups.com/article/view/10.21037/ccts-20-156/rc>).

Case presentation

A female patient aged 41 presented comatose to a local hospital after a motor vehicle accident. The emergency workup confirmed cranial trauma with intracranial hemorrhage and multiple contusions all over the body.




Figure 1 Images of preoperative CT and endoscopies. Preoperative CT showed that the levels of the tracheotomy stoma and the TEF were 2–3 cm apart (A,B,C); preoperative bronchoscopy (D) identified the fistula posterior to the cuff; preoperative esophagogastroscopy (E) identified the fistula through which the endotracheal cuff was visible. CT, computed tomography; TEF, tracheoesophageal fistula.

After an emergency decompressive craniotomy, she was still in coma and unable to re-establish spontaneous breathing, leading to a prolonged intubation in the intensive care unit. A nasojejun tube was thus placed to allow for enteral feeding. One week after intubation, the patient was stabilized and was switched from oral ventilation to ventilation via tracheostomy. Another 1 week after tracheostomy, increased amount of gas was noted in the gastric drainage and yellowish enteric contents were found in the tracheal aspirate. Based on these findings, a TEF was suspected and later confirmed by bronchoscopy. Therefore, a decision was made to replace the original tracheotomy tube with a 7.5 mm endotracheal tube via the stoma entrance. The tube was introduced until the cuff exceeded the fistula so as to avoid further leakage. With this management, the patient slowly gained consciousness and was then referred to our hospital for further treatment.

Upon admission, a cervico-thoracic computed tomography (CT) was ordered which indicated the presence of a TEF. The location and size of the fistula was further examined through additional bronchoscopy and esophagogastroscopy (*Figure 1*). After routine preoperative workup excluded contraindications, the patient was prepared for surgery.

During the entire treatment process, the patient uses ventilator to assist breathing when unconscious, and then gradually weaned off after being awake. The purpose of

indwelling the tracheal tube is to prevent digestive juice from entering the airway through the fistula. After the operation, the tracheal intubation was pulled out in the operating room. The patient has no ventilator dependence during treatment.

During operation, a horizontal cervical incision was made at the level of the tracheotomy stoma, in addition to a routine median sternotomy over the upper sternum. During the phase of dissection, a pedicled flap of left platysma muscles was preserved for later use. The surrounding scar tissue at the stoma site was first excised for adequate exposure of the trachea. Then, the dissection of the trachea started at this point and was carried both cephalad and caudad until the whole length of the fistula was fully exposed. Since the tracheal segment affected by the fistula was too long (the length is about 5 cm), a routine tracheal segment resection and reconstruction would result in excessive tension. To avoid excessive tension over the tracheal anastomosis, an oblique segment of the trachea was resected. Two oblique cut lines parallel to each other were drawn between the tracheostomy stoma anteriorly and the fistula posteriorly. Both the stoma and the fistula were included in this oblique segment of trachea to be resected. The fistula on the esophagus was exposed by trimming tissue adjacent to the membranous trachea and the fistula. The esophageal fistula was closed with two layers of interrupted 4-0 Vicryl sutures. The left platysma muscles




Figure 2 Schematic diagrams of the oblique resection and reconstruction. An oblique resection of the tracheal segment was performed. Two parallel cut lines were drawn between and encompassed the tracheal stoma and the fistula. (A); the scar tissue was trimmed off to gain exposure of the fistula (B); end-to-end anastomosis after oblique segmental resection (C); the fistula was closed with two layers of interrupted sutures (D).

prepared beforehand was placed between the tracheal anastomosis and the fistula closure on the esophagus. After the resected segment of trachea was removed, an end-to-end anastomosis with interrupted 4-0 Vicryl sutures was made to reconstruct the trachea. For posterior sutures, the knots were tied inside the lumen, while figure-of-eight sutures were placed in the anterior wall and knots tied outside of the lumen. Specifically, two horizontal mattress sutures were added to relieve tension in the area of cartilaginous-membranous junction. Through this method of reconstructing the trachea, we actually only cut about 2 cm of tracheal segment. Schematic diagrams are shown in *Figure 2*. Before closing, two 18Fr draining tubes were placed on both sides of the anastomosis on 10 cm H₂O suction.

Two months postoperatively, the follow-up CT demonstrated an intact trachea without any signs of malacia or stenosis. The anastomosis was satisfactory under bronchoscopy (*Figure 3*). At the most recent follow-up 1 year after surgery, the patient reported good recovery.

All procedures performed in this study were in accordance with the ethical standards of the institutional and national research committees and with the Helsinki Declaration (as revised in 2013). Written informed consent

was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the editorial office of this journal.

Discussion

Iatrogenic TEF following tracheotomy have unique etiologies: the fistulae are mostly caused by sandwiching between the endotracheal cuff and the gastric tube. With the increased use of decompressive endotracheal cuff and improved awareness of the etiologies, the incidence of this type of fistula declines markedly. The conventional cuff of a tracheotomy tube features a long oval shape. By this structure, the compression is unevenly distributed and is usually aggregated at the far end of the long axis, producing a continuous high pressure that will in turn cause a focal ischemia, predisposing the affected trachea and esophagus to a future fistula. Prevention of this iatrogenic fistula can be achieved by monitoring the cuff pressure, removal of gastric tube and upgrade to a decompressive cuff. If a fistula is identified early in its course, the abovementioned prophylactic measures can also be used for conservative treatment of this disease, which may be enough for




Figure 3 Images of follow-up CT and bronchoscopy. Follow-up CT demonstrated an intact trachea without any signs of malacia or stenosis (A,B); the anastomosis was with good healing under bronchoscopy (C). CT, computed tomography.

spontaneous healing of some lesions. In other fistulae where the diagnosis is delayed and a leakage of enteric fluid is present, conservative management is often ineffective and surgery is indicated (4).

There are differences in the anatomy of esophageal fistulae secondary to tracheotomy and to intubation. The incision of tracheotomy is usually located at the frontal trachea right beneath the cricoid cartilage. This adds a tracheal defect at a different level with the fistula, with the latter often being 2–3 cm inferior, at the membranous trachea where the cuff pressure is highest. Conventional surgical techniques require that the entire length of tracheal segment from the fistula to the stoma be resected. In this way, a direct single-stage end-to-end anastomosis is difficult to accomplish due to a long segment resection, and other reconstruction methods need to be considered (5). Previously, it was reported in a pediatric tracheal resection, the technique of oblique resection and reconstruction was utilized to alleviate anastomotic tension and prevent long-term stenosis. Inspired by this, we expanded the usage of this technique in a case of post-tracheotomy TEF, in order to minimize the resection, preserve more tissue, reduce anastomotic tension, and prevent long-term malacia and stenosis. The key of this technique is drawing

two transection lines with the upper one connecting the upper rims of the stoma and the fistula, and the lower one connecting the lower rims respectively. If the stoma and the fistula are similar in length, a tension-free anastomosis can be easily achieved. However, if the fistula is longer than the stoma, adjustment could be made such that part of the membranous trachea at the fistula is sutured to allow for good approximation (6).

Because TEF is a benign condition, surgical margin is not a concern, and normal tracheal tissue could be salvaged with every effort except that the scar tissue requires radical resection. Anatomically, the tracheotomy stoma is located in the cartilaginous trachea in the anterior, and the TEF is located in the membranous trachea in the posterior. To preserve adequate tracheal tissue and minimize anastomotic tension, we re-innovated the technique of oblique resection and reconstruction which features transection along the lines with the upper one connecting the upper rims of the stoma and the fistula, and the lower one connecting the lower rims respectively.

Conclusions

We believe this technique will effectively prevent

anastomotic complications postoperatively and recommend its use in cases of post-tracheotomy iatrogenic TEF.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the Guest Editor (Tutku Soyer) for the series “Tracheoesophageal Fistula” published in *Current Challenges in Thoracic Surgery*. The article has undergone external peer review.

Reporting Checklist: The authors have completed the CARE reporting checklist. Available at <https://ccts.amegroups.com/article/view/10.21037/ccts-20-156/rc>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://ccts.amegroups.com/article/view/10.21037/ccts-20-156/coif>). The series “Tracheoesophageal Fistula” was commissioned by the editorial office without any funding or sponsorship. The authors have no other conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. All procedures performed in this study were in accordance with the ethical standards of the institutional and national research committees and with the Helsinki Declaration (as revised in 2013). Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review

by the editorial office of this journal.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Malmejac C, Arnaud A, Houel J, et al. Tracheo-esophageal fistula after tracheotomy. *Ann Chir Thorac Cardiovasc* 1967;6:1167-9.
2. Toty L, Hertzog P, Diane C, et al. 5 cases of tracheo-esophageal fistula after tracheotomy. *Ann Chir Thorac Cardiovasc* 1967;6:1155-8.
3. Nicolas F, Dupon H, Guillon J, et al. Tracheo-esophageal fistula after tracheotomy. 3 cases. *Ann Chir Thorac Cardiovasc* 1967;6:1159-65.
4. Macchiarini P, Verhoye JP, Chapelier A, et al. Evaluation and outcome of different surgical techniques for postintubation tracheoesophageal fistulas. *J Thorac Cardiovasc Surg* 2000;119:268-76.
5. Puma F, Vannucci J, Santoprete S, et al. Surgery and perioperative management for post-intubation tracheoesophageal fistula: case series analysis. *J Thorac Dis* 2017;9:278-86.
6. Sethi P, Bhatia PK, Biyani G, et al. Acquired tracheo-oesophageal fistula: a challenging complication of tracheostomy. *J Coll Physicians Surg Pak* 2015;25 Suppl 2:S76-7.

doi: 10.21037/ccts-20-156

Cite this article as: Liu X, Li Q, Huang W, Liu M, Wang H, Jiang G. A novel technique of oblique resection and reconstruction in a post-tracheotomy tracheoesophageal fistula: a case report. *Curr Chall Thorac Surg* 2021;3:31.