



Idiopathic orofacial and nociplastic pain in elderly patients: a narrative review

Noboru Noma¹, Zhimin Yan², Kentaro Urata³, Chelsia Sim⁴, Yoko Tsukada⁵, Andrew Young⁶

¹Department of Oral Medicine, Nihon University School of Dentistry, Tokyo, Japan; ²Department of Oral Medicine, Peking University School and Hospital of Stomatology, Beijing, China; ³Department of Complete Denture Prosthodontics, Nihon University School of Dentistry, Tokyo, Japan; ⁴Department of Oral Maxillofacial Surgery, Faculty of Dentistry, National University of Singapore, Singapore, Singapore; ⁵Faculty of Dentistry, University of British Columbia, Vancouver, British Columbia, Canada; ⁶Department of Diagnostic Sciences, Arthur Dugoni School of Dentistry, University of the Pacific, San Francisco, CA, USA

Contributions: (I) Conception and design: N Noma; (II) Administrative support: N Noma; (III) Provision of study materials or patients: None; (IV) Collection and assembly of data: Z Yan; (V) Data analysis and interpretation: K Urata; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Noboru Noma. 1-8-13 Kandasurugadai, Chiyoda-ku, Tokyo 101-8310, Japan. Email: noma.noboru@nihon-u.ac.jp.

Background and Objective: Idiopathic orofacial pain (IOFP) includes burning mouth syndrome (BMS), persistent idiopathic facial pain (PIFP), and persistent idiopathic dentoalveolar pain (PIDAP). We aimed to perform a narrative review on the role of nociplastic pain mechanisms in IOFP in elderly patients.

Methods: We conducted a PubMed search using only English for studies published between April 1990 and September 2021 on IOFP and nociplastic pain using the following search terms: “(Burning Mouth Syndrome OR Persistent Idiopathic Dentoalveolar Pain OR Atypical Odontalgia) AND (Temporal Summation OR Conditioned Pain Modulation OR Quantitative Sensory Testing) AND (1990/04/01:2021/9/01[Date - Entry])”.

Key Content and Findings: We identified 43 potentially relevant articles, and upon review selected 13 studies for this review; 9 were for BMS, and 4 were for PIDAP or atypical odontalgia. The IOFPs are caused by a decrease in the function of descending pain inhibition. Conditioned pain modulation (CPM) represents a natural inhibitory process on pain, which decreases with age in the orofacial region. Reduced CPM may be one of the reasons for the increased prevalence of chronic pain in older individuals. Lack of brain structures or dysregulation in the endogenous opioid system could also be a factor in the lack of the CPM effect in older individuals.

Conclusions: This review implies the idiopathic oral pain syndromes may have more nociplastic pain components than psychogenic pain. Future research is needed to further assess the relationship between nociplastic pain and psychogenic pain on the orofacial regions.

Keywords: Burning mouth syndrome (BMS); persistent idiopathic dentoalveolar pain (PIDAP); descending pain inhibition; nociplastic pain

Received: 29 September 2021; Accepted: 20 May 2022; Published online: 09 September 2022.

doi: 10.21037/fomm-21-96

View this article at: <https://dx.doi.org/10.21037/fomm-21-96>

Introduction

Idiopathic orofacial pain (IOFP) includes burning mouth syndrome (BMS), persistent idiopathic facial pain (PIFP), and persistent idiopathic dentoalveolar pain (PIDAP). The International Classification of Orofacial Pain defines BMS as

“an intraoral burning or dysesthesia sensation, recurring daily for more than two hours per day for more than three months, without clinically evident causative lesions” (1). It is usually felt in the tip of the tongue, tongue rim, and palatal mucosa. BMS is most prevalent in elderly females. PIFP is defined

as “persistent facial pain of various characteristics, recurring daily for more than two hours per day for more than three months” (1). PIDAP which has been redefined most recently (ICOP, 2020), is characterized by persistent deep, dull, or pressure-like pain in the tooth or alveolar bone in the absence of local causes. Each of the IOFPs are classified into those with and those without somatosensory abnormalities. Somatosensory abnormalities can be positive (hypersensitivity) or negative (dulled sensations) symptoms. However, unlike neuropathic pain, the region of sensory abnormalities is not limited to the area controlled by the trigeminal nerve (2).

Traditionally, pain mechanisms have been divided into “nociceptive” and “neuropathic” categories, neither of which adequately describe IOFPs. The term “nociceptive pain” was introduced by the International Association for the Study of Pain (IASP) in 2017 as a third mechanistic pain descriptor (3).

It has recently been reported that IOFP includes an element of nociceptive pain that represents a modulation in somatosensory processing, such as the dysfunction of descending pain inhibition (2). Temporal summation (TS) and conditioned pain modulation (CPM) has been observed using a novel intraepidermal electrical stimulation (IES) process in patients with BMS and in healthy adults. There is growing recognition that not only changes in peripheral neurons but also changes in central neurons may be involved in the development of PIFP, suggesting the presence of neuropathy components (4-7). Regardless of the generator of peripheral pain, PIFP has been suggested to occur due to a central phenomenon/amplification caused by processes that occur in the periphery. Similar to those shown in fibromyalgia (8), low back pain (9), and peripheral neuropathic pain (10).

Objectives

This narrative review was conducted to address the following focus question: does IOFP include an element of nociceptive pain that represents a modulation in somatosensory processing? In this review, IOFP and nociceptive pain that occur in elderly patients are discussed. We present this article in accordance with the Narrative Review reporting checklist (available at <https://fomm.amegroups.com/article/view/10.21037/fomm-21-96/rc>).

Methods

Research selection

This narrative review was conducted based on the following

focus question: Does IOFP, such as BMS or PIDAP, include an element of nociceptive pain that represents a modulation in somatosensory processing?

A literature search in PubMed was performed for studies from April 1990 to September 2021. The search keywords included “(Burning Mouth Syndrome OR Persistent Idiopathic Dentoalveolar Pain OR Persistent Idiopathic Facial Pain OR Atypical Odontalgia) AND (Temporal Summation OR Conditioned Pain Modulation OR Quantitative Sensory Testing) AND (1990/04/01:2021/9/01[Date - Entry])”.

A total of 43 studies were retrieved in PubMed. According to the question in this narrative review, 13 studies were selected; 9 on BMS and 4 on PIDAP or atypical odontalgia (Table 1).

Discussion

Nociceptive pain

The IASP introduced the concept of nociceptive pain in 2018 as “pain that arises from altered nociception despite no clear evidence of actual or threatened tissue damage causing the activation of peripheral nociceptors or evidence for disease or lesion of the somatosensory system causing the pain” (11). Therefore, nociceptive pain is a pathological condition wherein a patient has abnormally heightened sensitivity to peripheral stimuli manifesting as allodynia (a somatosensory abnormality in which non-noxious stimuli are perceived as painful) and hyperalgesia (amplification of pain stimuli) in the absence of any local pain-causing disease or disorder. These abnormal amplifications of peripheral stimuli are characteristically observed in neuropathic pain (12) but can also be observed in nociceptive pain (13); however, tissue or nerve damage in the peripheral tissues resulting in pain is observed only in neuropathic pain. When noxious stimuli do not actually originate from peripheral and nerve tissues, the central nervous system amplifies pain or recognizes it with abnormally heightened sensitivity in patients with nociceptive pain. It is believed that the pain regulation mechanism in the brain, especially descending pain inhibition, is modulated, resulting in an overreaction to non-noxious input from the periphery (13).

A previous study has shown that CPM is reduced in older individuals (14). Recently, Khan *et al.* demonstrated that CPM in both the right and left gingival sites decreases as age increases (15). Reduced CPM may be one of the reasons for the increased prevalence of chronic pain in older individuals (16). It has been reported that in the absence of the descending pain inhibitory system, noxious stimuli can

Table 1 The search strategy summary

Items	Specification
Date of search	September 1, 2021
Databases and other sources searched	PubMed
Search terms used	(Burning Mouth Syndrome OR Persistent Idiopathic Dentoalveolar Pain OR Atypical Odontalgia OR Persistent Idiopathic Facial Pain) AND (Temporal Summation OR Conditioned Pain Modulation OR Quantitative Sensory Testing)
Timeframe	(1990/04/01:2021/9/01[Date - Entry])
Inclusion and exclusion criteria	We included review articles, meta-analyses, and original studies published in English. We excluded case reports, protocols, short communications, personal opinions, letters, conference abstracts, or laboratory research
Selection process	NN and ZY independently screened the full-text articles, with disagreements being resolved through consensus with KU, YT, CS and AY

be perceived as pain (15). The lack of brain structures or dysregulation in the endogenous opioid system could also be a factor in the decrease of CPM in older individuals. The neural and hormonal pain-modulatory systems are gradually influenced by the aging process along with the loss of noradrenergic and serotonergic fibers in the spinal dorsal horn and decline of the opioid system

Nociceptive pain and BMS

When selective C-fiber stimulation is repeatedly applied to the orofacial region, neuronal excitement rapidly increases due to the input signal of the pain stimulus, and a windup phenomenon occurs in the caudal subnucleus of the trigeminal spinal tract. N-methyl-D-aspartate (NMDA) receptors, on which glutamic acid acts as an excitatory neurotransmitter, are involved in the windup phenomenon (17). Previous studies have reported that TS can be induced by applying repeated noxious thermal stimuli to the forearms of patients with BMS (18). When repeated electrical stimulation (10 stimuli at 1 Hz) was applied to the lips of patients with BMS and healthy adults, the pain intensity with each stimulus after 10 stimulations was higher than that after the first stimulation, suggesting that TS was induced. However, this excessive activation of NMDA receptors is not exclusive to the pathophysiology of BMS, as the windup phenomenon was also observed in healthy adults (19).

CPM is a phenomenon wherein pain inhibits pain. A noxious stimulus [conditioning stimulus (CS)] at a distant site inhibits wide-acting neurons in the posterior horn of the spinal cord or the caudal subnucleus of the trigeminal spinal tract via the brain stem, thereby inhibiting pain

transmission (test stimulus). It has been reported that CPM is attenuated in patients with conditions that cause chronic pain such as fibromyalgia, irritable bowel syndrome, chronic tension headaches, and migraine (20,21).

CPM of a 40 °C CS was reported to be similar in patients with BMS and healthy adults, although CPM of a 47 °C CS was significantly decreased in patients with BMS compared to that in healthy adults (19). These results suggest that the pain control mechanism works similarly in patients with BMS and in healthy adults in response to non-noxious CS but not with noxious CS (19).

Psychological factors such as anxiety and depression may explain some of the individual variability in pain perception and therefore may also play a role in CPM, possibly due to the fact that serotonin and noradrenaline are involved in both anxiety and depression are involved in the CPM response. Recently, Ozasa *et al.* found a significant positive correlation of CPM 47 °C with state and trait anxiety in patients with BMS, suggesting that both state and trait anxiety negatively affect the descending pain modulation system (22).

In patients with BMS, paresthesia increased tongue pain when hot food and drinks were consumed. Shinozaki *et al.* analyzed the brain function of patients with BMS using functional magnetic resonance imaging (fMRI) (23). When noxious heat stimuli were applied to the lips of patients with BMS, strong activity in the pain matrix, which includes the anterior cingulate gyrus, insular cortex, and prefrontal cortex, was detected. In contrast, when noxious heat stimuli were applied to the lips of healthy adults, habituation was observed. In healthy adults, the descending pain inhibition circuit suppresses activity in the cerebral cortex (anterior

cingulate gyrus). Repeated noxious stimuli resulted in pain habituation, which affected the way pain was felt. However, in patients with BMS, this habituation was not observed. This lack of descending pain inhibition may lead to diminished pain habituation (23).

In a study by Watanabe *et al.* using somatosensory testing, they revealed that elderly patients with BMS had significantly higher cold pain thresholds (CPTs) than healthy adults, and that sex hormones influenced cold and pain perception (24). The neuroprotective and nerve regeneration ability of sex hormones, and the fact that rapid changes in sex hormones during menopause damage small diameter fibers (A δ) and cause cold hyperalgesia, are thought to play a role in the modulation of CPT (25). Yilmaz *et al.* reported that A δ fibers are more susceptible to damage than C fibers in patients with BMS and may induce central sensitization (26). Cold stimulation can be used to differentiate central sensitization in patients with chronic oral pain (27).

Wada *et al.* examined resting brain fMRI images in patients with BMS and healthy adults and found that the functional connections of the prefrontal cortex and orbitofrontal area, anterior cingulate gyrus, basal ganglia, thalamus, and brain stem were strengthened in patients with BMS (28). These sites are similar to the dopaminergic nerve fiber pathways involved in descending pain modulation. Morissette *et al.* reported that the administration of 17-estradiol or progesterone to animals in which the ovaries had been extracted increased dopamine uptake in the striatum (29). When pain is induced in healthy adults, increased binding of dopamine antagonists to the D2 receptors of the caudate nucleus, putamen (30), and striatum (31) occurs according to the pain intensity. Furthermore, binding to the nucleus accumbens increases depending on emotional factors (32). However, the uptake of F-dopamine in the putamen is lower in patients with BMS than in healthy adults (33). The reactive increase of dopamine transmission in the basal ganglia in response to the stresses of pain and emotion observed in healthy patients is suppressed in patients with BMS. CPM has been reported to be significantly enhanced when dopamine agonists are subcutaneously injected compared to when placebos are used, which may help explain the decreased CPM among patients with BMS. Furthermore, the amount of dopamine in the basal ganglia is significantly reduced in patients with untreated depression compared to healthy adults (34), suggesting that emotions may affect the dopaminergic nervous system in patients with BMS.

In contrast, brain stimulation therapy, such as

electroconvulsive therapy, repetitive transcranial magnetic stimulation, and transcranial direct current stimulation, was reported to be effective for patients with primary pain syndromes who present with nociplastic pain, including those with fibromyalgia, complex regional pain syndrome, and BMS (35-38). Additionally, dopamine release in the striatum increased in brains that underwent electroconvulsive therapy (39). However, more research is needed to identify effective treatments for chronic primary pain.

Nociplastic pain and PIDAP

PIDAP was first reported in 1,778 and is characterized by persistent pain in one tooth or at a site of tooth loss (dental alveolus) with no usual dental cause (40). Due to its unclear pathophysiology, PIDAP has previously been described as idiopathic periodontalgia, phantom tooth pain, atypical odontalgia, idiopathic toothache, post-traumatic painful trigeminal neuropathy, and persistent dentoalveolar pain disorder (12).

In the 2018 International Headache Classification, 3rd Edition (2), atypical odontalgia was listed as “a subtype of persistent idiopathic facial pain (PIPF)” under classification 13.12. PIPF is defined as persistent facial and/or oral pain with a variety of symptoms and poor localization, and most patients are female. In contrast, atypical odontalgia is clearly localized, has an early age of onset, and has no significant prevalence difference between sexes. In addition, PIPF is listed in category 6.2 of the International Orofacial Pain Classification, 1st Edition in 2020 (11), while pathological conditions previously referred to as atypical odontalgia are listed as PIDAP in category 6.3. As such, the two diseases are classified separately.

The pathophysiology of IOFP may include central sensitization and modulation of pain processing in the brain. Studies using brain functional imaging suggest that IOFP may be amplified during central pain processing, similar to the amplification of pain in fibromyalgia and irritable bowel syndrome (41). Woda *et al.* suggested that higher pain centers are involved in IOFP, and that the addition of emotional stress and peripheral stimuli to biochemically and psychologically vulnerable patients lead to increased neuropeptide levels in the target tissues, leading to disease development (42). Some patients with IOFP may present with paresthesia with warm or cold stimuli, pin pricks, or mild tactile stimuli on clinical somatosensory exams. IOFP can also be accompanied by modulation of

the somatosensory nervous system and may be related to modulation of the descending inhibitory system and nociplastic pain (2). Nasri-Heir *et al.* reported that CPM in patients with idiopathic persistent odontalgia was decreased compared to that in healthy adults, and that CPM was significantly attenuated in patients with brain disease of one year or longer (43). In addition, a relationship between CPM and post-root canal treatment has been reported (44). Patients with endodontic diseases underwent CPM evaluation prior to root canal treatment, and patients with low CPM had a high risk of developing chronic orofacial pain, and patients with higher CPM before treatment had less chronic pain 6 months postoperatively. These results suggest that CPM can be used to evaluate the endogenous analgesic system in patients with nociplastic pain (44).

This review has a limitation. Compared to PIFP, there were less studies on sensory changes, CPM, or MRI related to PIDAP in this region. The few studies on PIDAP may be attributed to the fact that PIDAP has a component of neuropathic pain as it is more common after receiving root canal treatment. Further research will be needed to elucidate the pathophysiology of PIDAP.

Summary

The idiopathic oral pain syndromes that were previously considered to be psychogenic pain are now considered as nociplastic pain components that are part of the chronic primary pain syndrome. Although the exact cause of pain is unknown, there is central dysregulation in information processing of peripheral stimuli in nociplastic pain. In particular, descending pain inhibition is impaired, resulting in pain amplification. Dysfunction of the dopaminergic nervous system plays a key role in nociplastic pain in the oral region.

It is important to recognize that nociplastic pain, unlike nociceptive pain and neuropathic pain, may be amplified by central nervous system conditions, such as fatigue, inadequate sleep, memory, mood anxiety and depression problems. Consistent with this central component, this type of pain does not typically respond to peripherally-directed therapies such as anti-inflammatory drugs and opioids, surgery, or injections.

Acknowledgments

Funding: This work was supported in part by research grants from the Sato and Uemura funds, the Dental

Research Center of Nihon University School of Dentistry, and KAKENHI (Grant-in-Aid for Scientific Research [C] [20K10171]).

Footnote

Provenance and Peer Review: This article was commissioned by the Guest Editors (Junad Khan and Davis Thomas) for the series “Orofacial Pain” published in *Frontiers of Oral and Maxillofacial Medicine*. The article has undergone external peer review.

Reporting Checklist: The authors have completed the Narrative Review reporting checklist. Available at <https://fomm.amegroups.com/article/view/10.21037/fomm-21-96/rc>

Peer Review File: Available at <https://fomm.amegroups.com/article/view/10.21037/fomm-21-96/prf>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://fomm.amegroups.com/article/view/10.21037/fomm-21-96/coif>). The series “Orofacial Pain” was commissioned by the editorial office without any funding or sponsorship. The authors have no other conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. International Classification of Orofacial Pain, 1st edition (ICOP). Cephalalgia 2020;40:129-221.
2. Headache Classification Committee of the International

- Headache Society (IHS) The International Classification of Headache Disorders, 3rd edition. *Cephalalgia* 2018;38:1-211.
3. Kosek E, Cohen M, Baron R, et al. Do we need a third mechanistic descriptor for chronic pain states? *Pain* 2016;157:1382-6.
 4. Woda A. Pain in the trigeminal system: from orofacial nociception to neural network modeling. *J Dent Res* 2003;82:764-8.
 5. Woda A, Pionchon P. A unified concept of idiopathic orofacial pain: pathophysiological features. *J Orofac Pain* 2000;14:196-212.
 6. Woda A. Neuropathic pain. *J Orofac Pain* 2007;22:257-8.
 7. Woda A. A "dysfunctional" pain group in addition to the "neuropathic" and "nociception/inflammatory" groups of orofacial pain entities? *J Orofac Pain* 2009;23:89-90.
 8. Mhalla A, de Andrade DC, Baudic S, et al. Alteration of cortical excitability in patients with fibromyalgia. *Pain* 2010;149:495-500.
 9. Hashmi JA, Baliki MN, Huang L, et al. Shape shifting pain: chronification of back pain shifts brain representation from nociceptive to emotional circuits. *Brain* 2013;136:2751-68.
 10. Lefaucheur JP. The use of repetitive transcranial magnetic stimulation (rTMS) in chronic neuropathic pain. *Neurophysiol Clin* 2006;36:117-24.
 11. Treede RD, Rief W, Barke A, et al. A classification of chronic pain for ICD-11. *Pain* 2015;156:1003-7.
 12. Finnerup NB, Haroutounian S, Kamerman P, et al. Neuropathic pain: an updated grading system for research and clinical practice. *Pain* 2016;157:1599-606.
 13. Trouvin AP, Perrot S. New concepts of pain. *Best Pract Res Clin Rheumatol* 2019;33:101415.
 14. Naugle KM, Ohlman T, Naugle KE, et al. Physical activity behavior predicts endogenous pain modulation in older adults. *Pain* 2017;158:383-90.
 15. Khan J, Korczeniewska O, Benoliel R, et al. Age and gender differences in mechanically induced intraoral temporal summation and conditioned pain modulation in healthy subjects. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2018;126:134-41.
 16. Johannes CB, Le TK, Zhou X, et al. The prevalence of chronic pain in United States adults: results of an Internet-based survey. *J Pain* 2010;11:1230-9.
 17. Price DD, Hu JW, Dubner R, et al. Peripheral suppression of first pain and central summation of second pain evoked by noxious heat pulses. *Pain* 1977;3:57-68.
 18. Nasri-Heir C, Shigdar D, Alnaas D, et al. Primary burning mouth syndrome: Literature review and preliminary findings suggesting possible association with pain modulation. *Quintessence Int* 2017;49:49-60.
 19. Nishihara C, Watanabe K, Ozasa K, et al. Altered pain modulation to noxious heat thermal stimuli in burning mouth syndrome. *Oral Dis* 2020;26:1777-82.
 20. Pielsticker A, Haag G, Zaudig M, et al. Impairment of pain inhibition in chronic tension-type headache. *Pain* 2005;118:215-23.
 21. Price DD, Staud R, Robinson ME, et al. Enhanced temporal summation of second pain and its central modulation in fibromyalgia patients. *Pain* 2002;99:49-59.
 22. Ozasa K, Noma N, Kobayashi M, et al. Association Between Anxiety and Descending Pain Modulation of Thermal Stimuli in Patients with Burning Mouth Syndrome: A Cross-Sectional Study. *J Oral Facial Pain Headache* 2022;36:67-77.
 23. Shinozaki T, Imamura Y, Kohashi R, et al. Spatial and Temporal Brain Responses to Noxious Heat Thermal Stimuli in Burning Mouth Syndrome. *J Dent Res* 2016;95:1138-46.
 24. Watanabe K, Noma N, Sekine N, et al. Association of somatosensory dysfunction with symptom duration in burning mouth syndrome. *Clin Oral Investig* 2019;23:3471-7.
 25. Jääskeläinen SK. Is burning mouth syndrome a neuropathic pain condition? *Pain* 2018;159:610-3.
 26. Yilmaz Z, Egbuniwe O, Renton T. The Detection of Small-Fiber Neuropathies in Burning Mouth Syndrome and Iatrogenic Lingual Nerve Injuries: Use of Quantitative Sensory Testing. *J Oral Facial Pain Headache* 2016;30:87-98.
 27. Zagury JG, Eliav E, Heir GM, et al. Prolonged gingival cold allodynia: a novel finding in patients with atypical odontalgia. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2011;111:312-9.
 28. Wada A, Shizukuishi T, Kikuta J, et al. Altered structural connectivity of pain-related brain network in burning mouth syndrome-investigation by graph analysis of probabilistic tractography. *Neuroradiology* 2017;59:525-32.
 29. Morissette M, Di Paolo T. Effect of chronic estradiol and progesterone treatments of ovariectomized rats on brain dopamine uptake sites. *J Neurochem* 1993;60:1876-83.
 30. Scott DJ, Heitzeg MM, Koeppe RA, et al. Variations in the human pain stress experience mediated by ventral and dorsal basal ganglia dopamine activity. *J Neurosci* 2006;26:10789-95.
 31. Hagelberg N, Jääskeläinen SK, Martikainen IK, et al.

- Striatal dopamine D2 receptors in modulation of pain in humans: a review. *Eur J Pharmacol* 2004;500:187-92.
32. Jääskeläinen SK, Rinne JO, Forssell H, et al. Role of the dopaminergic system in chronic pain -- a fluorodopa-PET study. *Pain* 2001;90:257-60.
 33. Anand A, Barkay G, Dziedzic M, et al. Striatal dopamine transporter availability in unmedicated bipolar disorder. *Bipolar Disord* 2011;13:406-13.
 34. Umezaki Y, Badran BW, DeVries WH, et al. The Efficacy of Daily Prefrontal Repetitive Transcranial Magnetic Stimulation (rTMS) for Burning Mouth Syndrome (BMS): A Randomized Controlled Single-blind Study. *Brain Stimul* 2016;9:234-42.
 35. Umezaki Y, Badran BW, Gonzales TS, et al. Daily left prefrontal repetitive transcranial magnetic stimulation for medication-resistant burning mouth syndrome. *Int J Oral Maxillofac Surg* 2015;44:1048-51.
 36. Villamar MF, Wivatvongvana P, Patumanond J, et al. Focal modulation of the primary motor cortex in fibromyalgia using 4x1-ring high-definition transcranial direct current stimulation (HD-tDCS): immediate and delayed analgesic effects of cathodal and anodal stimulation. *J Pain* 2013;14:371-83.
 37. Yang S, Chang MC. Transcranial Direct Current Stimulation for the Management of Neuropathic Pain: A Narrative Review. *Pain Physician* 2021;24:E771-81.
 38. Wolanin MW, Gulevski V, Schwartzman RJ. Treatment of CRPS with ECT. *Pain Physician* 2007;10:573-8.
 39. Strafella AP, Paus T, Fraraccio M, et al. Striatal dopamine release induced by repetitive transcranial magnetic stimulation of the human motor cortex. *Brain* 2003;126:2609-15.
 40. Malacarne A, Spierings ELH, Lu C, et al. Persistent Dentoalveolar Pain Disorder: A Comprehensive Review. *J Endod* 2018;44:206-11.
 41. Giesecke T, Gracely RH, Williams DA, et al. The relationship between depression, clinical pain, and experimental pain in a chronic pain cohort. *Arthritis Rheum* 2005;52:1577-84.
 42. Woda A, Pionchon P. A unified concept of idiopathic orofacial pain: clinical features. *J Orofac Pain* 1999;13:172-84; discussion 185-95.
 43. Nasri-Heir C, Khan J, Benoliel R, et al. Altered pain modulation in patients with persistent postendodontic pain. *Pain* 2015;156:2032-41.
 44. Khan J, Zusman T, Wang Q, et al. Acute and Chronic Pain in Orofacial Trauma Patients. *J Endod* 2019;45:S28-38.

doi: 10.21037/fomm-21-96

Cite this article as: Noma N, Yan Z, Urata K, Sim C, Tsukada Y, Young A. Idiopathic orofacial and nociplastic pain in elderly patients: a narrative review. *Front Oral Maxillofac Med* 2024;6:4.